HEDIS[®] 2024 Documentation and Coding Guidelines

ding Guidelines



EFFECTIVENESS OF CARE: PREVENTION AND SCREENING3
Care for Older Adults (COA)
Cervical Cancer Screening (CCS) 6
Childhood Immunization Status (CIS)8
Chlamydia Screening in Women (CHL)11
Immunizations for Adolescents (IMA) 12
Lead Screening Children (LSC) 14
Oral Evaluation, Dental Services (OED)15
Topical Fluoride for Children (TFC)15
Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (WCC)16
EFFECTIVENESS OF CARE: RESPIRATORY CONDITIONS
Appropriate Testing for Pharyngitis (CWP)19
Asthma Medication Ratio (AMR 20
Medication Management for People with Asthma (MMA) 22
Pharmacotherapy Management of COPD Exacerbation (PCE)
ACCESS AND AVAILABILITY
Adults' Access to Preventive/Ambulatory Health Services (AAP)
Children and Adolescents' Access to Primary Care (CAP) 24
Initiation and Engagement of Substance Use Disorder Treatment (IET) 24

	Prenatal and Postpartum Care (PPC) 27	7
	Use of First-Line Psychosocial Care for Children and Adolescents on Antipsychotics (APP)	C
E	FFECTIVENESS OF CARE: CARDIOVASCULAR CONDITIONS	1
	Controlling High Blood Pressure (CBP)	1
	Persistence of Beta Blocker Treatment After a Heart Attack (PBH) 33	3
	Cardiac Rehabilitation (CRE)	4
	Statin Therapy for Patients with Cardiovascular Disease (SPC)	5
E	FFECTIVENESS OF CARE: DIABETES	B
	Glycemic Status Assessment for Patients With Diabetes (GSD)	3
	Comprehensive Diabetes Care (CDC) Monitoring for Nephropathy 40	C
	Eye Exam for Patients with Diabetes (EED)	C
	Blood Pressure Control for Patients with Diabetes (BPD) 43	3
	Kidney Evaluation for Patients With Diabetes (KED) 44	4
	Statin Therapy for Patients with Diabetes (SPD) 45	5
E	FFECTIVENESS OF CARE: MUSCULOSKELETAL CONDITIONS	6
	Disease-Modifying Anti-Rheumatic Drug Therapy for Rheumatoid Arthritis (ART)	
	Osteoporosis Management in Women Who Had a Fracture (OMW) 47	7
	Osteoporosis Screening in Older Women (OSW)	3

El	FFECTIVENESS OF CARE: BEHAVIORAL HEALTH	.49
	Antidepressant Medication Management (AMM)	49
	Follow-Up After Hospitalization for Mental Illness (FUH)	50
	Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who are Using Antipsychotic Medications (SSD)	. 59
	Diabetes Monitoring for People with Diabetes and Schizophrenia (SMD)59
	Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC)	. 60
	Adherence to Antipsychotic Medications for Individuals with Schizophrenia (SAA)	. 60
	Follow-Up After Emergency Department Visit for Mental Illness (FUM) .	61
	Follow-Up After Emergency Department Visit for Substance Use (FUA) .	71
	Follow-Up After High-Intensity Care for Substance Use Disorder (FUI)	75
	Pharmacotherapy for Opioid Use Disorder (POD)	78
EI	FFECTIVENESS OF CARE: CARE COORDINATION	.79
	Advance Care Planning (ACP)	79
	Transition of Care (TRC)	80
	Follow-Up After Emergency Department Visit for People With Multiple High-Risk Chronic Conditions (FMC)	
EI	FFECTIVENESS OF CARE: OVERUSE/APPROPRIATENESS	.91
	Avoidance of Antibiotic Treatment for Acute Bronchitis (AAB)	91
	Non-Recommended PSA-Based Screening in Older Men (PSA)	92
	Appropriate Treatment for Upper Respiratory Infection (URI)	92
	Potentially Harmful Drug-Disease Interactions in Older Adults (DDE)	94
	Risk of Continued Opioid Use (COU)	96
	Use of High-Risk Medication in Older Adults (DAE)	97
w	ww.amerihealthcaritasoh.com	

Deprescribing of Benzodiazepines in Older Adults (DBO)	19
Use of Imaging Studies for Low Back Pain (LBP)	0
Use of Opioids at High Dosage (HDO)10)1
Use of Opioids From Multiple Providers (UOP)	13
UTILIZATION	4
Well-Child Visits in the First 30 Months of Life (W30) 10)4
Child and Adolescent Well-Care Visits (WCV))5
Annual Dental Visit (ADV)10)6
MEASURES COLLECTED USING ELECTRONIC CLINICAL DATA SYSTEMS 10	6
Follow-Up Care for Children Prescribed ADHD Medication (ADD-E) 10)6
Adult Immunization Status (AIS-E)	8
Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM-E))9
Breast Cancer Screening (BCS-E)11	0
Colorectal Cancer Screening (COL-E)11	.1
Depression Screening and Follow-Up for Adolescents and Adults (DSF-E)11	.3
Utilization of the PHQ-9 to Monitor Depression Symptoms for Adolescents and Adults (DMS-E)11	.4
Depression Remission or Response for Adolescents and Adults (DRR-E)	.6
Unhealthy Alcohol Use Screening and Follow-Up (ASF-E)	.7
Prenatal Immunization Status (PRS-E) 11	.8
Prenatal Depression Screening and Follow-Up (PND-E)	.9
Postpartum Depression Screening and Follow-Up (PDS-E)	0
Social Need Screening and Intervention (SNS-E)	2

2

HEDIS[®] 2024 Documentation and Coding Guidelines





Reproduced with permission from HEDIS Measurement Year (MY) 2024, Volume 2: Technical Specifications for Health Plans by the National Committee for Quality Assurance (NCQA). HEDIS[®] is a registered trademark of the National Committee for Quality Assurance (NCQA). To purchase copies of this publication, contact NCQA Customer Support at 888-275-7585 or visit <u>https://store.ncqa.org/.</u>

EFFECTIVENESS OF CARE: PREVENTION AND SCREENING

Measure	Measure Description	Measure Information/Documentation Required	Coding
Care for Older Adults (COA)	 The percentage of adults 66 years of age and older who had each of the following during the measurement year (MY): Medication review. Functional Status Assessment (FSA). Pain Assessment. 	 Required Medication Review: A review conducted by a prescribing practitioner or clinical pharmacist in the MY. The member does not need to be present for the medication review. Any of the following are acceptable: The presence of a medication list in the medical record with notation of the date reviewed. Dated notation that member is not taking any medications. Transitional care management services documented during the MY. Criteria is not met if review performed by an RN. Functional Status Assessment: At least one functional status assessment during the MY and the date it was performed. Functional status assessment must include one 	Functional Status Assessment: CPT: 99483 CPT-CAT-II: 1170F Pain Assessment: CPT-CAT-II: 1125F, 1126F Medication Review (with Medication List): CPT: 90863, 99483 CPT-CAT-II: 1160F Medication List (with Medication Review): CPT-CAT-II: 1159F Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

 Notation that Activities of Daily Living (ADLs) were assessed or that at least five of the following
were assessed: bathing, dressing,
eating, transferring (e.g., getting
in and out of chairs), using toilet,
walking. Notation that Instrumental
Activities of Daily Living (IADLs)
were assessed or at least four of
the following were assessed:
shopping for groceries, driving or
using public transportation, using the telephone, cooking or meal
preparation, housework, home
repair, laundry, taking
medications, handling finances.
Result of an assessment using a
standardized functional status assessment tool.
Criteria is not met by a fall
assessment.
Pain Assessment:
At least one pain assessment during
the MY and the date it was
performed.Documentation that the patient
was assessed for pain (which may
include positive or negative
findings for pain).
Result of assessment using a
standardized pain assessment tool.
 Criteria is not met by notation of
only a pain management plan or
only a pain treatment plan.
Criteria is not met by notation of
only screening for chest pain or

only documentation of chest pain.

Note:

- Telephone, e-visit, or virtual check-in visits are acceptable for FSA and Pain Assessment.
- Exclude services provided in an acute inpatient setting.

Required Exclusions:

Members who meet any of the following criteria are excluded from the measure:

- In hospice or using hospice services any time in the MY.
- Deceased at any time in the MY.

Common Chart Deficiencies:

- Medication Review: Medication review completed by RN.
- FSA: Documentation referencing patient living alone but not specifically that patient can perform ADLs or IADLs.
- FSA: Documentation of "normal" under review of systems without specifically addressing ADLs/IADLs.
- FSA: A functional status assessment limited to an acute or single condition, event, or body system.
- Pain: Patient not assessed for pain at visit.
- Pain: Diagnosis or medication related to pain or pain management plan but no

		documentation of pain assessment.	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Cervical Cancer Screening (CCS) This is also a measure (CCS-E) collected through claims and Electronic Clinical Data Systems. Please discuss options for a direct data feed with your Account Executive. Direct data feeds can improve provider quality performance and reduce the burden of medical record requests.	 The percentage of members 24 – 64 years of age in the MY who were recommended for routine cervical cancer screening using the following criteria: Ages 24 – 64: A cervical cytology (Pap) test within the last 3 years. Ages 30 – 64: A cervical high-risk human papillomavirus (hrHPV) test performed within the last 5 years. Ages 30 – 64: A cervical cytology (Pap test/high-risk human papillomavirus [hrHPV]) co-testing within the last 5 years. 	 Documentation using either of the following criteria meet: A note indicating the date when the cervical cytology was performed and the findings. A note indicating the date when the hrHPV test was performed and the findings. Note: Evidence of hrHPV testing within the last 5 years also captures patients who had cotesting. Do NOT Count: Lab results that indicate results "Unknown." Lab results that indicate the sample was inadequate or that "no cervical cells were present" is not a valid screening. Biopsies are diagnostic and are not valid as a primary cervical cancer screening. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. 	Cervical Cytology (Pap): CPT: 88141, 88142, 88143, 88147, 88148, 88150, 88152, 88153, 88164, 88165, 88166, 88167, 88174, 88175 HCPCS: G0123, G0143, G0144, G0145, G0147, G0148, Q0091 High-Risk HPV Testing: CPT: 87624, 87625 HCPCS: G0476 Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

 Receiving palliative care any time in the MY. Evidence of a hysterectomy with no residual cervix. Must specify "complete," "total," "radical," "abdominal," or "vaginal" hysterectomy. "Cervical agenesis" or "acquired absence of the cervix." Hysterectomy in combination with documentation that the patient no longer needs Pap testing/cervical cancer screening.
 Gender Exclusions: Evidence that a patient was born a male. Members with Male sex assigned at Birth. Documentation patient is "transitioning from male to female" or has undergone sex reassignment surgery from male to female. Documentation of "binary,"' "non-binary," "transgender," or "transsexual" would not be considered an exclusion.
 Common Chart Deficiencies: Unclear if member's cervix is absent. Hysterectomy is not documented in the chart sufficiently to exclude member from measure. Member-reported data not documented with sufficient information to show the

Childhood Immunization Status (CIS)The percentage of members 2 years of age in the MY who are up to date on recommended ucine administration services on the same diptheria, tetanus, and acellular pertussis (DTaP): polio (IPV); measles, mumps, and rubella (MMR);Children 2 years of age who had the following:Use applicable vaccination code or diagnosis indicating history of disease.0When coding E&M and vaccine administration services on the same diptheria, tetanus, and acellular pertussis (DTaP): polio (IPV); measles, mumps, and rubella (MMR); Haemophilus influenza type B (HiB): hepatitis B (CFE- collected through <i>Electronic Clinical Data</i> (VZV); pneumococcal conjugate (PCV); hepatitis A (HepA); rotavirus (RV); and influenza (Flu).Children 2 years of age who had the following:Use applicable vaccination code or diagnosis indicating history of disease.0When coding E&M and vaccine administration services on the £80 (DTaP): polio (IPV); measles, mumps, and rubella (MMR); Haemophilus influenza type B (HiB); hepatitis B (VZV); pneumococcal conjugate (PCV); hepatitis A (HepA); rotavirus (RV); and influenza (Flu).Children 2 years of age who had the following:Use applicable vaccination code or diagnosis indicating history of disease.Children 2 years of age influenza type B (MIMR); Haemophilus influenza type B (HiB); rotavirus (RV); and influenza (Flu).Children 2 years of age who had the rotavirus (RV); and influenza (Flu).Children 2 years of age who had the rotavirus (RV); and influenza (Flu).Children 2 years of age who had the rotavirus (RV); and influenza (Flu).Children 2 years of age who had the rotavirus (RV); and 	Measure	Measure Description	 screening was completed with a result in the measure time frame. Pap/HPV test completed but results not documented. Missing clear documentation on transgender patients (not clear that member is appropriate for the screening or if screening ordered/completed). Measure Information/Documentation Required 	Coding
	Status (CIS) When coding E&M and vaccine administration services on the same date, you must append modifier 25 to the E&M code effective 1/1/14. This is also a measure (CIS-E) collected through Electronic Clinical Data Systems. Please discuss options for a direct data feed with your Account Executive. Direct data feeds can improve provider quality performance and reduce the burden of medical	members 2 years of age in the MY who are up to date on recommended routine vaccines for diphtheria, tetanus, and acellular pertussis (DTaP); polio (IPV); measles, mumps, and rubella (MMR); Haemophilus influenza type B (HiB); hepatitis B (HepB); chicken pox (VZV); pneumococcal conjugate (PCV); hepatitis A (HepA); rotavirus (RV); and	 following: 1 MMR on or between the 1st and 2nd birthdays or history of measles, mumps, and rubella on or before the 2nd birthday. 1 VZV on or between the 1st and 2nd birthdays, history of chicken pox, or anaphylaxis due to the VZV vaccine on or before the 2nd birthday. 1 HepA on or between the 1st and 2nd birthdays, history of hepatitis A, or anaphylaxis due to the vaccine on or before the 2nd birthday. 3 HepB with different date of service on or before the 2nd birthday or history of the illness or anaphylaxis due to the vaccine. One of the 3 can be newborn (DOB to 7 days after birth). 3 IPV with different DOS on or before the 2nd birthday. Do not 	Diphtheria and Tetanus Toxoids and Acellular Pertussis vaccine (DTaP): CVX: 20, 50, 106, 107, 110, 120, 146 CPT: 90697, 90698, 90700, 90723 Haemophilus Influenza Type B (HiB): CVX: 17, 46, 47, 48, 49, 50, 51, 120, 146, 148 CPT: 90644, 90647, 90648, 90697, 90698, 90748 Hepatitis A Vaccine (HepA): CVX: 31, 83, 85 CPT: 90633 History of Hepatitis A: ICD10CM: B15.0, B15.9 Hepatitis B Vaccine (HepB): CVX: 08, 44, 45, 51, 110, 146 CPT: 90697, 90723, 90740, 90744, 90747, 90748 HCPCS: G0010 Hepatitis B Newborn Vaccine: ICD10PCS: 3E0234Z

 3 Hib with different DOS on or before the 2nd birthday or anaphylaxis due to the HiB vaccine. Do not count DOS prior to 42 days after birth. 4 PCV with different DOS or anaphylaxis due to the vaccine on or before the 2nd birthday. Do not count DOS prior to 42 days after birth. 4 DTaP different DOS on or before the 2nd birthday or anaphylaxis or encephalitis due to any of the vaccines. Do not count DOS prior to 42 days after birth. 2 or 3 RV on different DOS or anaphylaxis due to the vaccine on 	Inactivated Poliovirus Vaccine (IPV): CVX: 10, 89, 110, 120, 146 CPT: 90697, 90698, 90713, 90723 Influenza Vaccine: CVX: 88, 140, 141, 150, 153, 155, 158, 161, 171, 186 CPT: 90655, 90657, 90661, 90673, 90674, 90685, 90686, 90687, 90688, 90756 HCPCS: G0008 LAIV Immunization: CVX: 111, 149 CPT: 90660, 90672 Measles, Mumps, and Rubella Vaccine (MMR): CVX: 03, 94 CPT: 90707, 90710
or before the 2 nd birthday. Do not count DOS prior to 42 days after	History of Measles: ICD10CM: B05.0, B05.1, B05.2, B05.3, B05.4, B05.81, B05.89, B05.9
 birth. 2 Flu with different DOS or anaphylaxis due to the vaccine on or before 2nd birthday. Do not 	History of Mumps: ICD10CM: B26.0, B26.1, B26.2, B26.3, B26.81, B26.82, B26.83, B26.84, B26.85, B26.89, B26.9
count DOS prior to 6 months (180 days) after birth. One of the two vaccinations can be LAIV	History of Rubella: ICD10CM: B06.00, B06.01, B06.02, B06.09, B06.81, B06.82, B06.89, B06.9
administered ONLY on the 2 nd birthday.	Pneumococcal Conjugate Vaccine (PCV):
Documentation:	CVX: 109, 133, 152, 215 CPT: 90670, 90671
 A note indicating the name of the 	HCPS: G0009
specific antigen and the date of	
the immunization.	Rotavirus Vaccine (RV):
A certificate of immunization	CVX: 116, 122 (3 dose)
prepared by an authorized health	CPT: 90680 (3 dose), 90681 (2 dose)
care provider or agency, including	
the specific dates and types of	Varicella Zoster Virus (VZV):
immunizations administered.	CVX: 21, 94

 "nursery/hosp documented i record or indi immunization appropriate. Immunization using a generi vaccine) or "II 	record as B02.21, B02.22, B02.23, B02.24, B02.29, B02.30, B02.31, B02.32, B02.33, B02.34, B02.39, B02.7, B02.8, B02.9 s documented ic header (e.g., polio PV/OPV" can be
counted as ev	idence of IPV. Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
Required Exclusio	
Members who me	
following criteria a	
the measure:	
In hospice or	using hospice
services any t	ime in the MY.
Deceased at a	ny time in the MY.
	lowing on or before
the child's 2 nd	
	re combined
	unodeficiency.
	unodeficiency.
o HIV.	
	phoreticular cancer,
leuke	iple myeloma, or
o Intus	susception
Common Chart De	eficiencies:
Immunization	s administered
after the 2nd	birthday.
PCP charts do	not contain
immunization	
	eived elsewhere,
	given at health
	or those given in
the hospital a	t birth.

		 Rotavirus documentation does not specify if 2-dose or 3-dose. Flu Mist only meets criteria when administered on the 2nd birthday. A note that "member is up to date" with all immunizations does not constitute compliance due to insufficient data. Parental refusal does not meet compliance. 	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Chlamydia Screening in Women (CHL)	The percentage of women 16 – 24 years of age who were identified as sexually active and who had at least one test for chlamydia during the MY.	 Perform chlamydia screening each year on every 16- to 24-year-old female identified as sexually active. Chlamydia screening can be performed through a urine test. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. A pregnancy test in the MY and a prescription for isotretinoin (Retinoid) on the date of the pregnancy test. A pregnancy test in the MY and an X-ray on the date of the pregnancy test or the 6 days after the pregnancy test. 	Chlamydia Tests: CPT: 87110, 87270, 87320, 87490, 87491, 87492, 87810 Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

Immunizations for Adolescents (IMA)The percentage of adolescents 13 years of age who had the following: up to date on recommended routine vaccines for meningcoccal; tetanus, diptheria toxoids, and a cellular pertussis (IMA-E) collected through climics of feed with your Account Executive.The percentage of adolescents 13 years of age who had the following: the dolescents 13 years of age who had the following: or a between the 11 th and 13 th birthdays. or evidence of antigen, or a between the 13 th birthday.Meningcoccal Vaccine: CVX: 32, 108, 114, 136, 147, 167, 203 CVX: 32, 108, 0733, 90734When coding E&M and vaccines administration services on the same date, you must append modifier 25 to the E&M.The percentage of a dolescents 13 years of age who had the following: on or before the 13 th birthday.Meningcoccal Vaccine: CVX: 32, 108, 114, 136, 147, 167, 203 CVX: 32, 108, 114, 136, 147, 167, 203 CVX: 115This is also a measure (IMA-E) collected through climics of feed with your Account Executive. Direct data feed with your Account Executive. Direct data feed with your Account Executive. Direct data feed son in performance and reduce the burden of medical record requests.The percentage of adolescents (IMA) Adolescent 13 the inthday. O 2 doses with at least 146 days between the 9 th and 13 th birthdays. O 2 doses with at least 146 days between the 9 th and 13 th birthdays. O 2 doses with at least 146 days between the 9 th and 13 th birthdays. O 2 doses with at least 146 days between the 9 th and 13 th birthdays. O Direct data feed of medical record requests.Note: LOINC and SNOMED codes can be captured through electronic data data of ado son	Measure	Measure Description	 Common Chart Deficiencies: Not collecting/testing urine sample routinely at well-visits. Criteria is not met by notation of parental/patient refusal. Criteria is not met by notation that patient is not sexually active. Measure Information/Documentation Required 	Coding
Documentation:	Adolescents (IMA) When coding E&M and vaccine administration services on the same date, you must append modifier 25 to the E&M. This is also a measure (IMA-E) collected through claims and Electronic Clinical Data Systems. Please discuss options for a direct data feed with your Account Executive. Direct data feeds can improve provider quality performance and reduce the burden of medical	adolescents 13 years of age in the MY who are up to date on recommended routine vaccines for meningococcal; tetanus, diphtheria toxoids, and acellular pertussis (Tdap); and human	 Adolescents 13 years of age who had the following: Meningococcal MCV with DOS on or between the 11th and 13th birthdays or evidence of antigen or anaphylaxis due to the vaccine on or before the 13th birthday. Tdap or TD with DOS on or between the 10th and 13th birthdays or evidence of antigen, anaphylaxis, or encephalitis due to the vaccine on or before the 13th birthday. HPV — any of the following: 3 doses with different dates of service on or between the 9th and 13th birthdays. 2 doses with at least 146 days between the 1st and 2nd dose on or between the 9th and 13th birthdays. Anaphylaxis due to the vaccine on or before the 13th birthdays. 	CVX: 32, 108, 114, 136, 147, 167, 203 CPT: 90619, 90733, 90734 Tetanus, Diphtheria, & Acellular Pertussis Vaccine (Tdap): CVX: 115 CPT: 90715 HPV Vaccine: CVX: 62, 118, 137, 165 CPT: 90649, 90650, 90651 Note: LOINC and SNOMED codes can be captured through electronic data

A note indicating the name of the
specific antigen and the date of
the immunization.
A certificate of immunization
prepared by an authorized health
care provider or agency including
the specific dates and types of
immunizations administered.
Required Exclusions:
Members who meet any of the
following criteria are excluded from
the measure:
In hospice or using hospice
services any time in the MY.
 Deceased at any time in the MY.
- Deceased at any time in the Wit.
Common Chart Deficiencies:
Immunizations administered
outside of the appropriate time
frames.
PCP charts do not contain records
when immunizations
administered elsewhere (i.e.,
health departments, school
clinics, urgent care facilities).
 HPV doses are not at least 146
days apart when only 2 doses
administered.
 A note that "member is up to
date" with all immunizations
does not constitute compliance
due to insufficient data.
Parental refusal does not meet
compliance.
Td (Tetanus, Diphtheria Toxoids)
does not meet criteria for Tdap.
Meningococcal Recombinant
(serogroup B) (MenB) does not

		meet criteria for the Meningococcal vaccine.	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Lead Screening Children (LSC)	The percentage of children 2 years of age who had one or more capillary or venous lead blood test for lead poisoning at any time by their 2nd birthday.	 Documentation in the medical record must include both of the following on or before the 2nd birthday: A note indicating the date the test was performed. The result or finding. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. Common Chart Deficiencies: Lab results not documented in the record. Documentation of a lead assessment versus a lead screening. Lead screening not ordered, not completed, or result not documented. Lead screening after the child's 2nd birthday. Results of screening performed at an outside lab, health department, or WIC office not included in record. 	Lead Tests: CPT: 83655 Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

Measure	Measure Description	Measure Information/Documentation Required	Coding
Oral Evaluation, Dental Services (OED)	The percentage of members under 21 years of age who received a comprehensive or periodic oral evaluation with a dental provider during the MY.	 Documentation in the medical record must contain evidence of a comprehensive or periodic oral evaluation by a dental provider. Dental providers include dentist, dental hygienist, dental assistant, dental therapist, endodontist, denturist, oral medicinist, oral/maxillofacial dentist/surgeon. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. 	CDT: D0120, D0150 Dental Provider Taxonomy: 122300000X, 1223D0001X, 1223D0004X, 1223E0200X, 1223G0001X, 1223P0106X, 1223P0221X, 1223P0300X, 1223P0700X, 1223S0112X, 1223X0008X, 1223X0400X, 1223X2210X, 122400000X, 124Q00000X, 125J00000X, 125K00000X, 125Q00000X, 126800000X, 204E00000X, 261QD0000X, 261QF0400X, 261QR1300X, 261QS0112X
Measure	Measure Description	Measure Information/Documentation Required	Coding
Topical Fluoride for Children (TFC)	The percentage of members 1 – 4 years of age who received at least two fluoride varnish applications during the MY.	 Application of fluoride varnish on two different dates of service in the MY. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. 	CDT: 99188, D1206

Measure	Measure Description	Measure Information/Documentation	Coding
		Required	
Weight Assessment and	The percentage of	BMI Percentile:	BMI Percentile:
Counseling for Nutrition	members 3 – 17 years of	 Documentation must include 	ICD10CM: Z68.51, Z68.52, Z68.53, Z68.54
and Physical Activity for	age who had an	height, weight, and BMI	
Children/Adolescents	outpatient visit with a	percentile during the MY.	Nutrition Counseling:
(WCC)	PCP or OB/GYN and who	 The height, weight, and BMI must 	CPT: 97802, 97803, 97804
	had evidence of each of	be from the same data source.	HCPCS: G0447, S9452, S9470
	the following during the	BMI percentile can be	
	MY:	documented as a value or plotted	Encounter for Physical Activity Counseling: Z02.5, Z71.82
	BMI percentile	on an age-growth chart.	
	documentation.	 Member-reported values 	
	 Counseling for 	(weight, height, BMI) can be	
	nutrition.	captured during a telephone visit,	
	 Counseling for 	e-visit, or virtual check-in.	Note: LOINC and SNOMED codes can be captured through electronic data
	physical activity.		submissions. Please contact your Account Executive for more information.
		Counseling for Nutrition:	submissions. Theuse contact your Account Executive for more information.
		Documentation of counseling for	
		nutrition or referral for nutrition	
		education during the MY. Examples	
		include:	
		Discussion of current nutrition	
		behaviors (e.g., eating habits,	
		dieting behaviors).	
		Checklist indicating nutrition was	
		addressed.	
		Member received educational	
		materials on nutrition during a	
		face-to-face visit.	
		 Anticipatory guidance for 	
		nutrition.	
		Weight or obesity counseling.	
		Referral to the Special	
		Supplemental Nutrition Program	
		for Women, Infants, and Children	
		(WIC).	
		.	
		Counseling for Physical Activity:	

height, weight, BMI) documented in the chart.	
 activity (e.g., sports activities, exercise routines). Exam for sport participation/sports physical. Notes: Services may be rendered during a visit other than a well-child visit; however, services specific to the assessment or treatment of an acute or chronic condition do not count toward the "Counseling for Nutrition" and "Counseling for Physical Activity" indicators. Services may be delivered during a telephone visit, e-visit, or virtual check-in. This includes member-reported data (e.g., 	
 Documentation of counseling for physical activity or referral for physical activity during the MY. Examples include: Checklist indicating physical activity was addressed. Member received educational materials on physical activity during a face-to-face visit. Anticipatory guidance for physical activity or weight/obesity counseling. Weight or obesity counseling. Discussion of current physical 	

Members who meet any of the	
following criteria are excluded from	
the measure:	
In hospice or using hospice	
services any time in the MY.	
• Deceased at any time in the MY.	
 Diagnosis of pregnancy during 	
the MY.	
Common Chart Deficiencies:	
Height, weight, and BMI	
percentile not documented <i>each</i>	
year.	
BMI documented as a value and	
not as a percentile.	
BMI percentile documented as a	
range or threshold.	
BMI documented on an	
appropriate age-growth chart but	
without name, DOB, or	
discernible DOS on the chart.	
BMI documented on weight or	
stature for age charts.	
Documentation of developmental	
milestones without notation of	
anticipatory guidance or	
education for physical activity.	
Missing counseling/education on	
physical activity and/or nutrition.	
 Notation of "health education" or 	
"anticipatory guidance" without	
specific mention of nutrition	
and/or physical activity.	
 Counseling on safety (e.g., "wears 	
helmet" or "water safety")	
without specific mention of	
physical activity	
recommendations.	

		Required	
Appropriate Testing for	The percentage of	Outpatient, telephone, observation or	Group A Strep Test:
Pharyngitis (CWP)	episodes for members 3	ED visit, e-visit, or virtual check-in	CPT: 87070, 87071, 87081, 87430, 87650, 87651, 87652, 87880
	years and older where	with only a diagnosis of pharyngitis	
This is also a measure	the member was	and a dispensed antibiotic for that	Pharyngitis Diagnosis:
(CWP-E) collected through	diagnosed with	episode of care during the Intake	ICD10CM: J02.0, J02.8, J02.9, J03.00, J03.01, J03.80, J03.81, J03.90, J03.91
claims and Electronic	pharyngitis, dispensed	Period (IP), which is 3 days prior and 3	
Clinical Data Systems.	an antibiotic, and	days after the diagnosis.	
Please discuss options for	received a group A		Note: LOINC and SNOMED codes can be captured through electronic data
a direct data feed with	Streptococcus (Strep)	Visits that result in an inpatient stay	submissions. Please contact your Account Executive for more information.
your Account Executive.	test for the episode.	are excluded.	
Direct data feeds can			
improve provider quality			

www.amerihealthcaritasoh.com

performance and reduce the burden of medical record requests.	This is an episode-based event, so a member may be included multiple times.	 Telehealth visits are included in event/diagnosis criteria. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. Common Chart Deficiencies: Additional/competing diagnosis requiring antibiotics not documented in visit or coded on claim. 	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Asthma Medication Ratio (AMR)	The percentage of members 5 – 64 years of age who were identified as having persistent asthma and had a ratio of controller medications to total asthma medications of 50% or greater during the MY.	Oral medication-dispensing event: Multiple prescriptions for different medications dispensed on the same day are counted as separate dispensing events. If multiple prescriptions for the same medication are dispensed on the same day, sum the day's supply and divide by 30. Use the Drug ID to determine if the prescriptions are the same or different. Inhaler-dispensing event: All inhalers (i.e., canisters) of the same medication dispensed on the same day count as one dispensing event. Medications with different drug IDs dispensed on the same day are	 Population includes ED, IP, and/or observation visits billed with asthma diagnosis or 4 non-controller asthma medication-dispensing events during the MY and the year prior. Asthma Diagnosis:ICD10CM: J45.21, J45.22, J45.30, J45.31, J45.32, J45.40, J45.41, J45.42, J45.50, J45.51, J45.52, J45.901, J45.902, J45.909, J45.991, J45.998 Asthma Controller Medications: Antibody inhibitors: Omalizumab Anti-interleukin-4: Dupilumab Anti-interleukin-5: Benralizumab, Mepolizumab, Reslizumab Inhaled steroid combinations: Budesonide-formoterol, Fluticasone-salmeterol, Fluticasone-vilanterol, Formoterol-mometasone Inhaled corticosteroids: Beclomethasone, Budesonide, Ciclesonide, Flunisolide, Fluticasone, Mometasone Leukotriene modifiers: Montelukast, Zafirlukast, Zileuton Methylxanthines: Theophylline

	counted as different dispensing	
	events.	Asthma Reliever Medications:
	Injection-dispensing events: Each	Short-acting, inhaled beta-2 agonists: Albuterol, Levalbuterol
	injection counts as one dispensing event. Multiple dispensed injections of the same or different medications count as separate dispensing events.	Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
	Units of medications: When identifying medication units for the numerator, count each individual medication, defined as an amount lasting 30 days or less, as one medication unit. One medication unit equals one inhaler canister, one injection, or a 30-day or less supply of an oral medication.	
	 Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Members who had no asthma medications dispensed during the MY. Members who had a diagnosis of any of the following in the member's history through December 31 of the MY: emphysema, COPD, Obstructive Bronchitis, chronic respiratory conditions due to fumes/vapors, Cystic Fibrosis, acute respiratory failure. Deceased at any time in the MY. 	

Measure	Measure Description	Common Chart Deficiencies: No documentation of review of medications at every visit. Measure Information/Documentation Required	Coding
Medication Management for People with Asthma (MMA) Retired by NCQA in MY20 but may still apply in state quality reporting. Consult with your Account Executive.			Requires state-specific measure codes.
Measure	Measure Description	Measure Information/Documentation Required	Coding
Pharmacotherapy Management of COPD Exacerbation (PCE)	The percentage of members 40 years of age and older who had an acute inpatient discharge or ED visit on or between January 1 through November 30 of MY and who had evidence of an active prescription or were dispensed the appropriate medications: • A Systemic Corticosteroid within 14 days of the event, or	 Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. 	HEDIS rates are based on pharmacy claims. Systemic Corticosteroid Medications: Glucocorticoids: Cortisone, Dexamethasone, Hydrocortisone, Methylprednisolone, Prednisolone, Prednisone Bronchodilator Medications: Anticholinergic agents: Aclidinium bromide, Ipratropium, Tiotropium, Umeclidinium Beta 2-agonists: Albuterol, Arformoterol, Formoterol, Indacaterol, Levalbuterol, Metaproterenol, Olodaterol, Salmeterol Bronchodilator combinations: Albuterol-ipratropium, Budesonide-formoterol, Fluticasone-salmeterol, Fluticasone-vilanterol, Fluticasone furoate- umeclidinium-vilanterol, Formoterol-aclidinium, Formoterol-glycopyrrolate, Formoterol-mometasone, Glycopyrrolate-indacaterol, Olodaterol-tiotropium, Umeclidinium-vilanterol

ACCESS AND AVAILABILITY	 A Bronchodilator within 30 days of the event. This is an episode-based event, so a member may be included multiple times. 		
Measure	Measure Description	Measure Information/Documentation Required	Coding
Adults' Access to Preventive/Ambulatory Health Services (AAP)	The percentage of members 20 years and older who had an ambulatory or preventive care visit during the MY.	 One or more ambulatory or preventive care visits during the MY. Telephone and e-visits are acceptable. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. 	Ambulatory Visits: CPT: 92002, 92004, 92012, 92014, 98980, 98981, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99243, 99244, 99245, 99304, 99305, 99036, 99307, 99308, 99309, 99310, 99315, 99316, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99421, 99422, 99423, 99441, 99442, 99457, 99458, 99483 HCPCS: G0402, G0463, G2010, G2012, G2251, T1015 UBREV: 0510, 0513, 0515, 0516, 0517, 0519, 0900, 0904, 0911, 0914, 0915, 0916 0919 Reason for Ambulatory Visit: ICD10CM: Z00.00, Z00.01, Z00.121, Z00.129, Z00.3, Z00.5, Z00.8, Z02.0, Z02.1, Z02.2, Z02.3, Z02.4, Z02.5, Z02.6, Z02.71, Z02.79, Z02.81, Z02.82, Z02.83, Z02.89, Z02.9, Z76.1, Z76.2 Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

Measure	Measure Description	Measure Information/Documentation Required	Coding
Children and Adolescents' Access to Primary Care (CAP) Retired by NCQA in MY20 but may still apply in state quality reporting. Consult with your Account Executive.			Requires state-specific measure codes.
Measure	Measure Description	Measure Information/Documentation Required	Coding
Initiation and Engagement of Substance Use Disorder Treatment (IET)	The percentage of members with a new episode of substance use disorder (SUD) who received Initiation of SUD Treatment or Engagement of SUD Treatment. Two rates are reported: 1. Initiation of SUD Treatment: The percentage of new SUD episodes that result in treatment initiation through an inpatient SUD admission, outpatient visit, intensive outpatient encounter, partial hospitalization, telehealth, or medication treatment	 The MY is 1/1 – 12/31. Note: Methadone is not included in the medication lists for the measure. Medication treatment meets criteria for members being treated for alcohol or opioid abuse or dependence. It does not meet the criteria for treatment of other drug abuse or dependence. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. 	 Visit Setting Unspecified: (With Outpatient Place of Service (POS) and with Alcohol Abuse and Dependence, Opioid Abuse and Dependence, or Other Drug Abuse and Dependence): (with Partial Hospitalization POS and with Alcohol Abuse and Dependence, Opioid Abuse and Dependence, or Other Drug Abuse and Dependence): (With Behavioral Health (BH) Outpatient Visit and with Alcohol Abuse and Dependence, Opioid Abuse and Dependence, or Other Drug Abuse and Dependence, Opioid Abuse and Dependence, or Other Drug Abuse and Dependence): (with Non-Residential Substance Abuse Treatment Facility POS and with Alcohol Abuse and Dependence, Opioid Abuse and Dependence, or Other Drug Abuse and Dependence): (with Community Mental Health Center POS and with Alcohol Abuse and Dependence): (with Telehealth POS and with Alcohol Abuse and Dependence): (with Telehealth POS and with Alcohol Abuse and Dependence): (with Telehealth POS and with Alcohol Abuse and Dependence, Opioid Abuse and Dependence): (with Telehealth POS and with Alcohol Abuse and Dependence): (with Telehealth POS and with Alcohol Abuse and Dependence): (PT: 90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221, 99222, 99233, 99231, 99232, 99233, 99238, 99239, 99252, 99253, 99254, 99255

www.amerihealthcaritasoh.com

within 14 days of the	BH Outpatient Visit:
diagnosis.	(with Alcohol Abuse and Dependence, Opioid Abuse and Dependence, or Other
2. Engagement of SUD	Drug Abuse and Dependence):
Treatment:	CPT: 98960, 98961, 98962, 99078, 99202, 99203, 99204, 99205, 99211, 99212,
The percentage of new	99213, 99214, 99215, 99242, 99243, 99244, 99245, 99341, 99342, 99344,
SUD episodes that have	99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385,
evidence of treatment	99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401,
engagement within 34	99402, 99403, 99404, 99483, , 99510
days of the initiation	HCPCS: G0463, H0004, H0036, H0037, H0039, H2015, H2017, H2019, H2020,
visit.	T1015
Each qualifying episode	UBREV: 0513, 0515, 0516, 0517, 0519, 0900, 0904, 0911, 0914, 0915, 0916,
between 11/15 of the	0919
year prior to the MY and	
11/14 of the MY is	
included.	Partial Hospitalization or Intensive Outpatient Visit:
	(with Alcohol Abuse & Dependence, Opioid Abuse & Dependence, or Other
	Drug Abuse & Dependence):
	HCPCS: H2012, S9484, S9485
	UBREV: 0907, 0912, 0913
	Substance Use Disorder Services:
	(With AOD (Alcohol and Other Drug) Abuse and Dependence, Opioid Abuse and
	Dependence, or Other Drug Abuse and Dependence):
	HCPCS: G0396, G0397, H0001, H0005, H0015, H2036
	UBREV: 0906, 0944, 0945
	Substance Abuse Counseling and Surveillance:
	ICD10CM: Z71.41, Z71.51
	Telephone Visit:
	(with Alcohol Abuse and Dependence, Opioid Abuse and Dependence, or Other
	Drug Abuse and Dependence):
	CPT : 99442, 99443
	Online Assessments:
	(with Alcohol Abuse and Dependence, Opioid Abuse and Dependence, or Other
	Drug Abuse and Dependence):

	CPT: 99421, 99422, 99423, 99457, 99458
	HCPCS: G2010, G2012, G2251
	Outpatient POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72
	Non-Residential Substance Abuse POS: 57, 58
	Telehealth POS: 02, 10
	Alcohol Abuse and Dependence: ICD10CM: F10.10, F10.120, F10.121, F10.129, F10.130, F10.131, F10.132, F10.139, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232, F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281, F10.282, F10.288, F10.29
	Opioid Abuse and Dependence: ICD10CM: F11.10, F11.120, F11.121, F11.122, F11.129, F11.13, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.220, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29
	Other Drug Abuse and Dependence: ICD10CM: F12.10, F12.120, F12.121, F12.122, F12.129, F12.13, F12.150, F12.151, F12.159, F12.180, F12.188, F12.19, F12.20, F12.220, F12.221, F12.222, F12.229, F12.23, F12.250, F12.251, F12.259, F12.280, F12.288, F12.29, F13.10, F13.120, F13.121, F13.129, F13.130, F13.131, F13.132, F13.139, F13.14, F13.150, F13.151, F13.159, F13.180, F13.181, F13.182, F13.188, F13.19, F13.20, F13.220, F13.221, F13.229, F13.230, F13.231, F13.232, F13.239, F13.24, F13.250, F13.251, F13.259, F13.26, F13.27, F13.280, F13.281, F13.282, F13.288, F13.29, F14.10, F14.120, F14.121, F14.122, F14.129, F14.13, F14.14, F14.150, F14.151, F14.159, F14.180, F14.181, F14.182, F14.188, F14.19, F14.20, F14.220, F14.221, F14.222, F14.229, F14.23, F14.24, F14.250, F14.251, F14.259, F14.280, F14.281, F14.282, F14.288, F14.29, F15.10, F15.120, F15.121, F15.122, F15.129, F15.13, F15.14, F15.150, F15.151, F15.159, F15.180, F15.181, F15.182, F15.188,

			 F15.19, F15.20, F15.220, F15.221, F15.222, F15.229, F15.23, F15.24, F15.250, F15.251, F15.259, F15.280, F15.281, F15.282, F15.288, F15.29, F16.10, F16.120, F16.121, F16.122, F16.129, F16.14, F16.150, F16.151, F16.159, F16.180, F16.183, F16.188, F16.19, F16.20, F16.220, F16.221, F16.229, F16.24, F16.250, F16.251, F16.259, F16.280, F16.283, F16.288, F16.29, F18.10, F18.120, F18.121, F18.20, F18.20, F18.21, F18.259, F18.27, F18.20, F18.288, F18.29, F19.10, F19.120, F19.121, F19.122, F19.129, F19.130, F19.131, F19.132, F19.139, F19.14, F19.150, F19.151, F19.159, F19.16, F19.17, F19.180, F19.181, F19.182, F19.188, F19.19, F19.20, F19.20, F19.221, F19.222, F19.229, F19.230, F19.231, F19.232, F19.239, F19.24, F19.250, F19.251, F19.259, F19.26, F19.27, F19.280, F19.281, F19.282, F19.288, F19.29 Alcohol Use Disorder Treatment Medications List (if diagnosis from Alcohol Abuse and Dependence): Aldehyde dehydrogenase inhibitor: Disulfiram (oral) Antagonist: Naltrexone (oral and injectable) Other: Acamprosate (oral, delayed-release tablet) Opioid Use Disorder Treatment Medications (if diagnosis from Opioid Abuse and Dependence): Aldehyde dehydrogenase (oral and injectable) Partial Agonist: Buprenorphine (sublingual tablet, injection, implant), Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film) Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
Measure	Measure Description	Measure	Coding
		Information/Documentation Required	
Prenatal and Postpartum	The percentage of	Prenatal care visit to an OB/GYN or	Prenatal Indicator:
Care (PPC)	deliveries of live births	other prenatal care practitioner or	Stand Alone Prenatal Visits:
	on or between October	PCP. For visits to a PCP, a diagnosis of	CPT-CAT-II: 0500F, 0501F, 0502F
	8 of the year prior to the	pregnancy must be present.	HCPS: H1001, H1004
	MY and October 7 of the	Documentation in the medical record	
	MY. For these members,	must include a note indicating the	(Dates of service required to validate within measure time frame.)
	the measure assesses	date when the prenatal care visit	
	the following facets of	occurred, and evidence of one of the following:	Prenatal Visits (with Diagnosis of Pregnancy):

 prenatal and postpartum care. Timeliness of Prenatal Care. The percentage of deliveries that received a prenatal care visit in the first trimester, on or before the enrollment start date, or within 42 days of enrollment in the organization. Postpartum Care. The percentage of deliveries that had a postpartum visit on or between 7 and 84 days after delivery. 	 Documentation indicating pregnancy or reference to pregnancy (use of a standardized prenatal flow sheet, documentation of LMP, EDD, GA, a positive pregnancy test, gravidity and parity, a complete obstetrical history, prenatal risk assessment or counseling/education). A basic physical obstetrical examination that includes auscultation for fetal heart tone, pelvic exam with obstetric observations, or measurement of fundus height. Evidence that a prenatal care procedure was performed (OB panel, ultrasound, etc.). 	CPT: 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99241, 99242, 99243, 99244, 99245, 99421, 99422, 99423, 99441, 99442, 99443, 99457, 99458, 99483 HCPS: G0463, G2010, G2012, T1015 Postpartum Indica Encounter for Postpartum Care: Z01.411, Z01.419, Z01.42, Z30.430, Z39.1, Z39.2 Postpartum Care: CPT: 57170, 58300, 59430, 99501 CPT-CAT-II: 0503F HCPCS: G0101 Cervical Cytology Lab Test: CPT: 88141, 88142, 88143, 88147, 88148, 88150, 88152, 88153, 88164, 88165, 88166, 88167, 88174, 88175 HCPCS: G0123, G0143, G0144, G0145, G0147, G0148, Q0091
	 Postpartum visit to an OB/GYN or other prenatal care practitioner or PCP. Documentation in the medical record must include a note indicating the date when the postpartum care visit occurred, and evidence of one of the following: Pelvic Exam: Colposcopy is not acceptable for a postpartum visit. Evaluation of weight, BP, breast, and abdomen: Notation of "breastfeeding" is acceptable for the "evaluation of breasts" component. Notation of postpartum care, including, but not limited to: Notation of "postpartum care," "PP care," "PP Checks," "6-week check." 	Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

	A preprinted "Postpartum Care"
	form in which information was
	documented during the visit.
	Perineal or cesarean
	incision/wound check.
	Screening for depression, anxiety,
	tobacco use, substance use
	disorder, or preexisting mental
	health disorders.
	Glucose screening for women
	with gestational diabetes.
	Documentation of any of the
	following: infant care or
	breastfeeding, resumption of
	intercourse, birth spacing, family
	planning, sleep/fatigue,
	resumption of physical activity,
	attainment of healthy weight.
	Note:
	Services provided during a
	telephone visit, e-visit, or virtual
	check-in are acceptable.
	Services that occur over multiple
	visits count toward Timeliness of
	Prenatal Care if all services are
	within the time frame established
	in the measure. Ultrasound and
	in the measure. Ultrasound and lab results alone are not
	in the measure. Ultrasound and lab results alone are not considered a visit; they must be
	in the measure. Ultrasound and lab results alone are not considered a visit; they must be combined with an office visit with
	in the measure. Ultrasound and lab results alone are not considered a visit; they must be combined with an office visit with an appropriate practitioner in
	in the measure. Ultrasound and lab results alone are not considered a visit; they must be combined with an office visit with
	in the measure. Ultrasound and lab results alone are not considered a visit; they must be combined with an office visit with an appropriate practitioner in order to count for this measure.
	in the measure. Ultrasound and lab results alone are not considered a visit; they must be combined with an office visit with an appropriate practitioner in order to count for this measure.
	in the measure. Ultrasound and lab results alone are not considered a visit; they must be combined with an office visit with an appropriate practitioner in order to count for this measure.

		 In hospice or using hospice services any time in the MY. Deceased at any time in the MY. Non-live birth. Common Chart Deficiencies: Missing signature on charts so unable to determine provider type of services. Only initials on charts, so unable to determine provider type of services. Ultrasound and/or labs with no associated prenatal visit documented in measure time frame. Initial prenatal visit documented as intake with RN but no visit with OB/GYN or PCP. Diagnosis of pregnancy not documented in chart. Dates of service in progress notes do not align with dates on ONAF. 	
		 ONAF not filled out completely. Visit in postpartum time frame does not reference 	
		pregnancy/delivery.	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Use of First-Line	The percentage of	Documentation of psychosocial care	Psychosocial Care:
Psychosocial Care for	children and	in the 121-day period from 90 days	CPT: 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90846,
Children and Adolescents	adolescents 1 – 17 years	prior to the Rx dispensing date	90847, 90849, 90853, 90875, 90876, 90880
on Antipsychotics (APP)	of age who had a new	through 30 days after the Rx	
	prescription for an	dispensing date.	HCPCS: H0004, H0036, H0038, H0040, H2000, H2012, H2017, H2019, H2020,
	antipsychotic	Deguised Evolution	S9484, S9485
	medication and had documentation of	Required Exclusions:	Note: LOINC and SNOMED codes can be captured through electronic data
			submissions. Please contact your Account Executive for more information.

EFFECTIVENESS OF CARE: 0	psychosocial care as first-line treatment. CARDIOVASCULAR CONDITIO	 Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. 	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Controlling High Blood Pressure (CBP)	The percentage of members 18 – 85 years of age who had a diagnosis of hypertension (HTN) and whose BP was adequately controlled (<140/90) during the MY.	 BP must be latest reading in the MY and must occur on or after the diagnosis of HTN. BP readings taken on the same day as a diagnostic test or diagnostic or therapeutic procedure that requires a change in diet or change in medication on or one day before the test or procedure, with the exception of fasting blood tests, are not used. BP readings taken during an inpatient stay or ED visit are not used. When multiple BP measurements occur on the same date, the lowest systolic and lowest diastolic BP reading will be used. If no BP is recorded during the MY, the member is "not controlled." Services provided during a telephone visit, e-visit, or virtual check-in are acceptable. Member-reported data documented in medical record is acceptable if BP captured with a digital device and documented in 	 Systolic and Diastolic Result: CPT-CAT-II: Most Recent Systolic less than 130: 3074F Most Recent Systolic 130 – 139: 3075F Systolic greater than or equal to 140: 3077F Most Recent Diastolic less than 80: 3078F Most Recent Diastolic 80-89: 3079F Most Recent Diastolic greater than or equal to 90: 3080F Hypertension Diagnosis: ICD10CM: 110 Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

the medical record with date BP	
taken.	
Required Exclusions:	
Members who meet any of the	
following criteria are excluded from	
the measure:	
In hospice or using hospice	
services any time in the MY.	
• Deceased at any time in the MY.	
Receiving palliative care any time	
in the MY.	
• 66 years of age and older with	
frailty and advanced illness	
during the MY.	
Evidence of ESRD or kidney	
transplant on or prior to 12/31 of	
the MY. Documentation must	
include a dated note indicating	
evidence of ESRD, kidney	
transplant, or dialysis.	
• Diagnosis of pregnancy during the MY.	
 A nonacute inpatient admission 	
during the MY.	
Common Chart Deficiencies:	
Retake of BP that is 140/90 or	
above not documented.	
Member-reported BP is not	
documented with sufficient	
detail.	
Claim missing CPT II codes for BP	
results.	
• BP rounded up before	
documented in medical record.	
• BP documented as a range.	

Measure	Measure Description	 No documentation of follow-up appointment scheduled if BP elevated. Cardiology visits with no BP documented in the chart. Flowsheets missing member name and second identifier such as date of birth. Measure Information/Documentation Required 	Coding
Persistence of Beta Blocker Treatment After a Heart Attack (PBH)	The percentage of members 18 years of age and older during the MY who were hospitalized and discharged from 7/1 of the year prior to the MY to 6/30 of the MY with a diagnosis of acute myocardial infarction (AMI) and who received persistent beta-blocker treatment for 180 days (6 months) after discharge.	 Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. Receiving palliative care any time in the MY. 66 years of age and older with advanced illness during the MY. 81 years of age and older with frailty any time on or between 7/1 of the year prior to the MY and 12/31 of the MY. Documentation of any of the following: Asthma. COPD. Obstructive chronic bronchitis. Chronic respiratory conditions due to fumes or vapors. Hypotension. Heart block >1 degree. Sinus bradycardia. 	 HEDIS rates are based on pharmacy claims. Beta-Blocker Medications: Noncardioselective beta-blockers: Carvedilol, Labetalol, Nadolol, Pindolol, Propranolol, Timolol, Sotalol Cardioselective beta-blockers: Acebutolol, Atenolol, Betaxolol, Bisoprolol, Metoprolol, Nebivolol Antihypertensive combinations: Atenolol-chlorthalidone, Bendroflumethiazide-nadolol, Bisoprolol-hydrochlorothiazide, Hydrochlorothiazide-metoprolol, Hydrochlorothiazide-propranolol AMI Diagnosis: ICD10CM: 121.01, 121.02, 121.09, 121.11, 121.19, 121.21, 121.29, 121.3, 121.4 Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

		 A medication-dispensing event indicative of a history of asthma. Intolerance or allergy to beta-blocker therapy. Common Chart Deficiencies: Medication was ordered with no evidence that it was dispensed. 	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Cardiac Rehabilitation (CRE)	 The percentage of members 18 years and older who attended cardiac rehabilitation following a qualifying cardiac event, including: Myocardial infarction. Percutaneous coronary intervention. Coronary artery bypass grafting. Heart and heart/lung transplantation. Heart valve repair/replacement. Four rates are reported as the percentage of members who attended the specified number of cardiac rehabilitation sessions within the 	The MY is 1/1 – 12/31. The Intake Period (IP) is a 12-month window that begins on July 1 of the year prior to the MY and ends on June 30 of the MY. The Episode Date (EP) is the most recent cardiac event during the IP, including myocardial infarction (MI), coronary artery bypass graft (CABG), percutaneous coronary intervention (PCI), heart or heart/lung transplant, or heart valve repair/replacement. For MI, CABG, heart or heart/lung transplant or heart valve repair/replacement, the EP is the date of discharge. For PCI, the EP is the date of service. For inpatient claims, the EP is the date of discharge. Required Exclusions:	Cardiac Rehabilitation: CPT: 93797, 93798 HCPCS: G0422, G0423, S9472 Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

	 specified time after a qualifying event: 1. Initiation: 2 or more sessions within 30 days. 2. Engagement 1: 12 or more sessions within 90 days. 3. Engagement 2: 24 or more sessions within 180 days. 4. Achievement: 36 or more sessions within 180 days. 	 Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. Receiving palliative care during the IP through the end of the MY. 66 years of age and older with frailty and advanced illness during the MY. 81 years of age and older with frailty during the IP through the end of the MY. Discharged from an inpatient setting with the following during the 180 days after the EP: MI, CABG, heart or heart/lung transplant, heart valve repair or replacement. PCI in any setting during the 180 days after the EP. 	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Statin Therapy for	The percentage of males	The Index Prescription Start Date	High-intensity statin therapy: Atorvastatin (40 – 80 mg), Amlodipine-
Patients with	21 – 75 years of age and	(IPSD) is the earliest dispensing date	atorvastatin (40 – 80 mg), Rosuvastatin (20 – 40 mg), Simvastatin (80 mg),
Cardiovascular Disease	females 40 – 75 years of	for any statin medication of at least	Ezetimibe-simvastatin (80 mg)
(SPC)	age during the MY who	moderate intensity during the MY.	
	were identified as		Moderate-intensity statin therapy : Atorvastatin (10 – 20 mg), Amlodipine-
	having clinical atherosclerotic	The Treatment Period (TP) is the	Atorvastatin $(10 - 20 \text{ mg})$, Rosuvastatin $(5 - 10 \text{ mg})$, Simvastatin $(20 - 40 \text{ mg})$, Exotimibe cimulattatin $(20 - 40 \text{ mg})$, Brayastatin $(40 - 80 \text{ mg})$, Joyastatin $(40 - 10 \text{ mg})$
	cardiovascular disease	period beginning on the IPSD through 12/31 of the MY.	Ezetimibe-simvastatin (20 – 40 mg), Pravastatin (40 – 80 mg), Lovastatin (40 mg), Fluvastatin (40 – 80 mg), Pitavastatin (1 – 4 mg)
	(ASCVD) and met the		111g), 11avastatii (40 – 00 111g), Fitavastatii (1 – 4 111g)
	. ,	Required Exclusions:	MI Diagnosis:
	Tollowing criteria		
	following criteria.	•	-
	Two rates are reported:	Members who meet any of the following criteria are excluded from	ICD10CM: I21.01, I21.02, I21.09, I21.11, I21.19, I21.21, I21.29, I21.3, I21.4, I21.9, I21.41, I21.49, I22.0, I22.1, I22.2, I22.8, I22.9, I23.0, I23.1, I23.2, I23.3,

www.amerihealthcaritasoh.com

1. Received Statin Therapy: Members who were dispensed at least one high- or moderate- intensity statin medication during the	 In hospice or using hospice services any time in the MY. Deceased at any time in the MY. Receiving palliative care any time in the MY. 66 years of age and older with frailty and advanced illness 	ICD10PCS: 0210083, 0210088, 0210089, 0210093, 0210098, 0210099, 0211083, 0211088, 0211089, 0211093, 0211098, 0211099, 0212083, 0212088, 0212089, 0212093, 0212098, 0212099, 0213083, 0213088, 0213089, 0213093, 0213098, 0213099, 021008C, 021008F, 021008W, 021009C, 021009F, 021009W, 02100A3, 02100A8, 02100A9, 02100AC, 02100AF, 02100AW, 02100J3, 02100J8, 02100J9, 02100JC, 02100JF, 02100JW, 02100K3, 02100K8, 02100K9, 02100KC,
MY. 2. Statin Adherence 80%: Members who remained	 during the MY. Documentation of any of the following in the MY or year prior: Pregnancy, IVF treatment, 	02100KF, 02100KW, 02100Z3, 02100Z8, 02100K9, 02100KC, 02100ZF, 02100KF, 02100KW, 02100Z3, 02100Z8, 02100Z9, 02100ZC, 02100ZF, 021108C, 021108F, 021109F, 021109W, 02110JA, 02110A8, 02110A9, 02110AC, 02110AF, 02110AW, 02110J3, 02110J8, 02110J9, 02110JC, 02110JF, 02110JW, 02110K3, 02110K8, 02110K9, 02110KC, 02110KF, 02110KW,
on a high- or moderate- intensity statin medication for at least 80% of the treatment period.	 dispensed prescription for Clomiphene, cirrhosis, end stage renal disease (ESRD), or dialysis. Documentation of any of the following in the MY: Myalgia, myositis, myopathy, or rhabdomyolysis. 	02110Z3, 02110Z8, 02110Z9, 02110ZC, 02110ZF, 021208C, 021208F, 021208W, 021209C, 021209F, 021209W, 02120A3, 02120A8, 02120A9, 02120AC, 02120AF, 02120AW, 02120J3, 02120J8, 02120J9, 02120JC, 02120JF, 02120JW, 02120K3, 02120K8, 02120K9, 02120KC, 02120KF, 02120KW, 02120Z3, 02120Z8, 02120Z9, 02120ZC, 02120ZF, 021308C, 021308F, 021308W, 021309C, 021309F, 021309W, 02130A3, 02130A8, 02130A9, 02130AC, 02130AF, 02130AW, 02130J3, 02130J8, 02130J9, 02130JC, 02130JF, 02130JW, 02130K3, 02130K8, 02130K9, 02130KC, 02130KF, 02130KW, 02130Z3, 02130Z8, 02130Z9, 02130ZC,
	Common Chart Deficiencies: No documentation of review of 	O2130ZF PCI Diagnosis:
	medications at every visit.	CPT: 92920, 92924, 92928, 92933, 92937, 92941, 92943 HCPCS: C9600, C9602, C9604, C9606, C9607 ICD10PCS: 0270346, 0270356, 0270366, 0270376, 0270446, 0270456, 0270466, 0270476, 0271346, 0271356, 0271366, 0271376, 0271446, 0271456, 0271466, 0271476, 0272346, 0272356, 0272366, 0272376, 0272446, 0272456, 0272466, 0272476, 0273346, 0273356, 0273366, 0273376, 0273446, 0273456, 0273466, 0273476, 02703E6, 02704E6, 02713E6, 02714E6, 02723E6, 02724E6, 02733E6,
		02734E6, 027034Z, 027035Z, 027036Z, 027037Z, 02703D6, 02703DZ, 02703EZ, 02703F6, 02703FZ, 02703G6, 02703GZ, 02703T6, 02703TZ, 02703Z, 027044Z, 027045Z, 027046Z, 027047Z, 02704D6, 02704DZ, 02704EZ, 02704F6, 02704FZ, 02704G6, 02704GZ, 02704T6, 02704TZ, 02704Z6, 02704ZZ, 027134Z, 027135Z, 027136Z, 027137Z, 02713D6, 02713DZ, 02713ZZ, 02713F6, 02713FZ, 02713G6, 02713GZ, 02713TG, 02713TZ, 02713Z6, 02713ZZ, 02714FZ, 02714FZ, 02714F6, 02714FZ, 02714FG, 02714FZ, 02714FG, 02714FZ, 02714FG, 02714FZ, 02714FG, 02714FZ, 02714FG, 02714FZ, 02714FG, 02714FZ, 0271
		0271462, 0271472, 0271406, 0271402, 0271422, 0271462, 0271466, 0271466, 0271466, 0271466, 0271466, 0271472, 0271426, 0271427, 0272342, 0272352, 0272362, 0272372, 0272306, 0272302, 02723E2, 02723F6, 02723F2, 02723G6, 02723G2, 02723T6, 02723T2, 0272372, 02723Z2, 0272442, 02724452, 0272462, 0272472,

02724D6, 02724DZ, 02724EZ, 02724F6, 02724FZ, 02724G6, 02724GZ, 02724T6, 02724TZ, 02724Z6, 02724ZZ, 027334Z, 027335Z, 027336Z, 027337Z, 02733D6, 02733DZ, 02733EZ, 02733F6, 02733FZ, 02733G6, 02733GZ, 02733T6, 02733TZ, 02733Z6, 02733ZZ, 027344Z, 027345Z, 027346Z, 027347Z, 02734D6, 02734DZ, 02734EZ, 02734F6, 02734FZ, 02734G6, 02734GZ, 02734T6, 02734TZ, 02734Z6, 02734ZZ

Other Revascularization Diagnosis:

CPT: 37220, 37221, 37224, 37225, 37226, 37227, 37228, 37229, 37230, 37231

IVD Diagnosis:

ICD10CM: I20.0, I20.2, I20.8, I20.9, I24.0, I24.8, I24.9, I25.10, I25.110, I25.111, 125.112, 125.118, 125.119, 125.5, 125.6, 125.700, 125.701, 125.702, 125.708, 125.709, 125.710, 125.711, 125.712, 125.718, 125.719, 125.720, 125.721, 125.722, 125.728, 125.729, 125.730, 125.731, 125.732, 125.738, 125.739, 125.750, 125.751, 125.752, 125.758, 125.759, 125.760, 125.761, 125.762, 125.768, 125.769, 125.790, 125.791, 125.792, 125.798, 125.799, 125.810, 125.811, 125.812, 125.82, 125.83, 125.84, 125.89, 125.9, 163.20, 163.211, 163.212, 163.213, 163.219, 163.22, 163.231, 163.232, 163.233, 163.239, 163.29, 163.50, 163.511, 163.512, 163.513, 163.519, 163.521, 163.522, 163.523, 163.529, 163.531, 163.532, 163.533, 163.539, 163.541, 163.542, 163.543, 163.549, 163.59, 165.01, 165.02, 165.03, 165.09, 165.1, 165.21, 165.22, 165.23, 165.29, 165.8, 165.9, 166.01, 166.02, 166.03, 166.09, 166.11, 166.12, 166.13, 166.19, 166.21, 166.22, 166.23, 166.29, 166.3, 166.8, 166.9, 167.2, 170.1, 170.201, 170.202, 170.203, 170.208, 170.209, 170.211, 170.212, 170.213, 170.218, 170.219, 170.221, 170.222, 170.223, 170.228, 170.229, 170.231, 170.232, 170.233, 170.234, 170.235, 170.238, 170.239, 170.241, 170.242, 170.243, 170.244, 170.245, 170.248, 170.249, 170.25, 170.261, 170.262, 170.263, 170.268, 170.269, 170.291, 170.292, 170.293, 170.298, 170.299, 170.301, 170.302, 170.303, 170.308, 170.309, 170.311, 170.312, 170.313, 170.318, 170.319, 170.321, 170.322, 170.323, 170.328, 170.329, 170.331, 170.332, 170.333, 170.334, 170.335, 170.338, 170.339, 170.341, 170.342, 170.343, 170.344, 170.345, 170.348, 170.349, 170.35, 170.361, 170.362, 170.363, 170.368, 170.369, 170.391, 170.392, 170.393, 170.398, 170.399, 170.401, 170.402, 170.403, 170.408, 170.409, 170.411, 170.412, 170.413, 170.418, 170.419, 170.421, 170.422, 170.423, 170.428, 170.429, 170.431, 170.432, 170.433, 170.434, 170.435, 170.438, 170.439, 170.441, 170.442, 170.443, 170.444, 170.445, 170.448, 170.449, 170.45, 170.461, 170.462, 170.463, 170.468, 170.469, 170.491, 170.492, 170.493, 170.498, 170.499, 170.501, 170.502, 170.503, 170.508, 170.509, 170.511, 170.512, 170.513, 170.518, 170.519, 170.521, 170.522, 170.523, 170.528, 170.529,

170.531, 170.532, 170.533, 170.534, 170.535, 170.538, 170.539, 170.541, 170.542, 170.543, 170.544, 170.545, 170.548, 170.549, 170.55, 170.561, 170.562, 170.563, 170.568, 170.569, 170.591, 170.592, 170.593, 170.598, 170.599, 170.601, 170.602, 170.603, 170.608, 170.609, 170.611, 170.612, 170.613, 170.618, 170.619, 170.621, 170.622, 170.623, 170.628, 170.629, 170.631, 170.632, 170.633, 170.634, 170.635, 170.638, 170.639, 170.641, 170.642, 170.643, 170.644, 170.645, 170.648, 170.649, 170.65, 170.661, 170.662, 170.663, 170.668, 170.669, 170.691, 170.692, 170.693, 170.698, 170.699, 170.701, 170.702, 170.703, 170.708, 170.709, 170.711, 170.712, 170.732, 170.733, 170.734, 170.735, 170.738, 170.739, 170.741, 170.742, 170.743, 170.744, 170.745, 170.748, 170.749, 170.75, 170.761, 170.762, 170.763, 170.768, 170.769, 170.791, 170.792, 170.793, 170.798, 170.799, 170.92, 175.011, 175.012, 175.013, 175.019, 175.021, 175.022, 175.023, 175.029, 175.81, 175.89, T82.855A, T82.855D, T82.855A, T82.856D, T82.856D

Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

EFFECTIVENESS OF CARE: I	DIABETES		
Measure	Measure Description	Measure Information/Documentation Required	Coding
Glycemic Status Assessment for Patients With Diabetes (GSD) Formerly the HBD A1c Control for Patients with Diabetes indicator.	The percentage of members 18 – 75 years of age with diabetes (Type 1 or Type 2) whose most recent glycemic status (glucose management indicator [GMI] or hemoglobin A1c (HbA1c)) was at the following levels in the MY:	At a minimum, the documentation in the medical record must include a note indicating the date when the most recent HbA1c test was performed in the MY and the result or findings. Ranges and thresholds DO NOT meet criteria — a distinct numeric result is required.	 HbA1c Lab Test: CPT: 83036, 83037 HbA1c Test Result or Finding: CPT-CAT-II: Less than 7.0: 3044F Greater than or equal to 7.0 and less than 8.0: 3051F Greater than or equal to 8.0 and less than or equal to 9.0: 3052F Greater than 9.0: 3046F
	 Glycemic Status <8.0% Glycemic Status >9% 	Terms below, with date of service and result, can be used: A1c, Hemoglobin A1c, Glycated Hemoglobin, HbA1c, Glycohemoglobin A1c, Glycosylated Hemoglobin, HgA1c.	Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

A lower rate in Poor		
Control (>9%) indicates	Required Exclusions:	
better performance.	Members who meet any of the	
	following criteria are excluded from	
	the measure:	
	In hospice or using hospice	
	services any time in the MY.	
	• Deceased at any time in the MY.	
	Receiving palliative care any time	
	in the MY.	
	66 years of age or older who are	
	living long term in an institution	
	at any time during the	
	measurement year.	
	 66 years of age and older with 	
	frailty and advanced illness	
	during the MY.	
	Common Chart Deficiencies:	
	 A1c noted in the chart but 	
	without specific date.	
	In-house A1c noted in visit but no	
	result documented.	
	A1c result documented as a	
	range.	
	Diabetes diagnosis and	
	medication documented but	
	missing documentation of	
	treatment, follow-up, and/or	
	progress.	
	Flowsheets missing member	
	name and second identifier such	
	as date of birth.	
	Incomplete or missing	
	information from specialists or	
	consulting providers.	

Measure	Measure Description	Measure Information/Documentation Required	Coding
Comprehensive Diabetes Care (CDC) Monitoring for Nephropathy			Requires state-specific measure codes.
Retired by NCQA in MY22 but may still apply in state quality reporting. Consult with your Account Executive.			
Measure	Measure Description	Measure Information/Documentation Required	Coding
Eye Exam for Patients with Diabetes (EED) Formerly the CDC Eye Exam indicator.	The percentage of members 18 – 75 years of age with diabetes (Type 1 and Type 2) who had a retinal eye exam during the MY measurement year (MY), an exam with a negative result in the year prior to the MY, or documentation of bilateral eye enucleation any time prior to 12/31 of the MY.	 Documentation can include any of the following noted in the medical record: A note or letter during the MY prepared by an ophthalmologist, optometrist, PCP, or other health care provider indicating that an ophthalmoscopic exam was completed by an eye care provider, the date when the procedure was performed and the results. Documentation of a negative (or normal) retinal or dilated exam by an eye care provider in the year prior to the MY, where results indicate retinopathy was not present and the date when the exam was performed. A chart or photograph indicating the date when the fundus photography was performed and evidence that an eye care 	Diabetic Retinal Screening: CPT: 67028, 67030, 67031, 67036, 67039, 67040, 67041, 67042, 67043, 67101, 67105, 67107, 67108, 67110, 67113, 67121, 67141, 67145, 67208, 67210, 67218, 67220, 67221, 67227, 67228, 92002, 92004, 92012, 92014, 92018, 92019, 92134, 92201, 92202, 92227, 92228, 92230, 92235, 92240, 92250, 92260, 99203, 99204, 99205, 99213, 99214, 99215, 99242, 99243, 99244, 99245 Diabetes Mellitus without Complications (in Year Prior to MY with Diabetic Retinal Screening): ICD10CM: E10.9, E11.9, E13.9 Eye Exam without Evidence of Retinopathy: CPT-CAT-II: 2023F, 2025F, 2033F Eye Exam with Evidence of Retinopathy (in the MY Only): CPT-CAT-II: 2022F, 2024F, 2026F Unilateral Eye Enucleation (with Bilateral Modifier or 2 Unilateral Enucleations More than 14 Days Prior Apart): CPT: 65091, 65093, 65101, 65103, 65105, 65110, 65112, 65114

professional (optometrist or	
ophthalmologist) or qualified	
reading center reviewed the	
results, or that results were read	
by a system that provides	Note: LOINC and SNOMED codes can be captured through electronic data
artificial intelligence (AI)	submissions. Please contact your Account Executive for more information.
interpretation.	
Hypertensive retinopathy is handled	
the same as diabetic retinopathy	
when reporting the Eye Exam	
indicator.	
Positive for hypertensive	
retinopathy is counted as positive	
for diabetic retinopathy if	
diabetic retinopathy not	
documented.	
An eye exam documented as	
negative for hypertensive	
retinopathy is counted as	
negative for diabetic retinopathy	
if diabetic retinopathy not documented.	
documented.	
Common Abbreviations for	
Retinopathy:	
NPDR (Non-proliferative diabetic	
retinopathy).	
PDR (Proliferative diabetic	
retinopathy).	
BDR (Background diabetic	
retinopathy).	
Mild BDR or PDR.	
• Severe PDR.	
Examples of Negative Exam:	
Assessment of fundus and	
macula were "normal."	

 Diabetes mellitus without ophthalmic complication. Retinal exam documented as "normal" is considered negative for Retinopathy if diabetic retinopathy not documented. 	
Note: Notation limited to a statement that included "Diabetes without complications" does not meet criteria.	
 Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. Receiving palliative care any time in the MY. 66 years of age and older with frailty and advanced illness during the MY. 	
Blindness is not an exclusion for a diabetic eye exam.	
 Common Chart Deficiencies: Documentation of diabetic exam without results. Documentation of diabetic eye exam without provider (including credentials) of the exam. Documentation is not clear that patient had a dilated or retinal exam. Documentation not specific as to presence of retinopathy. 	

Measure	Measure Description	 Incomplete or missing information from specialists or consulting providers. Documentation of "diabetes without complications" does not meet criteria. Measure Information/Documentation Required 	Coding
Blood Pressure Control	The percentage of	BP must be latest reading in the	Systolic and Diastolic Result:
for Patients with	members 18 – 75 years	MY.	CPT-CAT-II:
Diabetes (BPD)	of age with diabetes	 BP readings taken on the same 	Most Recent Systolic less than 130: 3074F
	(Type 1 and Type 2) who	day as a diagnostic test or	 Most Recent Systolic 130 – 139: 3075F
Formerly the CDC BP	had a controlled BP of	diagnostic or therapeutic	Systolic greater than or equal to 140: 3077F
indicator.	<140/90 mm Hg during	procedure that requires a change	
	the MY.	in diet or change in medication on or one day before the test or	 Most Recent Diastolic less than 80: 3078F Most Recent Diastolic 80-89: 3079F
		procedure, with the exception of	 Most Recent Diastolic 80-89, 5079F Most Recent Diastolic greater than or equal to 90: 3080F
		fasting blood tests, are not used.	Wost Neterit Diastolie greater than of equal to 50. 50001
		 BP readings taken during an 	
		inpatient stay or ED visit are not	Note: LOINC and SNOMED codes can be captured through electronic data
		used.	submissions. Please contact your Account Executive for more information.
		When multiple BP measurements	
		occur on the same date, the	
		lowest systolic and lowest	
		diastolic BP reading will be used.	
		 If no BP is recorded during the MY, the member is "not 	
		controlled."	
		Member-reported data	
		documented in medical record is	
		acceptable if BP captured with a	
		digital device.	
		Required Exclusions:	
		Members who meet any of the	
		following criteria are excluded from	
		the measure:	

	(Type 1 and Type 2) who received a kidney health evaluation, defined by an estimated glomerular filtration rate (eGFR) and a urine albumin-	 Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. 	CPT: 80047, 80048, 80053, 80069, 82565 Quantitative Urine Albumin Lab Test: CPT: 82043 Urine Creatinine Lab Test: CPT: 82570
Patients With Diabetes (KED)	members 18 – 85 years of age with diabetes	required tests with result and date of service.	Estimated Glomerular Filtration Rate Lab Test:
Kidney Evaluation for	The percentage of	Required Documentation must include the	All three are required:
Measure	Measure Description	Measure Information/Documentation	Coding
		 BP documented as a range. Claim missing CPT II codes for BP results. Flowsheets missing member name and second identifier such as date of birth. Incomplete or missing information from specialists or consulting providers. 	
		 during the MY. Common Chart Deficiencies: Retake of BP that is 140/90 or above not documented. Member-reported BP is not documented with sufficient detail. BP rounded up before documented in medical record. 	
		 In hospice or using hospice services any time in the MY. Deceased at any time in the MY. Receiving palliative care any time in the MY. 66 years of age and older with frailty and advanced illness 	

	creatinine ratio (uACR), during the MY.	 Deceased at any time in the MY. Receiving palliative care any time in the MY. Evidence of ESRD or dialysis any time during the member's history through 12/31 of the MY. 66 years of age and older with frailty and advanced illness during the MY. 81 years of age and older with frailty during the MY. 	Service dates of Quantitative Urine Albumin Lab Test and Urine Creatinine Lab Test must be four or less days apart. <i>Note: LOINC and SNOMED codes can be captured through electronic data</i> <i>submissions. Please contact your Account Executive for more information.</i>
Measure	Measure Description	Measure Information/Documentation Required	Coding
Statin Therapy for Patients with Diabetes (SPD)	The percentage of members 40 – 75 years of age during the MY with diabetes who do not have clinical atherosclerotic cardiovascular disease (ASCVD) who met the following criteria. Two rates are reported: 1. Received statin therapy: Members who were dispensed at least one statin medication of any intensity during the MY. 2. Statin adherence 80%: Remained on a statin medication of any intensity for at least 80% of the treatment period.	 The Index Prescription Start Date (IPSD) is the earliest dispensing date for any statin medication of any intensity during the MY. The Treatment Period (TP) is the period beginning on the IPSD through 12/31 of the MY. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. Receiving palliative care any time in the MY. 66 years of age and older with frailty and advanced illness during the MY. Documentation of any of the following during the year prior to the MY: MI (myocardial 	Low-, Medium-, or High-Intensity Statin: Amlodipine-Atorvastatin, Atorvastatin, Ezetimibe-Simvastatin, Fluvastatin Lovastatin, Pitavastatin, Pravastatin, Rosuvastatin, Simvastatin

EFFECTIVENESS OF CARE: N		 infarction), CABG (coronary artery bypass graft), PCI (percutaneous coronary intervention), or other revascularization. Documentation of any of the following during the MY or the year prior: pregnancy, IVF, dispensed prescription for Clomiphene, ESRD, dialysis, or cirrhosis. Documentation of any of the following in the MY: myalgia, myositis, myopathy, or rhabdomyolysis. Diagnosis of ischemic vascular disease during the MY or the year prior who had at least one outpatient visit, telephone visit, online assessment, or acute inpatient encounter. No diagnosis of diabetes in any setting during the MY or the year prior and who had a diagnosis of polycystic ovarian syndrome, gestational diabetes, or steroid- induced diabetes during the MY or the year prior. 	
Measure	Measure Description	Measure Information/Documentation	Coding
		Required	
Disease-Modifying Anti- Rheumatic Drug Therapy			Requires state-specific measure codes.
for Rheumatoid Arthritis			
(ART)			
Retired by NCQA in MY20			
but may still apply in state			

quality reporting. Consult with your Account Executive.			
Measure	Measure Description	Measure Information/Documentation Required	Coding
Osteoporosis	The percentage of	The MY is 1/1 – 12/31.	HEDIS rates are based on pharmacy claims/BMD testing.
Management in Women	women 67 – 85 years of	The Intake Period (IP) is a 12-month	
Who Had a Fracture (OMW)	age who suffered a fracture and who had either a bone or mineral density (BMD) test or prescription for a drug to treat osteoporosis in the 180 days (6 months) after the fracture. Fractures of finger, toe, face, and skull are not	 window beginning 7/1 of the year prior to the MY and ending 6/30 of the MY. The IP is used to capture the first fracture. The Episode Date (EP) is an eligible encounter during the IP with a diagnosis of fracture. For outpatient, observation, or ER visit, the EP is the date of service. For inpatient stay, the EP is the date of discharge. 	Bone Mineral Density Tests: CPT: 76977, 77078, 77080, 77081, 77085, 77086 ICD10PCS: BP48ZZ1, BP49ZZ1, BP4GZZ1, BP4HZZ1, BP4HZZ1, BP4MZZ1, BP4NZZ1, BQ00ZZ1, BQ01ZZ1, BQ03ZZ1, BQ04ZZ1, BR00ZZ1, BR07ZZ1, BR09ZZ1, BR09ZZ1, BR0GZZ1 Osteoporosis Medication Therapy: HCPCS: J0897, J1740, J3110, J3111, J3489 Long-Acting Osteoporosis Medications: HCPCS: J0897, J1740, J3489
	included in this		Osteoporosis Medications List:
	measure.	Required Exclusions:	Bisphosphonates: Alendronate, Alendronate-cholecalciferol, Ibandronate,
		Members who meet any of the	Risedronate, Zoledronic acid
		 following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. 	Other Agents: Abaloparatide, Denosumab, Raloxifene, Romosozumab, Teriparatide
		 Deceased at any time in the MY. Receiving palliative care during the IP through the end of the MY. 67 – 80 years of age with frailty and advanced illness during the IP through the end of the MY. 81 years of age and older with frailty during the IP through the end of the MY. Had a BMD test during the 730 days prior to the ED. 	Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

		 Had a claim/encounter for osteoporosis therapy prior to the ED. Received a dispensed prescription or had an active prescription to treat Osteoporosis during the 365 days prior to the ED. 	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Osteoporosis Screening in Older Women (OSW)	The percentage of women 65 – 75 who received osteoporosis screening.	 One or more osteoporosis screening tests on or between the member's 65th birthday and 12/31 of the MY. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. Receiving palliative care any time in the MY. 66 years of age and older with frailty and advanced illness during the IP through the end of the MY. Had a claim/encounter for osteoporosis therapy any time in the mY. Had a dispensed dementia medication in the MY or the year prior to the MY. Had a dispensed prescription to treat osteoporosis any time from 1/1 three years prior to the MY 	Osteoporosis Screening Tests: CPT: 76977, 77078, 77080, 77081, 77085

		through 12/31 of the year prior to the MY.	
EFFECTIVENESS OF CARE: B	EHAVIORAL HEALTH		
Measure	Measure Description	Measure Information/Documentation Required	Coding
Medication Management (AMM)	members 18 years of age and older who were treated with antidepressant medication, had a diagnosis of major depression, and who remained on an	window starting on 5/1 of the year prior to the MY and ending on 4/30 of the MY. The Index Prescription Start Date (IPSD) is the earliest dispensing date for an antidepressant medication in the IP.	Major Depression Diagnosis: ICD10CM: F32.0, F32.1, F32.2, F32.3, F32.4, F32.9, F33.0, F33.1, F33.2, F33.3, F33.41, F33.9 Antidepressant Medications: Miscellaneous antidepressants: Bupropion, Vilazodone, Vortioxetine Monoamine oxidase inhibitors: Isocarboxazid, Phenelzine, Selegiline,
	antidepressant medication treatment. Two rates are reported: 1. Effective Acute Phase Treatment: The percentage of members who remained on an antidepressant medication for at least 84 days (12 weeks). 2. Effective Continuation Phase	 Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. No encounter with diagnosis of major depression during the 121- day period from 60 days prior to the IPSD, through 60 days after the IPSD. 	Tranylcypromine Phenylpiperazine antidepressants: Nefazodone, Trazodone Psychotherapeutic combinations: Amitriptyline-chlordiazepoxide, Amitriptyline-perphenazine, Fluoxetine-olanzapine SNRI antidepressants: Desvenlafaxine, Duloxetine, Levomilnacipran, Venlafaxine SSRI antidepressants: Citalopram, Escitalopram, Fluoxetine, Fluvoxamine, Paroxetine, Sertraline Tetracyclic antidepressants: Maprotiline, Mirtazapine Tricyclic antidepressants: Amitriptyline, Amoxapine, Clomipramine, Desipramine, Doxepin (>6mg), Imipramine, Nortriptyline, Protriptyline, Trimipramine
	Treatment: The percentage of members who remained on an antidepressant medication for at least 180 days (6 months).		Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

Measure	Measure Description	Measure Information/Documentation Required	Coding
Follow-Up After Hospitalization for Mental Illness (FUH)	Percentage of discharges for members 6 years of age and older who were hospitalized for treatment of selected mental illness or intentional self-harm diagnoses and who had a follow-up visit with a mental health provider.	 The MY is 1/1 – 12/31. An outpatient visit, with a mental health provider within 7 and 30 (calendar) days after discharge. Do not include visits that occur on the date of discharge. A visit with a mental health provider in any of the following settings: 	Visit Setting Unspecified: (With Outpatient POS Value Set and with a Mental Health Provider): (with Partial Hospitalization POS): (With Community Mental Health Center POS): (With Telehealth POS Value Set and with a Mental Health Provider): CPT: 90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221, 99222, 99223, 99231, 99232, 99233, 99238, 99239, 99252, 99253, 99254, 99255
	 Two rates are reported: 1. The percentage of discharges for which the member received follow-up within 30 (calendar) days of discharge. 2. The percentage of discharges for which the member received follow-up within 7 (calendar) days of discharge. 	 Outpatient. Behavioral health outpatient. Telehealth visit. Telephone visit. Observation visit. Transitional care management visit. A visit in any of the following settings: Intensive outpatient/partial hospitalization. Community mental 	BH Outpatient: (With a Mental Health Provider): (with Community Mental Health Center POS): CPT: 99078, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99243, 99244, 99245, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99483 HCPCS: G0463, H0004, H0036, H0040, H2000, H2017, H2019, H2020, T1015 UBREV: 0510, 0513, 0515, 0516, 0517, 0519, 0900, 0904, 0911, 0914, 0915, 0916, 0919
		 health center. Electroconvulsive therapy visit. Behavioral healthcare setting. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. 	Partial Hospitalization or Intensive Outpatient: HCPCS: H2012, S9484, S9485 UBREV: 0907, 0912, 0913 BH RCC Only Electroconvulsive Therapy: (with Ambulatory Surgical Center POS): (with Community Mental Health POS): (with Outpatient POS): (with Outpatient POS): (with Partial Hospitalization POS):

- Deceeded at any time in the MAY	CDT: 00070
• Deceased at any time in the MY.	CPT : 90870
	ICD10PCS: GZB0ZZZ, GZB1ZZZ, GZB2ZZZ, GZB3ZZZ, GZB4ZZZ
Common Chart Deficiencies:	
Follow-up visit more than 7 days	
or 30-days after discharge.	Behavioral Healthcare Setting Visit:
• Criteria is not met by a follow-up	UBREV: 0513, 0900, 0901, 0904, 0907, 0911, 0912, 0913,0914, 0915, 0916,
on the date of discharge.	0919
	Telephone Visit:
	(With a Mental Health Provider):
	CPT: 99441, 99442, 99443
	Ambulatory Surgical Center POS: 24
	Telehealth POS: 2
	Mental Illness and Intentional Self-Harm:
	ICD10CM: F20.0, F20.1, F20.2, F20.3, F20.5, F20.81, F20.89, F20.9, F21, F22,
	F23, F24, F25.0, F25.1, F25.8, F25.9, F28, F29, F30.10, F30.11, F30.12, F30.13,
	F30.2, F30.3, F30.4, F30.8, F30.9, F31.0, F31.10, F31.11, F31.12, F31.13, F31.2,
	F31.30, F31.31, F31.32, F31.4, F31.5, F31.60, F31.61, F31.62, F31.63, F31.64,
	F31.70, F31.71, F31.72, F31.73, F31.74, F31.75, F31.76, F31.77, F31.78, F31.81,
	F31.89, F31.9, F32.0, F32.1, F32.2, F32.3, F32.4, F32.5, F32.81, F32.89, F32.9,
	F32.A, F33.0, F33.1, F33.2, F33.3, F33.40, F33.41, F33.42, F33.8, F33.9, F34.0,
	F34.1, F34.81, F34.89, F34.9, F39, F42.2, F42.3, F42.4, F42.8, F42.9, F43.0,
	F43.10, F43.11, F43.12, F43.20, F43.21, F43.22, F43.23, F43.24, F43.25, F43.29,
	F43.8, F43.81, F43.89, F43.9, F44.89, F53.0, F53.1, F60.0, F60.1, F60.2, F60.3,
	F60.4, F60.5, F60.6, F60.7, F60.81, F60.89, F60.9, F63.0, F63.1, F63.2, F63.3,
	F63.81, F63.89, F63.9, F68.10, F68.11, F68.12, F68.13, F68.8, F68.A, F84.0,
	F84.2, F84.3, F84.5, F84.8, F84.9, F90.0, F90.1, F90.2, F90.8, F90.9, F91.0, F91.1,
	F91.2, F91.3, F91.8, F91.9, F93.0, F93.8, F93.9, F94.0, F94.1, F94.2, F94.8, F94.9,
	T14.91XA, T14.91XD, T14.91XS, T36.0X2A, T36.0X2D, T36.0X2S, T36.1X2A,
	T36.1X2D, T36.1X2S, T36.2X2A, T36.2X2D, T36.2X2S, T36.3X2D, T36.3X2D,
	T36.3X2S, T36.4X2A, T36.4X2D, T36.4X2S, T36.5X2A, T36.5X2D, T36.5X2S,
	T36.6X2A, T36.6X2D, T36.6X2S, T36.7X2A, T36.7X2D, T36.7X2S, T36.8X2A,
	T36.8X2D, T36.8X2S, T36.92XA, T36.92XD, T36.92XS, T37.0X2A, T37.0X2D,
	T37.0X2S, T37.1X2A, T37.1X2D, T37.1X2S, T37.2X2A, T37.2X2D, T37.2X2S,
	T37.3X2A, T37.3X2D, T37.3X2S, T37.4X2A, T37.4X2D, T37.4X2S, T37.5X2A,

· · · · · · · · · · · · · · · · · · ·	-
	T37.5X2D, T37.5X2S, T37.8X2A, T37.8X2D, T37.8X2S, T37.92XA, T37.92XD,
	T37.92XS, T38.0X2A, T38.0X2D, T38.0X2S, T38.1X2A, T38.1X2D, T38.1X2S,
	T38.2X2A, T38.2X2D, T38.2X2S, T38.3X2A, T38.3X2D, T38.3X2S, T38.4X2A,
	T38.4X2D, T38.4X2S, T38.5X2A, T38.5X2D, T38.5X2S, T38.6X2A, T38.6X2D,
	T38.6X2S, T38.7X2A, T38.7X2D, T38.7X2S, T38.802A, T38.802D, T38.802S,
	T38.812A, T38.812D, T38.812S, T38.892A, T38.892D, T38.892S, T38.902A,
	T38.902D, T38.902S, T38.992A, T38.992D, T38.992S, T39.012A, T39.012D,
	T39.012S, T39.092A, T39.092D, T39.092S, T39.1X2A, T39.1X2D, T39.1X2S,
	T39.2X2A, T39.2X2D, T39.2X2S, T39.312A, T39.312D, T39.312S, T39.392A,
	T39.392D, T39.392S, T39.4X2A, T39.4X2D, T39.4X2S, T39.8X2A, T39.8X2D,
	T39.8X2S, T39.92XA, T39.92XD, T39.92XS, T40.0X2A, T40.0X2D, T40.0X2S,
	T40.1X2A, T40.1X2D, T40.1X2S, T40.2X2A, T40.2X2D, T40.2X2S, T40.3X2A,
	T40.3X2D, T40.3X2S, T40.412A, T40.412D, T40.412S, T40.422A, T40.422D,
	T40.422S, T40.492A, T40.492D, T40.492S, T40.5X2A, T40.5X2D, T40.5X2S,
	T40.602A, T40.602D, T40.602S, T40.692A, T40.692D, T40.692S, T40.712A,
	T40.712D, T40.712S, T40.722A, T40.722D, T40.722S, T40.8X2A, T40.8X2D,
	T40.8X2S, T40.902A, T40.902D, T40.902S, T40.992A, T40.992D, T40.992S,
	T41.0X2A, T41.0X2D, T41.0X2S, T41.1X2A, T41.1X2D, T41.1X2S, T41.202A,
	T41.202D, T41.202S, T41.292A, T41.292D, T41.292S, T41.3X2A, T41.3X2D,
	T41.3X2S, T41.42XA, T41.42XD, T41.42XS, T41.5X2A, T41.5X2D, T41.5X2S,
	T42.0X2A, T42.0X2D, T42.0X2S, T42.1X2A, T42.1X2D, T42.1X2S, T42.2X2A,
	T42.2X2D, T42.2X2S, T42.3X2A, T42.3X2D, T42.3X2S, T42.4X2A, T42.4X2D,
	T42.4X2S, T42.5X2A, T42.5X2D, T42.5X2S, T42.6X2A, T42.6X2D, T42.6X2S,
	T42.72XA, T42.72XD, T42.72XS, T42.8X2A, T42.8X2D, T42.8X2S, T43.012A,
	T43.012D, T43.012S, T43.022A, T43.022D, T43.022S, T43.1X2A, T43.1X2D,
	T43.1X2S, T43.202A, T43.202D, T43.202S, T43.212A, T43.212D, T43.212S,
	T43.222A, T43.222D, T43.222S, T43.292A, T43.292D, T43.292S, T43.3X2A,
	T43.3X2D, T43.3X2S, T43.4X2A, T43.4X2D, T43.4X2S, T43.502A, T43.502D,
	T43.502S, T43.592A, T43.592D, T43.592S, T43.602A, T43.602D, T43.602S,
	T43.622A, T43.622D, T43.622S, T43.632A, T43.632D, T43.632S, T43.642A,
	T43.642D, T43.642S, T43.652A, T43.652D, T43.652S, T43.692A, T43.692D,
	T43.692S, T43.8X2A, T43.8X2D, T43.8X2S, T43.92XA, T43.92XD, T43.92XS,
	T44.0X2A, T44.0X2D, T44.0X2S, T44.1X2A, T44.1X2D, T44.1X2S, T44.2X2A,
	T44.2X2D, T44.2X2S, T44.3X2A, T44.3X2D, T44.3X2S, T44.4X2A, T44.4X2D,
	T44.4X2S, T44.5X2A, T44.5X2D, T44.5X2S, T44.6X2A, T44.6X2D, T44.6X2S,
	T44.7X2A, T44.7X2D, T44.7X2S, T44.8X2A, T44.8X2D, T44.8X2S, T44.902A,
	T44.902D, T44.902S, T44.992A, T44.992D, T44.992S, T45.0X2A, T45.0X2D,
	T45.0X2S, T45.1X2A, T45.1X2D, T45.1X2S, T45.2X2A, T45.2X2D, T45.2X2S,
	T45.3X2A, T45.3X2D, T45.3X2S, T45.4X2A, T45.4X2D, T45.4X2S, T45.512A,

T45.512D, T45.512S, T45.522A, T45.522D, T45.522S, T45.602A, T45.602D,
T45.602S, T45.612A, T45.612D, T45.612S, T45.622A, T45.622D, T45.622S,
T45.692A, T45.692D, T45.692S, T45.7X2A, T45.7X2D, T45.7X2S, T45.8X2A,
T45.8X2D, T45.8X2S, T45.92XA, T45.92XD, T45.92XS, T46.0X2A, T46.0X2D,
T46.0X2S, T46.1X2A, T46.1X2D, T46.1X2S, T46.2X2A, T46.2X2D, T46.2X2S,
T46.3X2A, T46.3X2D, T46.3X2S, T46.4X2A, T46.4X2D, T46.4X2S, T46.5X2A,
T46.5X2D, T46.5X2S, T46.6X2A, T46.6X2D, T46.6X2S, T46.7X2A, T46.7X2D,
T46.7X2S, T46.8X2A, T46.8X2D, T46.8X2S, T46.902A, T46.902D, T46.902S,
T46.992A, T46.992D, T46.992S, T47.0X2A, T47.0X2D, T47.0X2S, T47.1X2A,
T47.1X2D, T47.1X2S, T47.2X2A, T47.2X2D, T47.2X2S, T47.3X2A, T47.3X2D,
T47.3X2S, T47.4X2A, T47.4X2D, T47.4X2S, T47.5X2A, T47.5X2D, T47.5X2S,
T47.6X2A, T47.6X2D, T47.6X2S, T47.7X2A, T47.7X2D, T47.7X2S, T47.8X2A,
T47.8X2D, T47.8X2S, T47.92XA, T47.92XD, T47.92XS, T48.0X2A, T48.0X2D,
T48.0X2S, T48.1X2A, T48.1X2D, T48.1X2S, T48.202A, T48.202D, T48.202S,
T48.292A, T48.292D, T48.292S, T48.3X2A, T48.3X2D, T48.3X2S, T48.4X2A,
T48.4X2D, T48.4X2S, T48.5X2A, T48.5X2D, T48.5X2S, T48.6X2A, T48.6X2D,
T48.6X2S, T48.902A, T48.902D, T48.902S, T48.992A, T48.992D, T48.992S,
T49.0X2A, T49.0X2D, T49.0X2S, T49.1X2A, T49.1X2D, T49.1X2S, T49.2X2A,
T49.2X2D, T49.2X2S, T49.3X2A, T49.3X2D, T49.3X2S, T49.4X2A, T49.4X2D,
T49.4X2S, T49.5X2A, T49.5X2D, T49.5X2S, T49.6X2A, T49.6X2D, T49.6X2S,
T49.7X2A, T49.7X2D, T49.7X2S, T49.8X2A, T49.8X2D, T49.8X2S, T49.92XA,
T49.92XD, T49.92XS, T50.0X2A, T50.0X2D, T50.0X2S, T50.1X2A, T50.1X2D,
T50.1X2S, T50.2X2A, T50.2X2D, T50.2X2S, T50.3X2A, T50.3X2D, T50.3X2S,
T50.4X2A, T50.4X2D, T50.4X2S, T50.5X2A, T50.5X2D, T50.5X2S, T50.6X2A,
T50.6X2D, T50.6X2S, T50.7X2A, T50.7X2D, T50.7X2S, T50.8X2A, T50.8X2D,
T50.8X2S, T50.902A, T50.902D, T50.902S, T50.912A, T50.912D, T50.912S,
T50.992A, T50.992D, T50.992S, T50.A12A, T50.A12D, T50.A12S, T50.A22A,
T50.A22D, T50.A22S, T50.A92A, T50.A92D, T50.A92S, T50.B12A, T50.B12D,
T50.B12S, T50.B92A, T50.B92D, T50.B92S, T50.Z12A, T50.Z12D, T50.Z12S,
T50.Z92A, T50.Z92D, T50.Z92S, T51.0X2A, T51.0X2D, T51.0X2S, T51.1X2A,
T51.1X2D, T51.1X2S, T51.2X2A, T51.2X2D, T51.2X2S, T51.3X2A, T51.3X2D,
T51.3X2S, T51.8X2A, T51.8X2D, T51.8X2S, T51.92XA, T51.92XD, T51.92XS,
T52.0X2A, T52.0X2D, T52.0X2S, T52.1X2A, T52.1X2D, T52.1X2S, T52.2X2A,
T52.2X2D, T52.2X2S, T52.3X2A, T52.3X2D, T52.3X2S, T52.4X2A, T52.4X2D,
T52.4X2S, T52.8X2A, T52.8X2D, T52.8X2S, T52.92XA, T52.92XD, T52.92XS,
T53.0X2A, T53.0X2D, T53.0X2S, T53.1X2A, T53.1X2D, T53.1X2S, T53.2X2A,
T53.2X2D, T53.2X2S, T53.3X2A, T53.3X2D, T53.3X2S, T53.4X2A, T53.4X2D,
T53.4X2S, T53.5X2A, T53.5X2D, T53.5X2S, T53.6X2A, T53.6X2D, T53.6X2S,
T53.7X2A, T53.7X2D, T53.7X2S, T53.92XA, T53.92XD, T53.92XS, T54.0X2A,

T54.0X2D, T54.0X2S, T54.1X2A, T54.1X2D, T54.1X2S, T54.2X2A, T54.2X2D,
T54.2X2S, T54.3X2A, T54.3X2D, T54.3X2S, T54.92XA, T54.92XD, T54.92XS,
T55.0X2A, T55.0X2D, T55.0X2S, T55.1X2A, T55.1X2D, T55.1X2S, T56.0X2A,
T56.0X2D, T56.0X2S, T56.1X2A, T56.1X2D, T56.1X2S, T56.2X2A, T56.2X2D,
T56.2X2S, T56.3X2A, T56.3X2D, T56.3X2S, T56.4X2A, T56.4X2D, T56.4X2S,
T56.5X2A, T56.5X2D, T56.5X2S, T56.6X2A, T56.6X2D, T56.6X2S, T56.7X2A,
T56.7X2D, T56.7X2S, T56.812A, T56.812D, T56.812S, T56.892A, T56.892D,
T56.892S, T56.92XA, T56.92XD, T56.92XS, T57.0X2A, T57.0X2D, T57.0X2S,
T57.1X2A, T57.1X2D, T57.1X2S, T57.2X2A, T57.2X2D, T57.2X2S, T57.3X2A,
T57.3X2D, T57.3X2S, T57.8X2A, T57.8X2D, T57.8X2S, T57.92XA, T57.92XD,
T57.92XS, T58.02XA, T58.02XD, T58.02XS, T58.12XA, T58.12XD, T58.12XS,
T58.2X2A, T58.2X2D, T58.2X2S, T58.8X2A, T58.8X2D, T58.8X2S, T58.92XA,
T58.92XD, T58.92XS, T59.0X2A, T59.0X2D, T59.0X2S, T59.1X2A, T59.1X2D,
T59.1X2S, T59.2X2A, T59.2X2D, T59.2X2S, T59.3X2A, T59.3X2D, T59.3X2S,
T59.4X2A, T59.4X2D, T59.4X2S, T59.5X2A, T59.5X2D, T59.5X2S, T59.6X2A,
T59.6X2D, T59.6X2S, T59.7X2A, T59.7X2D, T59.7X2S, T59.812A, T59.812D,
T59.812S, T59.892A, T59.892D, T59.892S, T59.92XA, T59.92XD, T59.92XS,
T60.0X2A, T60.0X2D, T60.0X2S, T60.1X2A, T60.1X2D, T60.1X2S, T60.2X2A,
T60.2X2D, T60.2X2S, T60.3X2A, T60.3X2D, T60.3X2S, T60.4X2A, T60.4X2D,
T60.4X2S, T60.8X2A, T60.8X2D, T60.8X2S, T60.92XA, T60.92XD, T60.92XS,
T61.02XA, T61.02XD, T61.02XS, T61.12XA, T61.12XD, T61.12XS, T61.772A,
T61.772D, T61.772S, T61.782A, T61.782D, T61.782S, T61.8X2A, T61.8X2D,
T61.8X2S, T61.92XA, T61.92XD, T61.92XS, T62.0X2A, T62.0X2D, T62.0X2S,
T62.1X2A, T62.1X2D, T62.1X2S, T62.2X2A, T62.2X2D, T62.2X2S, T62.8X2A,
T62.8X2D, T62.8X2S, T62.92XA, T62.92XD, T62.92XS, T63.002A, T63.002D,
T63.002S, T63.012A, T63.012D, T63.012S, T63.022A, T63.022D, T63.022S,
T63.032A, T63.032D, T63.032S, T63.042A, T63.042D, T63.042S, T63.062A,
T63.062D, T63.062S, T63.072A, T63.072D, T63.072S, T63.082A, T63.082D,
T63.082S, T63.092A, T63.092D, T63.092S, T63.112A, T63.112D, T63.112S,
T63.122A, T63.122D, T63.122S, T63.192A, T63.192D, T63.192S, T63.2X2A,
T63.2X2D, T63.2X2S, T63.302A, T63.302D, T63.302S, T63.312A, T63.312D,
T63.312S, T63.322A, T63.322D, T63.322S, T63.332A, T63.332D, T63.332S,
T63.392A, T63.392D, T63.392S, T63.412A, T63.412D, T63.412S, T63.422A,
T63.422D, T63.422S, T63.432A, T63.432D, T63.432S, T63.442A, T63.442D,
T63.442S, T63.452A, T63.452D, T63.452S, T63.462A, T63.462D, T63.462S,
T63.482A, T63.482D, T63.482S, T63.512A, T63.512D, T63.512S, T63.592A,
T63.592D, T63.592S, T63.612A, T63.612D, T63.612S, T63.622A, T63.622D,
T63.622S, T63.632A, T63.632D, T63.632S, T63.692A, T63.692D, T63.692S,
T63.712A, T63.712D, T63.712S, T63.792A, T63.792D, T63.792S, T63.812A,

T63.812D, T63.812S, T63.822A, T63.822D, T63.822S, T63.832A, T63.832D,
T63.832S, T63.892A, T63.892D, T63.892S, T63.92XA, T63.92XD, T63.92XS,
T64.02XA, T64.02XD, T64.02XS, T64.82XA, T64.82XD, T64.82XS, T65.0X2A,
T65.0X2D, T65.0X2S, T65.1X2A, T65.1X2D, T65.1X2S, T65.212A, T65.212D,
T65.212S, T65.222A, T65.222D, T65.222S, T65.292A, T65.292D, T65.292S,
T65.3X2A, T65.3X2D, T65.3X2S, T65.4X2A, T65.4X2D, T65.4X2S, T65.5X2A,
T65.5X2D, T65.5X2S, T65.6X2A, T65.6X2D, T65.6X2S, T65.812A, T65.812D,
T65.812S, T65.822A, T65.822D, T65.822S, T65.832A, T65.832D, T65.832S,
T65.892A, T65.892D, T65.892S, T65.92XA, T65.92XD, T65.92XS, T71.112A,
T71.112D, T71.112S, T71.122A, T71.122D, T71.122S, T71.132A, T71.132D,
T71.132S, T71.152A, T71.152D, T71.152S, T71.162A, T71.162D, T71.162S,
T71.192A, T71.192D, T71.192S, T71.222A, T71.222D, T71.222S, T71.232A,
T71.232D, T71.232S
Intentional Self-Harm Diagnosis:
ICD10CM: T14.91XA, T14.91XD, T14.91XS, T36.0X2A, T36.0X2D, T36.0X2S,
T36.1X2A, T36.1X2D, T36.1X2S, T36.2X2A, T36.2X2D, T36.2X2S, T36.3X2A,
T36.3X2D, T36.3X2S, T36.4X2A, T36.4X2D, T36.4X2S, T36.5X2A, T36.5X2D,
T36.5X2S, T36.6X2A, T36.6X2D, T36.6X2S, T36.7X2A, T36.7X2D, T36.7X2S,
T36.8X2A, T36.8X2D, T36.8X2S, T36.92XA, T36.92XD, T36.92XS, T37.0X2A,
T37.0X2D, T37.0X2S, T37.1X2A, T37.1X2D, T37.1X2S, T37.2X2A, T37.2X2D,
T37.2X2S, T37.3X2A, T37.3X2D, T37.3X2S, T37.4X2A, T37.4X2D, T37.4X2S,
T37.5X2A, T37.5X2D, T37.5X2S, T37.8X2A, T37.8X2D, T37.8X2S, T37.92XA,
T37.92XD, T37.92XS, T38.0X2A, T38.0X2D, T38.0X2S, T38.1X2A, T38.1X2D,
T38.1X2S, T38.2X2A, T38.2X2D, T38.2X2S, T38.3X2A, T38.3X2D, T38.3X2S,
T38.4X2A, T38.4X2D, T38.4X2S, T38.5X2A, T38.5X2D, T38.5X2S, T38.6X2A,
T38.6X2D, T38.6X2S, T38.7X2A, T38.7X2D, T38.7X2S, T38.802A, T38.802D,
T38.802S, T38.812A, T38.812D, T38.812S, T38.892A, T38.892D, T38.892S,
T38.902A, T38.902D, T38.902S, T38.992A, T38.992D, T38.992S, T39.012A,
T39.012D, T39.012S, T39.092A, T39.092D, T39.092S, T39.1X2A, T39.1X2D,
T39.1X2S, T39.2X2A, T39.2X2D, T39.2X2S, T39.312A, T39.312D, T39.312S,
T39.392A, T39.392D, T39.392S, T39.4X2A, T39.4X2D, T39.4X2S, T39.8X2A,
T39.8X2D, T39.8X2S, T39.92XA, T39.92XD, T39.92XS, T40.0X2A, T40.0X2D,
T40.0X2S, T40.1X2A, T40.1X2D, T40.1X2S, T40.2X2A, T40.2X2D, T40.2X2S,
T40.3X2A, T40.3X2D, T40.3X2S, T40.412A, T40.412D, T40.412S, T40.422A,
T40.422D, T40.422S, T40.492A, T40.492D, T40.492S, T40.5X2A, T40.5X2D,
T40.5X2S, T40.602A, T40.602D, T40.602S, T40.692A, T40.692D, T40.692S,
T40.712A, T40.712D, T40.712S, T40.722A, T40.722D, T40.722S, T40.8X2A,
T40.8X2D, T40.8X2S, T40.902A, T40.902D, T40.902S, T40.992A, T40.992D,

T40.992S, T41.0X2A, T41.0X2D, T41.0X2S, T41.1X2A, T41.1X2D, T41.1X2S,
T41.202A, T41.202D, T41.202S, T41.292A, T41.292D, T41.292S, T41.3X2A,
T41.3X2D, T41.3X2S, T41.42XA, T41.42XD, T41.42XS, T41.5X2A, T41.5X2D,
T41.5X2S, T42.0X2A, T42.0X2D, T42.0X2S, T42.1X2A, T42.1X2D, T42.1X2S,
T42.2X2A, T42.2X2D, T42.2X2S, T42.3X2A, T42.3X2D, T42.3X2S, T42.4X2A,
T42.4X2D, T42.4X2S, T42.5X2A, T42.5X2D, T42.5X2S, T42.6X2A, T42.6X2D,
T42.6X2S, T42.72XA, T42.72XD, T42.72XS, T42.8X2A, T42.8X2D, T42.8X2S,
T43.012A, T43.012D, T43.012S, T43.022A, T43.022D, T43.022S, T43.1X2A,
T43.1X2D, T43.1X2S, T43.202A, T43.202D, T43.202S, T43.212A, T43.212D,
T43.212S, T43.222A, T43.222D, T43.222S, T43.292A, T43.292D, T43.292S,
T43.3X2A, T43.3X2D, T43.3X2S, T43.4X2A, T43.4X2D, T43.4X2S, T43.502A,
T43.502D, T43.502S, T43.592A, T43.592D, T43.592S, T43.602A, T43.602D,
T43.602S, T43.622A, T43.622D, T43.622S, T43.632A, T43.632D, T43.632S,
T43.642A, T43.642D, T43.642S, T43.652A, T43.652D, T43.652S, T43.692A,
T43.692D, T43.692S, T43.8X2A, T43.8X2D, T43.8X2S, T43.92XA, T43.92XD,
T43.92XS, T44.0X2A, T44.0X2D, T44.0X2S, T44.1X2A, T44.1X2D, T44.1X2S,
T44.2X2A, T44.2X2D, T44.2X2S, T44.3X2A, T44.3X2D, T44.3X2S, T44.4X2A,
T44.4X2D, T44.4X2S, T44.5X2A, T44.5X2D, T44.5X2S, T44.6X2A, T44.6X2D,
T44.6X2S, T44.7X2A, T44.7X2D, T44.7X2S, T44.8X2A, T44.8X2D, T44.8X2S,
T44.902A, T44.902D, T44.902S, T44.992A, T44.992D, T44.992S, T45.0X2A,
T45.0X2D, T45.0X2S, T45.1X2A, T45.1X2D, T45.1X2S, T45.2X2A, T45.2X2D,
T45.2X2S, T45.3X2A, T45.3X2D, T45.3X2S, T45.4X2A, T45.4X2D, T45.4X2S,
T45.512A, T45.512D, T45.512S, T45.522A, T45.522D, T45.522S, T45.602A,
T45.602D, T45.602S, T45.612A, T45.612D, T45.612S, T45.622A, T45.622D,
T45.622S, T45.692A, T45.692D, T45.692S, T45.7X2A, T45.7X2D, T45.7X2S,
T45.8X2A, T45.8X2D, T45.8X2S, T45.92XA, T45.92XD, T45.92XS, T46.0X2A,
T46.0X2D, T46.0X2S, T46.1X2A, T46.1X2D, T46.1X2S, T46.2X2A, T46.2X2D,
T46.2X2S, T46.3X2A, T46.3X2D, T46.3X2S, T46.4X2A, T46.4X2D, T46.4X2S,
T46.5X2A, T46.5X2D, T46.5X2S, T46.6X2A, T46.6X2D, T46.6X2S, T46.7X2A,
T46.7X2D, T46.7X2S, T46.8X2A, T46.8X2D, T46.8X2S, T46.902A, T46.902D,
T46.902S, T46.992A, T46.992D, T46.992S, T47.0X2A, T47.0X2D, T47.0X2S,
T47.1X2A, T47.1X2D, T47.1X2S, T47.2X2A, T47.2X2D, T47.2X2S, T47.3X2A,
T47.3X2D, T47.3X2S, T47.4X2A, T47.4X2D, T47.4X2S, T47.5X2A, T47.5X2D,
T47.5X2S, T47.6X2A, T47.6X2D, T47.6X2S, T47.7X2A, T47.7X2D, T47.7X2S,
T47.8X2A, T47.8X2D, T47.8X2S, T47.92XA, T47.92XD, T47.92XS, T48.0X2A,
T48.0X2D, T48.0X2S, T48.1X2A, T48.1X2D, T48.1X2S, T48.202A, T48.202D,
T48.202S, T48.292A, T48.292D, T48.292S, T48.3X2A, T48.3X2D, T48.3X2S,
T48.4X2A, T48.4X2D, T48.4X2S, T48.5X2A, T48.5X2D, T48.5X2S, T48.6X2A,
T48.6X2D, T48.6X2S, T48.902A, T48.902D, T48.902S, T48.992A, T48.992D,

T48.992S, T49.0X2A, T49.0X2D, T49.0X2S, T49.1X2A, T49.1X2D, T49.1X2S,
T49.2X2A, T49.2X2D, T49.2X2S, T49.3X2A, T49.3X2D, T49.3X2S, T49.4X2A,
T49.4X2D, T49.4X2S, T49.5X2A, T49.5X2D, T49.5X2S, T49.6X2A, T49.6X2D,
T49.6X2S, T49.7X2A, T49.7X2D, T49.7X2S, T49.8X2A, T49.8X2D, T49.8X2S,
T49.92XA, T49.92XD, T49.92XS, T50.0X2A, T50.0X2D, T50.0X2S, T50.1X2A,
T50.1X2D, T50.1X2S, T50.2X2A, T50.2X2D, T50.2X2S, T50.3X2A, T50.3X2D,
T50.3X2S, T50.4X2A, T50.4X2D, T50.4X2S, T50.5X2A, T50.5X2D, T50.5X2S,
T50.6X2A, T50.6X2D, T50.6X2S, T50.7X2A, T50.7X2D, T50.7X2S, T50.8X2A,
T50.8X2D, T50.8X2S, T50.902A, T50.902D, T50.902S, T50.912A, T50.912D,
T50.912S, T50.992A, T50.992D, T50.992S, T50.A12A, T50.A12D, T50.A12S,
T50.A22A, T50.A22D, T50.A22S, T50.A92A, T50.A92D, T50.A92S, T50.B12A,
T50.B12D, T50.B12S, T50.B92A, T50.B92D, T50.B92S, T50.Z12A, T50.Z12D,
T50.Z12S, T50.Z92A, T50.Z92D, T50.Z92S, T51.0X2A, T51.0X2D, T51.0X2S,
T51.1X2A, T51.1X2D, T51.1X2S, T51.2X2A, T51.2X2D, T51.2X2S, T51.3X2A,
T51.3X2D, T51.3X2S, T51.8X2A, T51.8X2D, T51.8X2S, T51.92XA, T51.92XD,
T51.92XS, T52.0X2A, T52.0X2D, T52.0X2S, T52.1X2A, T52.1X2D, T52.1X2S,
T52.2X2A, T52.2X2D, T52.2X2S, T52.3X2A, T52.3X2D, T52.3X2S, T52.4X2A,
T52.4X2D, T52.4X2S, T52.8X2A, T52.8X2D, T52.8X2S, T52.92XA, T52.92XD,
T52.92XS, T53.0X2A, T53.0X2D, T53.0X2S, T53.1X2A, T53.1X2D, T53.1X2S,
T53.2X2A, T53.2X2D, T53.2X2S, T53.3X2A, T53.3X2D, T53.3X2S, T53.4X2A,
T53.4X2D, T53.4X2S, T53.5X2A, T53.5X2D, T53.5X2S, T53.6X2A, T53.6X2D,
T53.6X2S, T53.7X2A, T53.7X2D, T53.7X2S, T53.92XA, T53.92XD, T53.92XS,
T54.0X2A, T54.0X2D, T54.0X2S, T54.1X2A, T54.1X2D, T54.1X2S, T54.2X2A,
T54.2X2D, T54.2X2S, T54.3X2A, T54.3X2D, T54.3X2S, T54.92XA, T54.92XD,
T54.92XS, T55.0X2A, T55.0X2D, T55.0X2S, T55.1X2A, T55.1X2D, T55.1X2S,
T56.0X2A, T56.0X2D, T56.0X2S, T56.1X2A, T56.1X2D, T56.1X2S, T56.2X2A,
T56.2X2D, T56.2X2S, T56.3X2A, T56.3X2D, T56.3X2S, T56.4X2A, T56.4X2D,
T56.4X2S, T56.5X2A, T56.5X2D, T56.5X2S, T56.6X2A, T56.6X2D, T56.6X2S,
T56.7X2A, T56.7X2D, T56.7X2S, T56.812A, T56.812D, T56.812S, T56.892A,
T56.892D, T56.892S, T56.92XA, T56.92XD, T56.92XS, T57.0X2A, T57.0X2D,
T57.0X2S, T57.1X2A, T57.1X2D, T57.1X2S, T57.2X2A, T57.2X2D, T57.2X2S,
T57.3X2A, T57.3X2D, T57.3X2S, T57.8X2A, T57.8X2D, T57.8X2S, T57.92XA,
T57.92XD, T57.92XS, T58.02XA, T58.02XD, T58.02XS, T58.12XA, T58.12XD,
T58.12XS, T58.2X2A, T58.2X2D, T58.2X2S, T58.8X2A, T58.8X2D, T58.8X2S,
T58.92XA, T58.92XD, T58.92XS, T59.0X2A, T59.0X2D, T59.0X2S, T59.1X2A,
T59.1X2D, T59.1X2S, T59.2X2A, T59.2X2D, T59.2X2S, T59.3X2A, T59.3X2D,
T59.3X2S, T59.4X2A, T59.4X2D, T59.4X2S, T59.5X2A, T59.5X2D, T59.5X2S,
T59.6X2A, T59.6X2D, T59.6X2S, T59.7X2A, T59.7X2D, T59.7X2S, T59.812A,
T59.812D, T59.812S, T59.892A, T59.892D, T59.892S, T59.92XA, T59.92XD,

T59.92XS, T60.0X2A, T60.0X2D, T60.0X2S, T60.1X2A, T60.1X2D, T60.1X2S,
T60.2X2A, T60.2X2D, T60.2X2S, T60.3X2A, T60.3X2D, T60.3X2S, T60.4X2A,
T60.4X2D, T60.4X2S, T60.8X2A, T60.8X2D, T60.8X2S, T60.92XA, T60.92XD,
T60.92XS, T61.02XA, T61.02XD, T61.02XS, T61.12XA, T61.12XD, T61.12XS,
T61.772A, T61.772D, T61.772S, T61.782A, T61.782D, T61.782S, T61.8X2A,
T61.8X2D, T61.8X2S, T61.92XA, T61.92XD, T61.92XS, T62.0X2A, T62.0X2D,
T62.0X2S, T62.1X2A, T62.1X2D, T62.1X2S, T62.2X2A, T62.2X2D, T62.2X2S,
T62.8X2A, T62.8X2D, T62.8X2S, T62.92XA, T62.92XD, T62.92XS, T63.002A,
T63.002D, T63.002S, T63.012A, T63.012D, T63.012S, T63.022A, T63.022D,
T63.022S, T63.032A, T63.032D, T63.032S, T63.042A, T63.042D, T63.042S,
T63.062A, T63.062D, T63.062S, T63.072A, T63.072D, T63.072S, T63.082A,
T63.082D, T63.082S, T63.092A, T63.092D, T63.092S, T63.112A, T63.112D,
T63.112S, T63.122A, T63.122D, T63.122S, T63.192A, T63.192D, T63.192S,
T63.2X2A, T63.2X2D, T63.2X2S, T63.302A, T63.302D, T63.302S, T63.312A,
T63.312D, T63.312S, T63.322A, T63.322D, T63.322S, T63.332A, T63.332D,
T63.332S, T63.392A, T63.392D, T63.392S, T63.412A, T63.412D, T63.412S,
T63.422A, T63.422D, T63.422S, T63.432A, T63.432D, T63.432S, T63.442A,
T63.442D, T63.442S, T63.452A, T63.452D, T63.452S, T63.462A, T63.462D,
T63.462S, T63.482A, T63.482D, T63.482S, T63.512A, T63.512D, T63.512S,
T63.592A, T63.592D, T63.592S, T63.612A, T63.612D, T63.612S, T63.622A,
T63.622D, T63.622S, T63.632A, T63.632D, T63.632S, T63.692A, T63.692D,
T63.692S, T63.712A, T63.712D, T63.712S, T63.792A, T63.792D, T63.792S,
T63.812A, T63.812D, T63.812S, T63.822A, T63.822D, T63.822S, T63.832A,
T63.832D, T63.832S, T63.892A, T63.892D, T63.892S, T63.92XA, T63.92XD,
T63.92XS, T64.02XA, T64.02XD, T64.02XS, T64.82XA, T64.82XD, T64.82XS,
T65.0X2A, T65.0X2D, T65.0X2S, T65.1X2A, T65.1X2D, T65.1X2S, T65.212A,
T65.212D, T65.212S, T65.222A, T65.222D, T65.222S, T65.292A, T65.292D,
T65.292S, T65.3X2A, T65.3X2D, T65.3X2S, T65.4X2A, T65.4X2D, T65.4X2S,
T65.5X2A, T65.5X2D, T65.5X2S, T65.6X2A, T65.6X2D, T65.6X2S, T65.812A,
T65.812D, T65.812S, T65.822A, T65.822D, T65.822S, T65.832A, T65.832D,
T65.832S, T65.892A, T65.892D, T65.892S, T65.92XA, T65.92XD, T65.92XS,
T71.112A, T71.112D, T71.112S, T71.122A, T71.122D, T71.122S, T71.132A,
T71.132D, T71.132S, T71.152A, T71.152D, T71.152S, T71.162A, T71.162D,
T71.162S, T71.192A, T71.192D, T71.192S, T71.222A, T71.222D, T71.222S,
T71.232A, T71.232D, T71.232S
Note: LOINC and SNOMED codes can be captured through electronic data
submissions. Please contact your Account Executive for more information.

Measure	Measure Description	Measure Information/Documentation Required	Coding
Diabetes Screening for People with Schizophrenia or Bipolar Disorder Who are Using Antipsychotic Medications (SSD)	The percentage of members 18 – 64 years of age with schizophrenia, schizoaffective disorder, or bipolar disorder who were dispensed an antipsychotic medication and had a diabetes screening test during the MY.	A glucose test or HbA1c test performed during the MY. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. Diabetes.	 Members are identified through administrative and pharmacy claims. Glucose Lab Test: CPT: 80047, 80048, 80053, 80069, 82947, 82950, 82951 HbA1C Lab Test: CPT: 83036, 83037 HbA1C Test Result or Finding: CPT-CAT-II: 3044F, 3046F, 3051F, 3052F Antipsychotics Medications: Miscellaneous antipsychotic agents: Aripiprazole, Asenapine, Brexpiprazole, Cariprazine, Clozapine, Haloperidol, Iloperidone, Loxapine, Lumateperone, Lurasidone, Molindone, Olanzapine, Paliperidone, Quetiapine, Risperidone, Ziprasidone Phenothiazine antipsychotics: Chlorpromazine, Fluphenazine, Perphenazine, Prochlorperazine, Thioridazine, Trifluoperazine Psychotherapeutic combinations: Amitriptyline-perphenazine Thioxanthenes: Thiothixene Long-acting injections: Aripiprazole, Fluphenazine decanoate, Haloperidol decanoate, Olanzapine, Paliperidone palmitate, Risperidone Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
Measure	Measure Description	Measure Information/Documentation Required	Coding
Diabetes Monitoring for People with Diabetes and Schizophrenia (SMD)	The percentage of members 18 – 64 years of age with schizophrenia or schizoaffective disorder,	An HbA1c test and an LDL-C test performed in the MY. Required Exclusions:	Members are identified through administrative and pharmacy claims. HbA1C Lab Test: CPT: 83036, 83037

	and diabetes who had both an LDL-C test and an HbA1c test during the MY.	 Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. 	HbA1C Test Result or Finding: CPT-CAT-II: 3044F, 3046F, 3051F, 3052FLDL-C Lab Test: CPT: 80061, 83700, 83701, 83704, 83721LDL-C Test Result or Finding: CPT-CAT-II: 3048F, 3049F, 3050FMust have both A1c and LDL.Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
Measure	Measure Description	Measure Information/Documentation Required	Coding
Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC)	The percentage of members 18 – 64 years of age with schizophrenia or schizoaffective disorder and cardiovascular (IVD, CABG, PCI, AMI) disease who had an LDL-C test during the MY.	 An LDL-C test performed during the MY. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. 	LDL-C Lab Test: CPT: 80061, 83700, 83701, 83704, 83721 LDL-C Test Result or Finding: CPT-CAT-II: 3048F, 3049F, 3050F Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
Measure	Measure Description	Measure Information/Documentation Required	Coding
Adherence to Antipsychotic Medications for Individuals with Schizophrenia (SAA)	The percentage of members 18 years of age and older during the MY with schizophrenia or schizoaffective disorder who were dispensed and remained	The Index Prescription Start Date (ISPD) is the earliest prescription- dispensing date during the MY. The Treatment period is the ISPD through the last day of the MY. Required Exclusions:	Schizophrenia Diagnosis: ICD10CM: F20.0, F20.1, F20.2, F20.3, F20.5, F20.81, F20.89, F20.9, F25.0, F25.1, F25.8, F25.9 Long-Acting Injections 28-Day Supply: HCPCS: J0401, J1631, J1943, J1944, J2358, J2426, J2680

	on an oral or long-acting injection antipsychotic medication at least 80% of their treatment period.	 Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. 66 – 80 years of age with frailty and advanced illness during the MY. 81 years of age and older with frailty. Diagnosis of dementia in the MY. 	 Oral Antipsychotic Medications: Miscellaneous antipsychotic agents: Aripiprazole, Asenapine, Brexpiprazole, Cariprazine, Clozapine, Haloperidol, Iloperidone, Loxapine, Lumateperone, Lurasidone, Molindone, Olanzapine, Paliperidone, Quetiapine, Risperidone, Ziprasidone, Phenothiazine antipsychotics: Chlorpromazine, Fluphenazine, Perphenazine, Prochlorperazine, Thioridazine, Trifluoperazine Psychotherapeutic combinations: Amitriptyline-perphenazine Thioxanthenes: Thiothixene Long-Acting Injections: 28-day supply: Aripiprazole, Fluphenazine decanoate, Haloperidol decanoate, Olanzapine, Paliperidone Palmitate Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
Measure	Measure Description	Measure Information/Documentation Required	Coding
Follow-Up After Emergency Department Visit for Mental Illness (FUM)	The percentage of emergency department (ED) visits for members 6 years of age and older with a principal diagnosis of mental illness or intentional self-harm, who had a follow-up visit for mental illness. Two rates are reported: 1. The percentage of ED visits for which the member received follow-up within 30 days of	 A follow-up visit with any practitioner, with a principal diagnosis of a mental health disorder or with a principal diagnosis of intentional self-harm and any diagnosis of a mental health disorder within 7 and 30 days after ED visit. Include outpatient visits, behavioral health outpatient visits, intensive outpatient visits, partial hospitalizations, community mental health visits, electroconvulsive therapy visits, telehealth visits, and observation visits. Includes visits that occur on the date of the ED visit Telephone visits, e-visits, and virtual check-ins are acceptable. 	 Visit Setting Unspecified: (With Outpatient POS and Principal Diagnosis of Mental Health or Principal Diagnosis of Intentional Self-Harm with Any Diagnosis of Mental Health): (With Partial Hospitalization POS and Principal Diagnosis of Mental Health or Principal Diagnosis of Intentional Self-Harm with any Diagnosis of Mental Health): (With Community Mental Health Center POS and Principal Diagnosis of Mental Health): (With Community Mental Health Center POS and Principal Diagnosis of Mental Health or Principal Diagnosis of Intentional Self-Harm with Any Diagnosis of Mental Health or Principal Diagnosis of Intentional Self-Harm with Any Diagnosis of Mental Health): (With Telehealth POS and Principal Diagnosis of Mental Health or Principal Diagnosis of Intentional Self-Harm with Any Diagnosis of Mental Health): (With Telehealth POS and Principal Diagnosis of Mental Health): CPT: 90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221, 99222, 99233, 99231, 99232, 99233, 99238, 99239, 99252, 99253, 99254, 99255

Г	the ED vieit (24 total	Deguized Evolution	PUL Outpatient
	the ED visit (31 total	Required Exclusions:	BH Outpatient:
	days).	Members who meet any of the	(With Principal Diagnosis of Mental Health or Principal Diagnosis of Intentional
	2. The percentage of ED visits for which	following criteria are excluded from	Self-Harm with Any Diagnosis of Mental Health):
	the member received follow-up within 7 days of the ED visit (8 total days).	 the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. 	CPT: 99078, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99243, 99244, 99245, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99483
			HCPCS: G0463, H0004, H0036, H0040, H2000, H2017, H2019, H2020, T1015
			UBREV: 0510, 0513, 0515, 0516, 0517, 0519, 0900, 0904, 0911, 0914, 0915, 0916, 0919
			Partial Hospitalization or Intensive Outpatient: (With Principal Diagnosis of Mental Health or Principal Diagnosis of Intentional
			Self-Harm with Any Diagnosis of Mental Health of Principal Diagnosis of Interitional HCPCS: H2012, S9484, S9485
			UBREV: 0907, 0912, 0913 only covered in BH RCC
			Electroconvulsive Therapy: (With Ambulatory Surgical Center POS, Community Mental Health POS, Outpatient POS, or Partial Hospitalization POS and Principal Diagnosis of Mental Health or Principal Diagnosis of Intentional Self-Harm with Any Diagnosis of Mental Health): CPT: 99381, 99382, 99391, 99392
			Observation: (with Principal Diagnosis of Mental Health or Principal Diagnosis of Intentional Self-Harm with Any Diagnosis of Mental Health): CPT: 99217, 99218, 99219, 99220
			Telephone Visits: (with Principal Diagnosis of Mental Health or Principal Diagnosis of Intentional Self-Harm with Any Diagnosis of Mental Health): CPT: 99441, 99442, 99443

Online Assessments:
(With Principal Diagnosis of Mental Health or Principal Diagnosis of Intentional
Self-Harm with Any Diagnosis of Mental Health):
CPT: 99421, 99422, 99423, 99457, 99458
HCPCS: G2010, G2012, G2251,
Ambulatory Surgical Center POS: 24
Outpatient POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72
Telehealth POS: 2
Mental Illness and Intentional Self-Harm:
ICD10CM: F20.0, F20.1, F20.2, F20.3, F20.5, F20.81, F20.89, F20.9, F21, F22,
F23, F24, F25.0, F25.1, F25.8, F25.9, F28, F29, F30.10, F30.11, F30.12, F30.13,
F30.2, F30.3, F30.4, F30.8, F30.9, F31.0, F31.10, F31.11, F31.12, F31.13, F31.2,
F31.30, F31.31, F31.32, F31.4, F31.5, F31.60, F31.61, F31.62, F31.63, F31.64,
F31.70, F31.71, F31.72, F31.73, F31.74, F31.75, F31.76, F31.77, F31.78, F31.81,
F31.89, F31.9, F32.0, F32.1, F32.2, F32.3, F32.4, F32.5, F32.81, F32.89, F32.9,
F32.A, F33.0, F33.1, F33.2, F33.3, F33.40, F33.41, F33.42, F33.8, F33.9, F34.0,
F34.1, F34.81, F34.89, F34.9, F39, F42.2, F42.3, F42.4, F42.8, F42.9, F43.0,
F43.10, F43.11, F43.12, F43.20, F43.21, F43.22, F43.23, F43.24, F43.25, F43.29,
F43.8, F43.81, F43.89, F43.9, F44.89, F53.0, F53.1, F60.0, F60.1, F60.2, F60.3,
F60.4, F60.5, F60.6, F60.7, F60.81, F60.89, F60.9, F63.0, F63.1, F63.2, F63.3,
F63.81, F63.89, F63.9, F68.10, F68.11, F68.12, F68.13, F68.8, F68.A, F84.0,
F84.2, F84.3, F84.5, F84.8, F84.9, F90.0, F90.1, F90.2, F90.8, F90.9, F91.0, F91.1,
F91.2, F91.3, F91.8, F91.9, F93.0, F93.8, F93.9, F94.0, F94.1, F94.2, F94.8, F94.9,
T14.91XA, T14.91XD, T14.91XS, T36.0X2A, T36.0X2D, T36.0X2S, T36.1X2A,
T36.1X2D, T36.1X2S, T36.2X2A, T36.2X2D, T36.2X2S, T36.3X2A, T36.3X2D,
T36.3X2S, T36.4X2A, T36.4X2D, T36.4X2S, T36.5X2A, T36.5X2D, T36.5X2S,
T36.6X2A, T36.6X2D, T36.6X2S, T36.7X2A, T36.7X2D, T36.7X2S, T36.8X2A,
T36.8X2D, T36.8X2S, T36.92XA, T36.92XD, T36.92XS, T37.0X2A, T37.0X2D,
T37.0X2S, T37.1X2A, T37.1X2D, T37.1X2S, T37.2X2A, T37.2X2D, T37.2X2S,
T37.3X2A, T37.3X2D, T37.3X2S, T37.4X2A, T37.4X2D, T37.4X2S, T37.5X2A,
T37.5X2D, T37.5X2S, T37.8X2A, T37.8X2D, T37.8X2S, T37.92XA, T37.92XD,
T37.92XS, T38.0X2A, T38.0X2D, T38.0X2S, T38.1X2A, T38.1X2D, T38.1X2S,
T38.2X2A, T38.2X2D, T38.2X2S, T38.3X2A, T38.3X2D, T38.3X2S, T38.4X2A,
T38.4X2D, T38.4X2S, T38.5X2A, T38.5X2D, T38.5X2S, T38.6X2A, T38.6X2D,

T38.6X2S, T38.7X2A, T38.7X2D, T38.7X2S, T38.802A, T38.802D, T38.802S,
T38.812A, T38.812D, T38.812S, T38.892A, T38.892D, T38.892S, T38.902A,
T38.902D, T38.902S, T38.992A, T38.992D, T38.992S, T39.012A, T39.012D,
T39.012S, T39.092A, T39.092D, T39.092S, T39.1X2A, T39.1X2D, T39.1X2S,
T39.2X2A, T39.2X2D, T39.2X2S, T39.312A, T39.312D, T39.312S, T39.392A,
T39.392D, T39.392S, T39.4X2A, T39.4X2D, T39.4X2S, T39.8X2A, T39.8X2D,
T39.8X2S, T39.92XA, T39.92XD, T39.92XS, T40.0X2A, T40.0X2D, T40.0X2S,
T40.1X2A, T40.1X2D, T40.1X2S, T40.2X2A, T40.2X2D, T40.2X2S, T40.3X2A,
T40.3X2D, T40.3X2S, T40.412A, T40.412D, T40.412S, T40.422A, T40.422D,
T40.422S, T40.492A, T40.492D, T40.492S, T40.5X2A, T40.5X2D, T40.5X2S,
T40.602A, T40.602D, T40.602S, T40.692A, T40.692D, T40.692S, T40.712A,
T40.712D, T40.712S, T40.722A, T40.722D, T40.722S, T40.8X2A, T40.8X2D,
T40.8X2S, T40.902A, T40.902D, T40.902S, T40.992A, T40.992D, T40.992S,
T41.0X2A, T41.0X2D, T41.0X2S, T41.1X2A, T41.1X2D, T41.1X2S, T41.202A,
T41.202D, T41.202S, T41.292A, T41.292D, T41.292S, T41.3X2A, T41.3X2D,
T41.3X2S, T41.42XA, T41.42XD, T41.42XS, T41.5X2A, T41.5X2D, T41.5X2S,
T42.0X2A, T42.0X2D, T42.0X2S, T42.1X2A, T42.1X2D, T42.1X2S, T42.2X2A,
T42.2X2D, T42.2X2S, T42.3X2A, T42.3X2D, T42.3X2S, T42.4X2A, T42.4X2D,
T42.4X2S, T42.5X2A, T42.5X2D, T42.5X2S, T42.6X2A, T42.6X2D, T42.6X2S,
T42.72XA, T42.72XD, T42.72XS, T42.8X2A, T42.8X2D, T42.8X2S, T43.012A,
T43.012D, T43.012S, T43.022A, T43.022D, T43.022S, T43.1X2A, T43.1X2D,
T43.1X2S, T43.202A, T43.202D, T43.202S, T43.212A, T43.212D, T43.212S,
T43.222A, T43.222D, T43.222S, T43.292A, T43.292D, T43.292S, T43.3X2A,
T43.3X2D, T43.3X2S, T43.4X2A, T43.4X2D, T43.4X2S, T43.502A, T43.502D,
T43.502S, T43.592A, T43.592D, T43.592S, T43.602A, T43.602D, T43.602S,
T43.622A, T43.622D, T43.622S, T43.632A, T43.632D, T43.632S, T43.642A,
T43.642D, T43.642S, T43.652A, T43.652D, T43.652S, T43.692A, T43.692D,
T43.692S, T43.8X2A, T43.8X2D, T43.8X2S, T43.92XA, T43.92XD, T43.92XS,
T44.0X2A, T44.0X2D, T44.0X2S, T44.1X2A, T44.1X2D, T44.1X2S, T44.2X2A,
T44.2X2D, T44.2X2S, T44.3X2A, T44.3X2D, T44.3X2S, T44.4X2A, T44.4X2D,
T44.4X2S, T44.5X2A, T44.5X2D, T44.5X2S, T44.6X2A, T44.6X2D, T44.6X2S,
T44.7X2A, T44.7X2D, T44.7X2S, T44.8X2A, T44.8X2D, T44.8X2S, T44.902A,
T44.902D, T44.902S, T44.992A, T44.992D, T44.992S, T45.0X2A, T45.0X2D,
T45.0X2S, T45.1X2A, T45.1X2D, T45.1X2S, T45.2X2A, T45.2X2D, T45.2X2S,
T45.3X2A, T45.3X2D, T45.3X2S, T45.4X2A, T45.4X2D, T45.4X2S, T45.512A,
T45.512D, T45.512S, T45.522A, T45.522D, T45.522S, T45.602A, T45.602D,
T45.602S, T45.612A, T45.612D, T45.612S, T45.622A, T45.622D, T45.622S,
T45.692A, T45.692D, T45.692S, T45.7X2A, T45.7X2D, T45.7X2S, T45.8X2A,
T45.8X2D, T45.8X2S, T45.92XA, T45.92XD, T45.92XS, T46.0X2A, T46.0X2D,

T46.0X2S, T46.1X2A, T46.1X2D, T46.1X2S, T46.2X2A, T46.2X2D, T46.2X2S,
T46.3X2A, T46.3X2D, T46.3X2S, T46.4X2A, T46.4X2D, T46.4X2S, T46.5X2A,
T46.5X2D, T46.5X2S, T46.6X2A, T46.6X2D, T46.6X2S, T46.7X2A, T46.7X2D,
T46.7X2S, T46.8X2A, T46.8X2D, T46.8X2S, T46.902A, T46.902D, T46.902S,
T46.992A, T46.992D, T46.992S, T47.0X2A, T47.0X2D, T47.0X2S, T47.1X2A,
T47.1X2D, T47.1X2S, T47.2X2A, T47.2X2D, T47.2X2S, T47.3X2A, T47.3X2D,
T47.3X2S, T47.4X2A, T47.4X2D, T47.4X2S, T47.5X2A, T47.5X2D, T47.5X2S,
T47.6X2A, T47.6X2D, T47.6X2S, T47.7X2A, T47.7X2D, T47.7X2S, T47.8X2A,
T47.8X2D, T47.8X2S, T47.92XA, T47.92XD, T47.92XS, T48.0X2A, T48.0X2D,
T48.0X2S, T48.1X2A, T48.1X2D, T48.1X2S, T48.202A, T48.202D, T48.202S,
T48.292A, T48.292D, T48.292S, T48.3X2A, T48.3X2D, T48.3X2S, T48.4X2A,
T48.4X2D, T48.4X2S, T48.5X2A, T48.5X2D, T48.5X2S, T48.6X2A, T48.6X2D,
T48.6X2S, T48.902A, T48.902D, T48.902S, T48.992A, T48.992D, T48.992S,
T49.0X2A, T49.0X2D, T49.0X2S, T49.1X2A, T49.1X2D, T49.1X2S, T49.2X2A,
T49.2X2D, T49.2X2S, T49.3X2A, T49.3X2D, T49.3X2S, T49.4X2A, T49.4X2D,
T49.4X2S, T49.5X2A, T49.5X2D, T49.5X2S, T49.6X2A, T49.6X2D, T49.6X2S,
T49.7X2A, T49.7X2D, T49.7X2S, T49.8X2A, T49.8X2D, T49.8X2S, T49.92XA,
T49.92XD, T49.92XS, T50.0X2A, T50.0X2D, T50.0X2S, T50.1X2A, T50.1X2D,
T50.1X2S, T50.2X2A, T50.2X2D, T50.2X2S, T50.3X2A, T50.3X2D, T50.3X2S,
T50.4X2A, T50.4X2D, T50.4X2S, T50.5X2A, T50.5X2D, T50.5X2S, T50.6X2A,
T50.6X2D, T50.6X2S, T50.7X2A, T50.7X2D, T50.7X2S, T50.8X2A, T50.8X2D,
T50.8X2S, T50.902A, T50.902D, T50.902S, T50.912A, T50.912D, T50.912S,
T50.992A, T50.992D, T50.992S, T50.A12A, T50.A12D, T50.A12S, T50.A22A,
T50.A22D, T50.A22S, T50.A92A, T50.A92D, T50.A92S, T50.B12A, T50.B12D,
T50.B12S, T50.B92A, T50.B92D, T50.B92S, T50.Z12A, T50.Z12D, T50.Z12S,
T50.Z92A, T50.Z92D, T50.Z92S, T51.0X2A, T51.0X2D, T51.0X2S, T51.1X2A,
T51.1X2D, T51.1X2S, T51.2X2A, T51.2X2D, T51.2X2S, T51.3X2A, T51.3X2D,
T51.3X2S, T51.8X2A, T51.8X2D, T51.8X2S, T51.92XA, T51.92XD, T51.92XS,
T52.0X2A, T52.0X2D, T52.0X2S, T52.1X2A, T52.1X2D, T52.1X2S, T52.2X2A,
T52.2X2D, T52.2X2S, T52.3X2A, T52.3X2D, T52.3X2S, T52.4X2A, T52.4X2D,
T52.4X2S, T52.8X2A, T52.8X2D, T52.8X2S, T52.92XA, T52.92XD, T52.92XS,
T53.0X2A, T53.0X2D, T53.0X2S, T53.1X2A, T53.1X2D, T53.1X2S, T53.2X2A,
T53.2X2D, T53.2X2S, T53.3X2A, T53.3X2D, T53.3X2S, T53.4X2A, T53.4X2D,
T53.4X2S, T53.5X2A, T53.5X2D, T53.5X2S, T53.6X2A, T53.6X2D, T53.6X2S,
T53.7X2A, T53.7X2D, T53.7X2S, T53.92XA, T53.92XD, T53.92XS, T54.0X2A,
T54.0X2D, T54.0X2S, T54.1X2A, T54.1X2D, T54.1X2S, T54.2X2A, T54.2X2D,
T54.2X2S, T54.3X2A, T54.3X2D, T54.3X2S, T54.92XA, T54.92XD, T54.92XS,
T55.0X2A, T55.0X2D, T55.0X2S, T55.1X2A, T55.1X2D, T55.1X2S, T56.0X2A,
T56.0X2D, T56.0X2S, T56.1X2A, T56.1X2D, T56.1X2S, T56.2X2A, T56.2X2D,

T56.2X2S, T56.3X2A, T56.3X2D, T56.3X2S, T56.4X2A, T56.4X2D, T56.4X2S,
T56.5X2A, T56.5X2D, T56.5X2S, T56.6X2A, T56.6X2D, T56.6X2S, T56.7X2A,
T56.7X2D, T56.7X2S, T56.812A, T56.812D, T56.812S, T56.892A, T56.892D,
T56.892S, T56.92XA, T56.92XD, T56.92XS, T57.0X2A, T57.0X2D, T57.0X2S,
T57.1X2A, T57.1X2D, T57.1X2S, T57.2X2A, T57.2X2D, T57.2X2S, T57.3X2A,
T57.3X2D, T57.3X2S, T57.8X2A, T57.8X2D, T57.8X2S, T57.92XA, T57.92XD,
T57.92XS, T58.02XA, T58.02XD, T58.02XS, T58.12XA, T58.12XD, T58.12XS,
T58.2X2A, T58.2X2D, T58.2X2S, T58.8X2A, T58.8X2D, T58.8X2S, T58.92XA,
T58.92XD, T58.92XS, T59.0X2A, T59.0X2D, T59.0X2S, T59.1X2A, T59.1X2D,
T59.1X2S, T59.2X2A, T59.2X2D, T59.2X2S, T59.3X2A, T59.3X2D, T59.3X2S,
T59.4X2A, T59.4X2D, T59.4X2S, T59.5X2A, T59.5X2D, T59.5X2S, T59.6X2A,
T59.6X2D, T59.6X2S, T59.7X2A, T59.7X2D, T59.7X2S, T59.812A, T59.812D,
T59.812S, T59.892A, T59.892D, T59.892S, T59.92XA, T59.92XD, T59.92XS,
T60.0X2A, T60.0X2D, T60.0X2S, T60.1X2A, T60.1X2D, T60.1X2S, T60.2X2A,
T60.2X2D, T60.2X2S, T60.3X2A, T60.3X2D, T60.3X2S, T60.4X2A, T60.4X2D,
T60.4X2S, T60.8X2A, T60.8X2D, T60.8X2S, T60.92XA, T60.92XD, T60.92XS,
T61.02XA, T61.02XD, T61.02XS, T61.12XA, T61.12XD, T61.12XS, T61.772A,
T61.772D, T61.772S, T61.782A, T61.782D, T61.782S, T61.8X2A, T61.8X2D,
T61.8X2S, T61.92XA, T61.92XD, T61.92XS, T62.0X2A, T62.0X2D, T62.0X2S,
T62.1X2A, T62.1X2D, T62.1X2S, T62.2X2A, T62.2X2D, T62.2X2S, T62.8X2A,
T62.8X2D, T62.8X2S, T62.92XA, T62.92XD, T62.92XS, T63.002A, T63.002D,
T63.002S, T63.012A, T63.012D, T63.012S, T63.022A, T63.022D, T63.022S,
T63.032A, T63.032D, T63.032S, T63.042A, T63.042D, T63.042S, T63.062A,
T63.062D, T63.062S, T63.072A, T63.072D, T63.072S, T63.082A, T63.082D,
T63.082S, T63.092A, T63.092D, T63.092S, T63.112A, T63.112D, T63.112S,
T63.122A, T63.122D, T63.122S, T63.192A, T63.192D, T63.192S, T63.2X2A,
T63.2X2D, T63.2X2S, T63.302A, T63.302D, T63.302S, T63.312A, T63.312D,
T63.312S, T63.322A, T63.322D, T63.322S, T63.332A, T63.332D, T63.332S,
T63.392A, T63.392D, T63.392S, T63.412A, T63.412D, T63.412S, T63.422A,
T63.422D, T63.422S, T63.432A, T63.432D, T63.432S, T63.442A, T63.442D,
T63.442S, T63.452A, T63.452D, T63.452S, T63.462A, T63.462D, T63.462S,
T63.482A, T63.482D, T63.482S, T63.512A, T63.512D, T63.512S, T63.592A,
T63.592D, T63.592S, T63.612A, T63.612D, T63.612S, T63.622A, T63.622D,
T63.622S, T63.632A, T63.632D, T63.632S, T63.692A, T63.692D, T63.692S,
T63.712A, T63.712D, T63.712S, T63.792A, T63.792D, T63.792S, T63.812A,
T63.812D, T63.812S, T63.822A, T63.822D, T63.822S, T63.832A, T63.832D,
T63.832S, T63.892A, T63.892D, T63.892S, T63.92XA, T63.92XD, T63.92XS,
T64.02XA, T64.02XD, T64.02XS, T64.82XA, T64.82XD, T64.82XS, T65.0X2A,
T65.0X2D, T65.0X2S, T65.1X2A, T65.1X2D, T65.1X2S, T65.212A, T65.212D,

T65.212S, T65.222A, T65.222D, T65.222S, T65.292A, T65.292D, T65.292S,
T65.3X2A, T65.3X2D, T65.3X2S, T65.4X2A, T65.4X2D, T65.4X2S, T65.5X2A,
T65.5X2D, T65.5X2S, T65.6X2A, T65.6X2D, T65.6X2S, T65.812A, T65.812D,
T65.812S, T65.822A, T65.822D, T65.822S, T65.832A, T65.832D, T65.832S,
T65.892A, T65.892D, T65.892S, T65.92XA, T65.92XD, T65.92XS, T71.112A,
T71.112D, T71.112S, T71.122A, T71.122D, T71.122S, T71.132A, T71.132D,
T71.132S, T71.152A, T71.152D, T71.152S, T71.162A, T71.162D, T71.162S,
T71.192A, T71.192D, T71.192S, T71.222A, T71.222D, T71.222S, T71.232A,
T71.232D, T71.232S,
Intentional Self-Harm Diagnosis:
ICD10CM: T14.91XA, T14.91XD, T14.91XS, T36.0X2A, T36.0X2D, T36.0X2S,
T36.1X2A, T36.1X2D, T36.1X2S, T36.2X2A, T36.2X2D, T36.2X2S, T36.3X2A,
T36.3X2D, T36.3X2S, T36.4X2A, T36.4X2D, T36.4X2S, T36.5X2A, T36.5X2D,
T36.5X2S, T36.6X2A, T36.6X2D, T36.6X2S, T36.7X2A, T36.7X2D, T36.7X2S,
T36.8X2A, T36.8X2D, T36.8X2S, T36.92XA, T36.92XD, T36.92XS, T37.0X2A,
T37.0X2D, T37.0X2S, T37.1X2A, T37.1X2D, T37.1X2S, T37.2X2A, T37.2X2D,
T37.2X2S, T37.3X2A, T37.3X2D, T37.3X2S, T37.4X2A, T37.4X2D, T37.4X2S,
T37.5X2A, T37.5X2D, T37.5X2S, T37.8X2A, T37.8X2D, T37.8X2S, T37.92XA,
T37.92XD, T37.92XS, T38.0X2A, T38.0X2D, T38.0X2S, T38.1X2A, T38.1X2D,
T38.1X2S, T38.2X2A, T38.2X2D, T38.2X2S, T38.3X2A, T38.3X2D, T38.3X2S,
T38.4X2A, T38.4X2D, T38.4X2S, T38.5X2A, T38.5X2D, T38.5X2S, T38.6X2A,
T38.6X2D, T38.6X2S, T38.7X2A, T38.7X2D, T38.7X2S, T38.802A, T38.802D,
T38.802S, T38.812A, T38.812D, T38.812S, T38.892A, T38.892D, T38.892S,
T38.902A, T38.902D, T38.902S, T38.992A, T38.992D, T38.992S, T39.012A,
T39.012D, T39.012S, T39.092A, T39.092D, T39.092S, T39.1X2A, T39.1X2D,
T39.1X2S, T39.2X2A, T39.2X2D, T39.2X2S, T39.312A, T39.312D, T39.312S,
T39.392A, T39.392D, T39.392S, T39.4X2A, T39.4X2D, T39.4X2S, T39.8X2A,
T39.8X2D, T39.8X2S, T39.92XA, T39.92XD, T39.92XS, T40.0X2A, T40.0X2D,
T40.0X2S, T40.1X2A, T40.1X2D, T40.1X2S, T40.2X2A, T40.2X2D, T40.2X2S,
T40.3X2A, T40.3X2D, T40.3X2S, T40.412A, T40.412D, T40.412S, T40.422A,
T40.422D, T40.422S, T40.492A, T40.492D, T40.492S, T40.5X2A, T40.5X2D,
T40.5X2S, T40.602A, T40.602D, T40.602S, T40.692A, T40.692D, T40.692S,
T40.712A, T40.712D, T40.722A, T40.722D, T40.722S, T40.8X2A, T40.8X2D,
T40.8X2S, T40.902A, T40.902D, T40.902S, T40.992A, T40.992D, T40.992S,
T41.0X2A, T41.0X2D, T41.0X2S, T41.1X2A, T41.1X2D, T41.1X2S, T41.202A,
T41.202D, T41.202S, T41.292A, T41.292D, T41.292S, T41.3X2A, T41.3X2D,
T41.3X2S, T41.42XA, T41.42XD, T41.42XS, T41.5X2A, T41.5X2D, T41.5X2S,
T42.0X2A, T42.0X2D, T42.0X2S, T42.1X2A, T42.1X2D, T42.1X2S, T42.2X2A,

 -
T42.2X2D, T42.2X2S, T42.3X2A, T42.3X2D, T42.3X2S, T42.4X2A, T42.4X2D,
T42.4X2S, T42.5X2A, T42.5X2D, T42.5X2S, T42.6X2A, T42.6X2D, T42.6X2S,
T42.72XA, T42.72XD, T42.72XS, T42.8X2A, T42.8X2D, T42.8X2S, T43.012A,
T43.012D, T43.012S, T43.022A, T43.022D, T43.022S, T43.1X2A, T43.1X2D,
T43.1X2S, T43.202A, T43.202D, T43.202S, T43.212A, T43.212D, T43.212S,
T43.222A, T43.222D, T43.222S, T43.292A, T43.292D, T43.292S, T43.3X2A,
T43.3X2D, T43.3X2S, T43.4X2A, T43.4X2D, T43.4X2S, T43.502A, T43.502D,
T43.502S, T43.592A, T43.592D, T43.592S, T43.602A, T43.602D, T43.602S,
T43.622A, T43.622D, T43.622S, T43.632A, T43.632D, T43.632S, T43.642A,
T43.642D, T43.642S, T43.652A, T43.652D, T43.652S, T43.692A, T43.692D,
T43.692S, T43.8X2A, T43.8X2D, T43.8X2S, T43.92XA, T43.92XD, T43.92XS,
T44.0X2A, T44.0X2D, T44.0X2S, T44.1X2A, T44.1X2D, T44.1X2S, T44.2X2A,
T44.2X2D, T44.2X2S, T44.3X2A, T44.3X2D, T44.3X2S, T44.4X2A, T44.4X2D,
T44.4X2S, T44.5X2A, T44.5X2D, T44.5X2S, T44.6X2A, T44.6X2D, T44.6X2S,
T44.7X2A, T44.7X2D, T44.7X2S, T44.8X2A, T44.8X2D, T44.8X2S, T44.902A,
T44.902D, T44.902S, T44.992A, T44.992D, T44.992S, T45.0X2A, T45.0X2D,
T45.0X2S, T45.1X2A, T45.1X2D, T45.1X2S, T45.2X2A, T45.2X2D, T45.2X2S,
T45.3X2A, T45.3X2D, T45.3X2S, T45.4X2A, T45.4X2D, T45.4X2S, T45.512A,
T45.512D, T45.512S, T45.522A, T45.522D, T45.522S, T45.602A, T45.602D,
T45.602S, T45.612A, T45.612D, T45.612S, T45.622A, T45.622D, T45.622S,
T45.692A, T45.692D, T45.692S, T45.7X2A, T45.7X2D, T45.7X2S, T45.8X2A,
T45.8X2D, T45.8X2S, T45.92XA, T45.92XD, T45.92XS, T46.0X2A, T46.0X2D,
T46.0X2S, T46.1X2A, T46.1X2D, T46.1X2S, T46.2X2A, T46.2X2D, T46.2X2S,
T46.3X2A, T46.3X2D, T46.3X2S, T46.4X2A, T46.4X2D, T46.4X2S, T46.5X2A,
T46.5X2D, T46.5X2S, T46.6X2A, T46.6X2D, T46.6X2S, T46.7X2A, T46.7X2D,
T46.7X2S, T46.8X2A, T46.8X2D, T46.8X2S, T46.902A, T46.902D, T46.902S,
T46.992A, T46.992D, T46.992S, T47.0X2A, T47.0X2D, T47.0X2S, T47.1X2A,
T47.1X2D, T47.1X2S, T47.2X2A, T47.2X2D, T47.2X2S, T47.3X2A, T47.3X2D,
T47.3X2S, T47.4X2A, T47.4X2D, T47.4X2S, T47.5X2A, T47.5X2D, T47.5X2S,
T47.6X2A, T47.6X2D, T47.6X2S, T47.7X2A, T47.7X2D, T47.7X2S, T47.8X2A,
T47.8X2D, T47.8X2S, T47.92XA, T47.92XD, T47.92XS, T48.0X2A, T48.0X2D,
T48.0X2S, T48.1X2A, T48.1X2D, T48.1X2S, T48.202A, T48.202D, T48.202S,
T48.292A, T48.292D, T48.292S, T48.3X2A, T48.3X2D, T48.3X2S, T48.4X2A,
T48.4X2D, T48.4X2S, T48.5X2A, T48.5X2D, T48.5X2S, T48.6X2A, T48.6X2D,
T48.6X2S, T48.902A, T48.902D, T48.902S, T48.992A, T48.992D, T48.992S,
T49.0X2A, T49.0X2D, T49.0X2S, T49.1X2A, T49.1X2D, T49.1X2S, T49.2X2A,
T49.2X2D, T49.2X2S, T49.3X2A, T49.3X2D, T49.3X2S, T49.4X2A, T49.4X2D,
T49.4X2S, T49.5X2A, T49.5X2D, T49.5X2S, T49.6X2A, T49.6X2D, T49.6X2S,
T49.7X2A, T49.7X2D, T49.7X2S, T49.8X2A, T49.8X2D, T49.8X2S, T49.92XA,

	Τ4	9.92XD, T49.92XS, T50.0X2A, T50).0X2D, T50.0X2S, T50.1X2A, T50.1X2D,	
	T5	0.1X2S, T50.2X2A, T50.2X2D, T50).2X2S, T50.3X2A, T50.3X2D, T50.3X2S,	
	T5	0.4X2A, T50.4X2D, T50.4X2S, T50).5X2A, T50.5X2D, T50.5X2S, T50.6X2A,	
	T5	0.6X2D, T50.6X2S, T50.7X2A, T50).7X2D, T50.7X2S, T50.8X2A, T50.8X2D,	
	T5	0.8X2S, T50.902A, T50.902D, T50).902S, T50.912A, T50.912D, T50.912S,	
	T5	0.992A, T50.992D, T50.992S, T50).A12A, T50.A12D, T50.A12S, T50.A22A,	
	T5	0.A22D, T50.A22S, T50.A92A, T5	0.A92D, T50.A92S, T50.B12A, T50.B12D,	
	T5	0.B12S, T50.B92A, T50.B92D, T50	0.B92S, T50.Z12A, T50.Z12D, T50.Z12S,	
	T5	0.Z92A, T50.Z92D, T50.Z92S, T51	0X2A, T51.0X2D, T51.0X2S, T51.1X2A,	
	T5	1.1X2D, T51.1X2S, T51.2X2A, T51	L.2X2D, T51.2X2S, T51.3X2A, T51.3X2D,	
	Т5	1.3X2S, T51.8X2A, T51.8X2D, T51	L.8X2S, T51.92XA, T51.92XD, T51.92XS,	
	T5	2.0X2A, T52.0X2D, T52.0X2S, T52	2.1X2A, T52.1X2D, T52.1X2S, T52.2X2A,	
	T5	2.2X2D, T52.2X2S, T52.3X2A, T52	2.3X2D, T52.3X2S, T52.4X2A, T52.4X2D,	
	T5	2.4X2S, T52.8X2A, T52.8X2D, T52	2.8X2S, T52.92XA, T52.92XD, T52.92XS,	
	T5	3.0X2A, T53.0X2D, T53.0X2S, T53	3.1X2A, T53.1X2D, T53.1X2S, T53.2X2A,	
	T5	3.2X2D, T53.2X2S, T53.3X2A, T53	3.3X2D, T53.3X2S, T53.4X2A, T53.4X2D,	
	T5.	3.4X2S, T53.5X2A, T53.5X2D, T53	3.5X2S, T53.6X2A, T53.6X2D, T53.6X2S,	
	T5	3.7X2A, T53.7X2D, T53.7X2S, T53	3.92XA, T53.92XD, T53.92XS, T54.0X2A,	
	T5-	4.0X2D, T54.0X2S, T54.1X2A, T54	4.1X2D, T54.1X2S, T54.2X2A, T54.2X2D,	
	T5	4.2X2S, T54.3X2A, T54.3X2D, T54	4.3X2S, T54.92XA, T54.92XD, T54.92XS,	
	T5.	5.0X2A, T55.0X2D, T55.0X2S, T55	5.1X2A, T55.1X2D, T55.1X2S, T56.0X2A,	
	T5	6.0X2D, T56.0X2S, T56.1X2A, T56	5.1X2D, T56.1X2S, T56.2X2A, T56.2X2D,	
	T5	6.2X2S, T56.3X2A, T56.3X2D, T56	5.3X2S, T56.4X2A, T56.4X2D, T56.4X2S,	
	T5	6.5X2A, T56.5X2D, T56.5X2S, T56	5.6X2A, T56.6X2D, T56.6X2S, T56.7X2A,	
	T5	6.7X2D, T56.7X2S, T56.812A, T56	5.812D, T56.812S, T56.892A, T56.892D,	
	T5	6.892S, T56.92XA, T56.92XD, T56	5.92XS, T57.0X2A, T57.0X2D, T57.0X2S,	
	T5	7.1X2A, T57.1X2D, T57.1X2S, T57	7.2X2A, T57.2X2D, T57.2X2S, T57.3X2A,	
	T5	7.3X2D, T57.3X2S, T57.8X2A, T57	7.8X2D, T57.8X2S, T57.92XA, T57.92XD,	
	T5	7.92XS, T58.02XA, T58.02XD, T58	3.02XS, T58.12XA, T58.12XD, T58.12XS,	
	T5	8.2X2A, T58.2X2D, T58.2X2S, T58	3.8X2A, T58.8X2D, T58.8X2S, T58.92XA,	
	T5	8.92XD, T58.92XS, T59.0X2A, T59	9.0X2D, T59.0X2S, T59.1X2A, T59.1X2D,	
	T5	9.1X2S, T59.2X2A, T59.2X2D, T59	9.2X2S, T59.3X2A, T59.3X2D, T59.3X2S,	
	T5	9.4X2A, T59.4X2D, T59.4X2S, T59	9.5X2A, T59.5X2D, T59.5X2S, T59.6X2A,	
	T5	9.6X2D, T59.6X2S, T59.7X2A, T59	9.7X2D, T59.7X2S, T59.812A, T59.812D,	
	T5	9.812S, T59.892A, T59.892D, T59	9.892S, T59.92XA, T59.92XD, T59.92XS,	
).1X2A, T60.1X2D, T60.1X2S, T60.2X2A,	
	Т6	0.2X2D, T60.2X2S, T60.3X2A, T60).3X2D, T60.3X2S, T60.4X2A, T60.4X2D,	
			D.8X2S, T60.92XA, T60.92XD, T60.92XS,	
	Т6	1.02XA, T61.02XD, T61.02XS, T61	1.12XA, T61.12XD, T61.12XS, T61.772A,	

T61.772D, T61.772S, T61.782A, T61.782D, T61.782S, T61.8X2A, T61.8X2D,
T61.8X2S, T61.92XA, T61.92XD, T61.92XS, T62.0X2A, T62.0X2D, T62.0X2S,
T62.1X2A, T62.1X2D, T62.1X2S, T62.2X2A, T62.2X2D, T62.2X2S, T62.8X2A,
T62.8X2D, T62.8X2S, T62.92XA, T62.92XD, T62.92XS, T63.002A, T63.002D,
T63.002S, T63.012A, T63.012D, T63.012S, T63.022A, T63.022D, T63.022S,
T63.032A, T63.032D, T63.032S, T63.042A, T63.042D, T63.042S, T63.062A,
T63.062D, T63.062S, T63.072A, T63.072D, T63.072S, T63.082A, T63.082D,
T63.082S, T63.092A, T63.092D, T63.092S, T63.112A, T63.112D, T63.112S,
T63.122A, T63.122D, T63.122S, T63.192A, T63.192D, T63.192S, T63.2X2A,
T63.2X2D, T63.2X2S, T63.302A, T63.302D, T63.302S, T63.312A, T63.312D,
T63.312S, T63.322A, T63.322D, T63.322S, T63.332A, T63.332D, T63.332S,
T63.392A, T63.392D, T63.392S, T63.412A, T63.412D, T63.412S, T63.422A,
T63.422D, T63.422S, T63.432A, T63.432D, T63.432S, T63.442A, T63.442D,
T63.442S, T63.452A, T63.452D, T63.452S, T63.462A, T63.462D, T63.462S,
T63.482A, T63.482D, T63.482S, T63.512A, T63.512D, T63.512S, T63.592A,
T63.592D, T63.592S, T63.612A, T63.612D, T63.612S, T63.622A, T63.622D,
T63.622S, T63.632A, T63.632D, T63.632S, T63.692A, T63.692D, T63.692S,
T63.712A, T63.712D, T63.712S, T63.792A, T63.792D, T63.792S, T63.812A,
T63.812D, T63.812S, T63.822A, T63.822D, T63.822S, T63.832A, T63.832D,
T63.832S, T63.892A, T63.892D, T63.892S, T63.92XA, T63.92XD, T63.92XS,
T64.02XA, T64.02XD, T64.02XS, T64.82XA, T64.82XD, T64.82XS, T65.0X2A,
T65.0X2D, T65.0X2S, T65.1X2A, T65.1X2D, T65.1X2S, T65.212A, T65.212D,
T65.212S, T65.222A, T65.222D, T65.222S, T65.292A, T65.292D, T65.292S,
T65.3X2A, T65.3X2D, T65.3X2S, T65.4X2A, T65.4X2D, T65.4X2S, T65.5X2A,
T65.5X2D, T65.5X2S, T65.6X2A, T65.6X2D, T65.6X2S, T65.812A, T65.812D,
T65.812S, T65.822A, T65.822D, T65.822S, T65.832A, T65.832D, T65.832S,
T65.892A, T65.892D, T65.892S, T65.92XA, T65.92XD, T65.92XS, T71.112A,
T71.112D, T71.112S, T71.122A, T71.122D, T71.122S, T71.132A, T71.132D,
T71.132S, T71.152A, T71.152D, T71.152S, T71.162A, T71.162D, T71.162S,
T71.192A, T71.192D, T71.192S, T71.222A, T71.222D, T71.222S, T71.232A,
T71.232D, T71.232S
Note: LOINC and SNOMED codes can be captured through electronic data
submissions. Please contact your Account Executive for more information.

Measure	Measure Description	Measure Information/Documentation Required	Coding
Follow-Up After Emergency Department Visit for Substance Use (FUA)	 The percentage of emergency department (ED) visits for members 13 years of age and older with a principal diagnosis of substance use disorder (SUD), or any diagnosis of drug overdose, for which there was follow up. Two rates are reported: 1. The percentage of ED visits for which the member received follow-up within 30 days of the ED visit (31 total days). 2. The percentage of ED visits for which the member received follow up within 7 days of the ED visit (8 total days). 	 A follow-up visit or a pharmacotherapy dispensing event within 30 days after the ED visit (31 total days). Includes visits that occur on the date of the ED visit. A follow-up visit or pharmacotherapy dispensing event within 7 days after the ED visit (8 total days). Include visits that occur on the date of the ED visit. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. 	 Visit Setting Unspecified: (With Outpatient POS and with a Principal Diagnosis of AOD Abuse and Dependence, Substance Induced Disorders or Unintentional Drug Overdose, or with Mental Health Provider) (With Partial Hospitalization POS and with a Principal Diagnosis of AOD Abuse and Dependence, Substance Induced Disorders or Unintentional Drug Overdose, or with Mental Health Provider) (With Non-residential Substance Abuse Treatment Facility POS and with any Diagnosis of AOD Abuse and Dependence, Substance Induced Disorders or Unintentional Drug Overdose, or with Mental Health Provider) (With Community Mental Health Center POS, and with Any Diagnosis of AOD Abuse and Dependence, Substance Induced Disorders or Unintentional Drug Overdose, or with A Mental Health Provider) (With Community Mental Health Provider) (With Telehealth POS, and with Any Diagnosis of AOD Abuse and Dependence, Substance Induced Disorders or Unintentional Drug Overdose, or with a Mental Health Provider) (With Telehealth POS, and with Any Diagnosis of AOD Abuse and Dependence, Substance Induced Disorders or Unintentional Drug Overdose, or with Mental Health Provider) (With Telehealth POS, and with Any Diagnosis of AOD Abuse and Dependence, Substance Induced Disorders or Unintentional Drug Overdose, or with Mental Health Provider): CPT: 90791, 90792, 90832, 90833, 90875, 90876, 99221, 99222, 99233, 99234, 99232, 99233, 99234, 99252, 99253, 99254, 99255 BH Outpatient: (With Any Diagnosis of AOD Abuse and Dependence, Substance Induced Disorders or Unintentional Drug Overdose, or with a Mental Health Provider) CPT: 98960, 98861, 98962, 99078, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99243, 99244, 99245, 99341, 99342, 99344, 99345, 99347, 99348, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99344, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99483

		Partial Hospitalization or Intensive Outpatient Visit:
		(With Any Diagnosis of AOD Abuse and Dependence, Substance Induced
		Disorders or Unintentional Drug Overdose, or with a Mental Health Provider)
		HCPCS: H2001, H2012, S0201, S9480, S9484, S9485
		UBREV: 0907, 0912, 0913 covered at BH Rural Care Clinic
		Peer Support Service:
		(With Any Diagnosis of AOD Abuse and Dependence, Substance Induced
		Disorders or Unintentional Drug Overdose):
		HCPCS: H0038, H0040,S9445
		Teleshawa Matta
		Telephone Visits:
		(With Any Diagnosis of AOD Abuse and Dependence, Substance Induced
		Disorders or Unintentional Drug Overdose, or with Mental Health Provider):
		CPT: 99441, 99442, 99443
		Online Assessments:
		(With Any Diagnosis of AOD Abuse and Dependence, Substance Induced
		Disorders or Unintentional Drug Overdose, or with Mental Health Provider):
		CPT: 98980, 98981,99421, 99422, 99423, 99457, 99458
		HCPCS: G2010, G201, G2251
		Substance Abuse Counseling and Surveillance:
		ICD10CM: Z71.41, Z71.51
		Substance Use Disorder Services:
		HCPCS: G0396, G0397, H0001, H0005, H0015, H2036
		UBREV: 0906, 0944, 0945
		Behavioral Health Assessment:
		HCPCS: G0396, G0397, G2011, H0001
		Substance Use Services:
		HCPCS: H0006
		Pharmacotherapy-Dispensing Event:
		Alcohol Use Disorder Treatment Medications:
	l	

Aldehyde dehydrogenase inhibitor: Disulfiram (oral)
Antagonist: Naltrexone (oral and injectable)
Other: Acamprosate (oral and delayed-release tablet)
Opioid Use Disorder Treatment Medications:
Antagonist: Naltrexone (oral and injectable)
Partial agonist: Buprenorphine (sublingual tablet, injection, implant),
Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film)
AOD Medication Treatment:
HCPCS: J0570, J0571, J0572, J0573, J0574, J0575, J2315, Q9991, Q9992
Outpatient POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49,
50, 71, 72
Non-Residential Substance Abuse POS: 57, 58
Community Mental Health POS: 53
Telehealth POS: 02, 10
AOD Abuse and Dependence Diagnosis:
ICD10CM: F10.10, F10.120, F10.121, F10.129, F10.130, F10.131, F10.132,
F10.139, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182,
F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232,
F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281,
F10.282, F10.288, F10.29, F11.10, F11.120, F11.121, F11.122, F11.129, F11.13,
F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20,
F11.220, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259,
F11.281, F11.282, F11.288, F11.29, F12.10, F12.120, F12.121, F12.122, F12.129,
F12.13, F12.150, F12.151, F12.159, F12.180, F12.188, F12.19, F12.20, F12.220,
F12.221, F12.222, F12.229, F12.23, F12.250, F12.251, F12.259, F12.280,
F12.288, F12.29, F13.10, F13.120, F13.121, F13.129, F13.130, F13.131, F13.132,
F13.139, F13.14, F13.150, F13.151, F13.159, F13.180, F13.181, F13.182,
F13.188, F13.19, F13.20, F13.220, F13.221, F13.229, F13.230, F13.231, F13.232,
F13.239, F13.24, F13.250, F13.251, F13.259, F13.26, F13.27, F13.280, F13.281,
F13.282, F13.288, F13.29, F14.10, F14.120, F14.121, F14.122, F14.129, F14.13,
F14.14, F14.150, F14.151, F14.159, F14.180, F14.181, F14.182, F14.188, F14.19,
F14.20, F14.220, F14.221, F14.222, F14.229, F14.23, F14.24, F14.250, F14.251,
F14.259, F14.280, F14.281, F14.282, F14.288, F14.29, F15.10, F15.120, F15.121,
F15.122, F15.129, F15.13, F15.14, F15.150, F15.151, F15.159, F15.180, F15.181,

· · · · · · · · · · · · · · · · · · ·	
	F15.182, F15.188, F15.19, F15.20, F15.220, F15.221, F15.222, F15.229, F15.23,
	F15.24, F15.250, F15.251, F15.259, F15.280, F15.281, F15.282, F15.288, F15.29,
	F16.10, F16.120, F16.121, F16.122, F16.129, F16.14, F16.150, F16.151, F16.159,
	F16.180, F16.183, F16.188, F16.19, F16.20, F16.220, F16.221, F16.229, F16.24,
	F16.250, F16.251, F16.259, F16.280, F16.283, F16.288, F16.29, F18.10, F18.120,
	F18.121, F18.129, F18.14, F18.150, F18.151, F18.159, F18.17, F18.180, F18.188,
	F18.19, F18.20, F18.220, F18.221, F18.229, F18.24, F18.250, F18.251, F18.259,
	F18.27, F18.280, F18.288, F18.29, F19.10, F19.120, F19.121, F19.122, F19.129,
	F19.130, F19.131, F19.132, F19.139, F19.14, F19.150, F19.151, F19.159, F19.16,
	F19.17, F19.180, F19.181, F19.182, F19.188, F19.19, F19.20, F19.220, F19.221,
	F19.222, F19.229, F19.230, F19.231, F19.232, F19.239, F19.24, F19.250,
	F19.251, F19.259, F19.26, F19.27, F19.280, F19.281, F19.282, F19.288, F19.29
	Substance Induced Disorders:
	ICD10CM: F10.90, F10.920, F10.921, F10.929, F10.930, F10.931, F10.932,
	F10.939, F10.94, F10.950, F10.951, F10.959, F10.96, F10.97, F10.980, F10.981,
	F10.982, F10.988, F10.99, F11.90, F11.920, F11.921, F11.922, F11.929, F11.93,
	F11.94, F11.950, F11.951, F11.959, F11.981, F11.982, F11.988, F11.99, F12.90,
	F12.920, F12.921, F12.922, F12.929, F12.93, F12.950, F12.951, F12.959,
	F12.980, F12.988, F12.99, F13.90, F13.920, F13.921, F13.929, F13.930, F13.931,
	F13.932, F13.939, F13.94, F13.950, F13.951, F13.959, F13.96, F13.97, F13.980,
	F13.981, F13.982, F13.988, F13.99, F14.90, F14.920, F14.921, F14.922, F14.929,
	F14.93, F14.94, F14.950, F14.951, F14.959, F14.980, F14.981, F14.982, F14.988,
	F14.99, F15.90, F15.920, F15.921, F15.922, F15.929, F15.93, F15.94, F15.950,
	F15.951, F15.959, F15.980, F15.981, F15.982, F15.988, F15.99, F16.90, F16.920,
	F16.921, F16.929, F16.94, F16.950, F16.951, F16.959, F16.980, F16.983,
	F16.988, F16.99, F18.90, F18.920, F18.921, F18.929, F18.94, F18.950, F18.951,
	F18.959, F18.97, F18.980, F18.988, F18.99, F19.90, F19.920, F19.921, F19.922,
	F19.929, F19.930, F19.931, F19.932, F19.939, F19.94, F19.950, F19.951,
	F19.959, F19.96, F19.97, F19.980, F19.981, F19.982, F19.988, F19.99
	Unintentional Drug Overdose:
	ICD10CM: T40.0X1A, T40.0X1D, T40.0X1S, T40.0X4A, T40.0X4D, T40.0X4S,
	T40.1X1A, T40.1X1D, T40.1X1S, T40.1X4A, T40.1X4D, T40.1X4S, T40.2X1A,
	T40.2X1D, T40.2X1S, T40.2X4A, T40.2X4D, T40.2X4S, T40.3X1A, T40.3X1D,
	T40.3X1S, T40.3X4A, T40.3X4D, T40.3X4S, T40.411A, T40.411D, T40.411S,
	T40.414A, T40.414D, T40.414S, T40.421A, T40.421D, T40.421S, T40.424A,
	T40.424D, T40.424S, T40.491A, T40.491D, T40.491S, T40.494A, T40.494D,
	T40.494S, T40.5X1A, T40.5X1D, T40.5X1S, T40.5X4A, T40.5X4D, T40.5X4S,

			T40.601A, T40.601D, T40.601S, T40.604A, T40.604D, T40.604S, T40.691A, T40.691D, T40.691S, T40.694A, T40.694D, T40.694S, T40.711A, T40.711D,
			T40.711S, T40.714A, T40.714D, T40.721A, T40.721S, T40.724A, T40.724D,
			T40.724S, T40.8X1A, T40.8X1D, T40.8X1S, T40.8X4A, T40.8X4D, T40.8X4S,
			T40.901A, T40.901D, T40.901S, T40.904A, T40.904D, T40.904S, T40.904S, T40.991A,
			T40.991D, T40.991S, T40.994A, T40.994D, T40.994S, T41.0X1A, T41.0X1D,
			T41.0X1S, T41.0X4A, T41.0X4D, T41.0X4S, T41.1X1A, T41.1X1D, T41.1X1S,
			T41.1X4A, T41.1X4D, T41.1X4S, T41.201A, T41.201D, T41.201S, T41.204A,
			T41.204D, T41.204S, T41.291A, T41.291D, T41.291S, T41.294A, T41.294D,
			T41.294S, T41.3X1A, T41.3X1D, T41.3X1S, T41.3X4A, T41.3X4D, T41.3X4S,
			T41.41XA, T41.41XD, T41.41XS, T41.44XA, T41.44XD, T41.44XS, T41.5X1A,
			T41.5X1D, T41.5X1S, T41.5X4A, T41.5X4D, T41.5X4S, T42.3X1A, T42.3X1D,
			T42.3X1S, T42.3X4A, T42.3X4D, T42.3X4S, T42.4X1A, T42.4X1D, T42.4X1S,
			T42.4X4A, T42.4X4D, T42.4X4S, T43.601A, T43.601D, T43.601S, T43.604A,
			T43.604D, T43.604S, T43.621A, T43.621D, T43.621S, T43.624A, T43.624D,
			T43.624S, T43.631A, T43.631D, T43.631S, T43.634A, T43.634D, T43.634S,
			T43.641A, T43.641D, T43.641S, T43.644A, T43.644D, T43.644S, T43.651A,
			T43.651D, T53.651S, T43.654A, T43.654D, T43.654S, T43.691A, T43.691D,
			T43.691S, T43.694A, T43.694D, T43.694S, T51.0X1A, T51.0X1D, T51.0X1S,
			T51.0X4A, T51.0X4D, T51.0X4S
			Note: LOINC and SNOMED codes can be captured through electronic data
			submissions. Please contact your Account Executive for more information.
Measure	Measure Description	Measure	Coding
		Information/Documentation	
		Required	
Follow-Up After High-	The percentage of	The percentage of acute inpatient	Visit Setting Unspecified:
Intensity Care for	members 13 years of	hospitalizations, residential	(with Outpatient POS and with a Principal Diagnosis of AOD Abuse and
Substance Use Disorder	age or older who had an	treatment, or withdrawal	Dependence):
(FUI)	acute inpatient	management visits for a diagnosis of	(with BH Outpatient Visit and with a Principal Diagnosis of AOD Abuse and
	hospitalization, residential treatment or	substance use disorder among members 13 years of age and older	Dependence): (with Partial Hospitalization POS and with a Principal Diagnosis of AOD Abuse
	withdrawal	that result in a follow-up visit or	and Dependence):
	management visit for a	service for substance use disorder:	(with Non-Residential Substance Abuse Treatment Facility POS and with a
	diagnosis of substance		Principal Diagnosis of AOD Abuse and Dependence):
	use disorder (SUD) that	7-Day Follow-Up: A follow-up visit or	(with Community Mental Health Center POS and with a Principal Diagnosis of
		event with any practitioner for a	AOD Abuse and Dependence):
	1	event with any practitioner for a	Nouse and Dependence).

resulted in a follow-up	principal diagnosis of substance use	(with Telehealth POS and with a Principal Diagnosis of AOD Abuse and
visit or service for SUD.	disorder within the 7 days after an	Dependence):
	episode for substance use disorder.	CPT: 90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221, 99222, 99223, 99231,
	30-Day Follow-Up: A follow-up visit or	99232, 99233, 99238, 99239, 99252, 99253, 99254, 99255
	event with any practitioner for a	
	principal diagnosis of substance use	
	disorder within the 30 days after an	BH Outpatient:
	episode for substance use disorder.	(with Principal Diagnosis of AOD Abuse and Dependence):
		CPT: 98960, 98961, 98962, 99078, 99202, 99203, 99204, 99205, 99211, 99212,
	Note:	99213, 99214, 99215, 99242, 99243, 99244, 99245, 99341, 99342, 99344,
	• Methadone is not included in the	99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385,
	medication lists for the measure.	99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401,
	Follow-up does not include	99402, 99403, 99404, 99411, 99412, 99483
	withdrawal management.	HCPCS: G0463, H0036, H0040, H2000, H2017, H2019, H2020, T1015
		UBREV: 0510, 0513, 0515, 0516, 0517, 0519, 0900, 0904, 0911, 0914, 0915,
	Required Exclusions:	0916, 0919
	Members who meet any of the	
	following criteria are excluded from	Partial Hospitalization or Intensive Outpatient Visit:
	the measure:	(with a Principal Diagnosis of AOD Abuse and Dependence):
	In hospice or using hospice	HCPCS: H2001, H2012, S0201, S9480, S9484, S9485
	services any time in the MY.	
	• Deceased at any time in the MY.	Substance Use Disorder Services:
		(with a Principal Diagnosis of AOD Abuse and Dependence):
		HCPCS: G0396, G0397, H0001, H0005, H0015, H2036, T1002
		UBREV: 0906, 0944, 0945 OP PT ONLY
		Substance Abuse Counseling and Surveillance:
		ICD10CM: Z71.41, Z71.51
		Residential Behavioral Health Treatment:
		(with a Principal Diagnosis of AOD Abuse and Dependence):
		Telephone Visit (with a Principal Diagnosis of AOD Abuse and Dependence): CPT: 99441, 99442, 99443
		Online Assessments:
		(with a Principal Diagnosis of AOD Abuse and Dependence):
		CPT: 99421, 99422, 99423, 99457, 99458

	HCPCS: G2010, G2012, G2251
	Pharmacotherapy-Dispensing Event:
	Alcohol Use Disorder Treatment Medications:
	Aldehyde dehydrogenase inhibitor: Disulfiram (oral)
	Antagonist: Naltrexone (oral and injectable)
	Other: Acamprosate (oral and delayed-release tablet)
	Opioid Use Disorder Treatment Medications:
	Antagonist: Naltrexone (oral and injectable)
	Partial agonist: Buprenorphine (sublingual tablet, injection, implant),
	Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film)
	AOD Medication Treatment:
	HCPCS: J0570, J0571, J0572, J0573, J0574, J0575, J2315, Q9991, Q9992
	Outpatient POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49,
	50, 71, 72
	Non-Residential Substance Abuse POS: 57, 58
	Community Montel Health DOC: 52
	Community Mental Health POS: 53
	Telehealth POS: 02
	AOD Abuse and Dependence Diagnosis:
	ICD10CM: F10.10, F10.120, F10.121, F10.129, F10.130, F10.131, F10.132,
	F10.139, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182,
	F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232,
	F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281,
	F10.282, F10.288, F10.29, F11.10, F11.120, F11.121, F11.122, F11.129, F11.13,
	F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20,
	F11.220, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259,
	F11.281, F11.282, F11.288, F11.29, F12.10, F12.120, F12.121, F12.122, F12.129,
	F12.13, F12.150, F12.151, F12.159, F12.180, F12.188, F12.19, F12.20, F12.220,
	F12.221, F12.222, F12.229, F12.23, F12.250, F12.251, F12.259, F12.280,
	F12.288, F12.29, F13.10, F13.120, F13.121, F13.129, F13.130, F13.131, F13.132,
	F13.139, F13.14, F13.150, F13.151, F13.159, F13.180, F13.181, F13.182,

			F13.188, F13.19, F13.20, F13.220, F13.221, F13.229, F13.230, F13.231, F13.232, F13.239, F13.24, F13.250, F13.251, F13.259, F13.26, F13.27, F13.280, F13.281, F13.282, F13.288, F13.29, F14.10, F14.120, F14.121, F14.122, F14.129, F14.13, F14.14, F14.150, F14.151, F14.159, F14.180, F14.181, F14.182, F14.188, F14.19, F14.20, F14.220, F14.221, F14.222, F14.229, F14.23, F14.24, F14.250, F14.251, F14.259, F14.280, F14.281, F14.282, F14.288, F14.29, F15.10, F15.120, F15.121, F15.122, F15.129, F15.13, F15.14, F15.150, F15.151, F15.159, F15.180, F15.181, F15.182, F15.188, F15.19, F15.20, F15.220, F15.221, F15.222, F15.229, F15.23, F15.24, F15.250, F15.251, F15.259, F15.280, F15.281, F15.282, F15.288, F15.29, F16.10, F16.120, F16.121, F16.122, F16.129, F16.14, F16.150, F16.151, F16.159, F16.180, F16.183, F16.188, F16.19, F16.20, F16.220, F16.221, F16.229, F16.24, F16.250, F16.251, F16.259, F16.280, F16.283, F16.28, F16.29, F18.10, F18.120, F18.121, F18.129, F18.14, F18.150, F18.151, F18.159, F18.17, F18.180, F18.188, F18.19, F18.20, F18.220, F18.221, F18.229, F18.24, F18.250, F18.251, F18.259, F18.27, F18.280, F18.288, F18.29, F19.10, F19.120, F19.121, F19.122, F19.129, F19.130, F19.131, F19.132, F19.139, F19.14, F19.150, F19.151, F19.159, F19.16, F19.17, F19.180, F19.181, F19.182, F19.188, F19.19, F19.20, F19.220, F19.221, F19.222, F19.229, F19.230, F19.231, F19.232, F19.239, F19.24, F19.282, F19.288, F19.29, F19.251, F19.259, F19.26, F19.27, F19.280, F19.281, F19.282, F19.288, F19.29
			submissions. Please contact your Account Executive for more information.
Measure	Measure Description	Measure Information/Documentation Required	Coding
Pharmacotherapy for Opioid Use Disorder (POD)	The percentage of opioid use disorder (OUD) pharmacotherapy events that lasted at least 180 days among members 16 years of age and older with a diagnosis of OUD and a new pharmacotherapy event.	Intake period: 12-month period that begins on 7/1 of the year prior to the MY and ends on 6/30 of the MY. The Treatment Period (TP) is the date of an OUD dispensing event or OUD medication administration event during the IP. No more than an 8-day gap is allowed during the TP. Note:	Members are identified through administrative and pharmacy claims. Opioid Abuse and Dependence Diagnosis: ICD10CM: F11.10, F11.120, F11.121, F11.122, F11.129, F11.13, F11.14, F11.150, F11.151, F11.159, F11.181, F11.182, F11.188, F11.19, F11.20, F11.220, F11.221, F11.222, F11.229, F11.23, F11.24, F11.250, F11.251, F11.259, F11.281, F11.282, F11.288, F11.29 Opioid Use Disorder Treatment Medications: Antagonist: Naltrexone (oral) Antagonist: Naltrexone (injectable)

		• Methadone is not included in the	Partial agonist: Buprenorphine (sublingual tablet), Buprenorphine (injection),
		medication lists for the measure.	Buprenorphine (implant), Buprenorphine/naloxone (sublingual tablet, buccal
			film, sublingual film)
		Required Exclusions:	Agonist: Methadone (oral) is only acceptable when billed on a medical claim. A
		Members who meet any of the	pharmacy claim would be indicative of treatment for pain rather than OUD.
		following criteria are excluded from	
		the measure:	Buprenorphine Implant:
		• In hospice or using hospice	HCPCS: J0570
		services any time in the MY.	
		• Deceased at any time in the MY.	Buprenorphine Injection:
			HCPCS: Q9991, Q9992
			Buprenorphine Naloxone:
			HCPCS: J0572, J0573, J0574, J0575
			Buprenorphine Oral:
			HCPCS: J0571
			Methadone Oral:
			HCPCS: H0020,
			Naltrexone Injection:
			HCPCS: J2315
			HCrC3. 32313
			Note: LOINC and SNOMED codes can be captured through electronic data
			submissions. Please contact your Account Executive for more information.
			submissions. I lease contact your Account Executive for more information.
EFFECTIVENESS OF CARE: C	ARE COORDINATION	l 	
Measure	Measure Description	Measure	Coding
		Information/Documentation	
		Required	
Advance Care Planning	The percentage of	Advance Care Plan or discussion of	CPT : 99483, 99497
(ACP)	members 66 – 80 years	Advance Care Planning documented	CPT-CAT-II: 1123F, 1124F, 1157F, 1158F
	of age with advanced	in the medical record on or before	ICD10CM: Z66
1 st Year Measure	illness, frailty, or	12/31 of the MY.	
(MY2022)	receiving palliative care,		
[11112022]			1

	and adults 81 years of age or older, who had advance care planning during the MY.	 Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services in the MY. Deceased at any time in the MY. 	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Transition of Care (TRC)	 The percentage of discharges for members 18 years of age and older who had each of the following. Four rates are reported 1. Notification of Inpatient Admission. 2. Receipt of Discharge Information. 3. Patient Engagement After Inpatient Discharge. 4. Medication Reconciliation Post- Discharge. Each qualifying discharge in the MY is measured. 	 Notification of Inpatient Admission (NIA): Documentation must include evidence of receipt of notification of inpatient admission on the day of admission through the 2 days following admission (3 total days). Admission refers to the date of inpatient admission or date of admission for an observation stay that turns into an inpatient admission. Documentation must include evidence of receipt of notification of inpatient admission that includes evidence of the date when the documentation was received. Any of the following examples meet criteria: Communication between inpatient providers or staff and the member's PCP or ongoing care provider (e.g., phone call, email, fax). Communication about admission between emergency department and the member's PCP or ongoing care provider (e.g., phone call, email, fax). 	Patient Engagement Indicator: Outpatient and Telehealth: CPT: 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99241, 99242, 99243, 99344, 99345, 99347, 99348, 99348, 99335, 99384, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99421, 99422, 99423, 99441, 99442, 99443, 99457, 99458, 99483 HCPCS: G0402, G0463, G2010, G2012, G2251, T1015 Online Assessments: CPT: 99421, 99422, 99423, 99457, 99458 HCPCS: G2010, G2012, G2251 Medication Reconciliation Post-Discharge Indicator: Medication Reconciliation Intervention: CPT: 99483 Medication of Inpatient Admission and Receipt of Discharge Information has no administrative reporting option. They are based on medical record review only.

 Communication about admission to the member's PCP or ongoing care provider through a health information exchange; an automated admission, discharge, and transfer (ADT) alert system; or a shared electronic medical record system. Communication about admission to the member's PCP or ongoing care provider from the member's health plan. Indication that the member's PCP or ongoing care provider admitted the member to the hospital. Indication that a specialist admitted the member to the hospital. Indication that a specialist admitted the member to the hospital. Indication that a specialist admitted the member's PCP or ongoing care provider placed orders for tests and treatments any time during the member's inpatient stay. Documentation that the PCP or ongoing care provider performed a preadmission exam or received communication about a planned inpatient admission. The time frame that the planned inpatient admitsen must be
a preadmission exam or received communication about a planned inpatient admission. The time

notification of a planned	
admission prior to the admit date	
also meets criteria. The planned	
admission documentation or	
preadmission exam must clearly	
pertain to the admission.	
Receipt of Discharge Information	
(RDI):	
Documentation must include	
evidence of receipt of discharge	
information on the day of discharge	
through the 2 days following	
discharge.	
Discharge information may be	
included in, but not limited to, a	
discharge summary or summary of	
care record or be located in	
structured fields in an Electronic	
Health Record (EHR). At a minimum,	
the discharge information must	
include all of the following:	
The practitioner responsible for	
the member's care during the	
inpatient stay.	
Procedures or treatment provided	
provided.	
Diagnoses at discharge.	
Current medication list.	
Testing results, or documentation	
of pending tests or no tests	
pending.	
Instructions for patient care post-	
discharge.	
Detient Engagement After Innetient	
Patient Engagement After Inpatient	
Discharge (PE):	
Documentation must include	
evidence of patient engagement	

I	
	within 30 days following discharge.
	Any of the following meets criteria:
	An outpatient visit, including
	office visits and home visits.
	A telephone visit.
	A synchronous telehealth visit
	where real-time interaction
	occurred between the member
	and provider via telephone or
	video conferencing. Do not
	include patient engagement that
	occurs on the date of discharge.
	An e-visit or virtual check-in.
	Medication Reconciliation Post-
	Discharge (Med Rec):
	Documentation in the outpatient
	medical record must include evidence
	of medication reconciliation and the
	date it was performed by a
	prescribing practitioner (including
	physician assistant), clinical
	pharmacist, or registered nurse, as
	documented on the date of discharge
	through 30 days after discharge (31
	total days). Any of the following meet
	criteria:
	Documentation of the current
	medications with a notation that
	the provider reconciled the
	current and discharge
	medications.
	Documentation of the current
	medications with a notation that
	references the discharge
	medications (e.g., no changes in
	medications since discharge,
	same medications at discharge,

discontinue all discharge
medications).
Documentation of the member's
current medications with a
notation that the discharge
medications were reviewed.
Documentation of a current
medication list, a discharge
medication list, and notation that
both lists were reviewed on the
same date of service.
Documentation of the current
medications with evidence that
the member was seen for post-
discharge hospital follow-up with
evidence of medication
reconciliation or review.
Documentation in the discharge
summary that the discharge
medications were reconciled with
the most recent medication list in
the outpatient medical record.
There must be evidence that the
discharge summary was filed in
the outpatient chart on the date
of discharge through 30 days
after discharge (31 total days).
Notation that no medications
were prescribed or ordered upon
discharge.
Only documentation in the
outpatient chart meets the intent
of the rate, but an outpatient
visit is not required, and the
member does not have to be
present.
The following notations or examples
of documentation do not count as

· · · · · · · · · · · · · · · · · · ·		
	numerator compliant for Notification	
	of Inpatient Admission and	
	Notification of Inpatient Discharge:	
	Documentation that the member	
	or the member's family notified	
	the member's PCP or ongoing	
	care provider of the admission or	
	discharge.	
	Required Exclusions:	
	Members who meet any of the	
	following criteria are excluded from	
	the measure:	
	In hospice or using hospice	
	services any time in the MY.	
	• Deceased at any time in the MY.	
	Remained in an acute or	
	nonacute facility from discharge	
	through 12/1 of the MY.	
	Common Chart Deficiencies:	
	Inpatient records cannot be used	
	for TRC.	
	NIA: Documentation that a	
	provider sent the member to the	
	ED does not meet criteria.	
	NIA: Documentation that the	
	member or the member's family	
	member notified the PCP or	
	ongoing care provider of the	
	admission does not meet criteria.	
	NIA: Documentation of	
	notification that does not include	
	a time frame or date when the	
	documentation was received	
	documentation was received	
	does not meet criteria.	

		 PCP does not meet criteria — documentation of receipt is required. RDI: Discharge Summary not included in outpatient record or missing one or more of the 6 required elements. RDI: Documentation on Discharge Summary that communication was sent to the PCP does not meet criteria — documentation of receipt is required. PE: Patient engagement that occurs on the date of discharge, or more than 30 days after discharge, does not meet criteria. Med Rec: Completed by incorrect provider type. Med Rec: Documentation of current medications reviewed without reference to the hospitalization. Med Rec: Medication list found in both the discharge summary and outpatient record but no evidence the two were reconciled. 	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Follow-Up After Emergency Department Visit for People With	The percentage of emergency department (ED) visits for members	The MP is 1/1 through 12/24. ED visits that result in an inpatient	COPD Diagnosis: ICD10CM: J41.0, J41.1, J41.8, J42, J43.0, J43.1, J43.2, J43.8, J43.9, J44.0, J44.1, J44.9, J47.0, J47.1, J47.9
Multiple High-Risk Chronic Conditions (FMC)	18 years and older who have multiple high-risk chronic conditions who	stay or that are followed by admission to acute or nonacute inpatient care within 7 days are excluded.	Asthma Diagnosis:

had a fallow we are dea		
had a follow-up service	Chronic conditions include:	ICD10CM: J45.21, J45.22, J45.31, J45.32, J45.41, J45.42, J45.51, J45.52, J45.901,
within 7 days of the ED		J45.902, J45.990, J45.991, J45.998
visit.	COPD and asthma.Alzheimer's Disease and related	
Fach world in FD in the		Demontin
Each qualifying ED in the	disorders (dementia,	Dementia:
Measurement Period	frontotemporal dementia).	ICD10CM: F01.50, F01.51, F01.511, F01.518, F01.52, F01.53, F01.54, F01.A0,
(MP) is measured.	Chronic kidney disease.	F01.A11, F01.A18, F01.A2, F01.A3, F01.A4, F01.B0, F01.B11, F01.B2, F01.B3,
	Major depression.	F01.B4, F01.C0, F01.C11, F01.C18, F01.C2, F01.C3, F01.C4, F02.80, F02.81,
	Dysthymic disorder.	F02.811, F02.818, F02.82, F02.83, F02.84, F02.A0, F02.A11, F02.A18, F02.A2,
	Heart failure and chronic heart	F02.A3, F02.A4, F02.B0, F02.B11, F02.B18, F02.B2, F02.B3, F02.B4, F02.C11,
	failure.	F02.C18, F02.C2, F02.C3, F02.C4, F03.90, F03.91, F03.911, F03.918, F03.92,
	Acute myocardial infarction.	F03.93, F03.94, F03.A0, F03.A11, F03.A18, F03.A2, F03.A3, F03.A4, F03.B0,
	• Atrial fibrillation.	F03.B0, F03.B11, F03.B18, F03.B2, F03.B3, F03.B4, F03.C0, F03.C11, F03.C18,
	Stroke and transient ischemic	F03.C2, F03.C3, F03.C4, F04, F10.27, F10.97, F13.27, F13.97, F18.17, F18.27,
	attack.	F18.97, F19.17, F19.27, F19.97, G30.0, G30.1, G30.8, G30.9, G31.83
	Required Exclusions:	Frontotemporal Dementia:
	Members who meet any of the	ICD10CM: G31.01, G31.09
	following criteria are excluded from	
	the measure:	Chronic Kidney Disease:
	 In hospice or using hospice 	ICD10CM: A18.11, A52.75, B52.0, C64.1, C64.2, C64.9, C68.9, D30.00, D30.01,
	services any time in the MY.	D30.02, D41.00, D41.01, D41.02, D41.10, D41.11, D41.12, D41.20, D41.21,
	• Deceased at any time in the MY.	D41.22, D59.30, D59.31, D59.32, D59.39, E08.21, E08.22, E08.29, E08.65,
		E09.21, E09.22, E09.29, E10.21, E10.22, E10.29, E10.65, E11.21, E11.22, E11.29,
		E11.65, E13.21, E13.22, E13.29, E74.8, E74.810, E74.818, E74.819, E74.89, I12.0,
		113.11, 113.2, 170.1, 172.2, K76.7, M10.30, M10.311, M10.312, M10.319,
		M10.321, M10.322, M10.329, M10.331, M10.332, M10.339, M10.341,
		M10.322, M10.322, M10.329, M10.331, M10.332, M10.339, M10.341, M10.342, M10.349, M10.351, M10.352, M10.359, M10.361, M10.362,
		M10.369, M10.371, M10.372, M10.379, M10.38, M10.39, M32.14, M32.15,
		M35.04, N00.0, N00.1, N00.2, N00.3, N00.4, N00.5, N00.6, N00.7, N00.8, N00.9,
		N00.A, N01.0, N01.1, N01.2, N01.3, N01.4, N01.5, N01.6, N01.7, N01.8, N01.9,
		N01.A, N02.0, N02.1, N02.2, N02.3, N02.4, N02.5, N02.6, N02.7, N02.8, N02.9,
		N02.A, N03.0, N03.1, N03.2, N03.3, N03.4, N03.5, N03.6, N03.7, N03.8, N03.9,
		N03.A, N04.0, N04.1, N04.2, N04.3, N04.4, N04.5, N04.6, N04.7, N04.8, N04.9,
		N04.A, N05.0, N05.1, N05.2, N05.3, N05.4, N05.5, N05.6, N05.7, N05.8, N05.9,
		N05.A, N06.0, N06.1, N06.2, N06.3, N06.4, N06.5, N06.6, N06.7, N06.8, N06.9,
		N05.A, N06.0, N06.1, N06.2, N06.3, N06.4, N06.5, N06.6, N06.7, N06.8, N06.9, N06.A, N07.0, N07.1, N07.2, N07.3, N07.4, N07.5, N07.6, N07.7, N07.8, N07.9,

N14.2, N14.3, N14.4, N15.0, N15.8, N15.9, N16, N17.0, N17.1, N17.2, N17.8,
N17.9, N18.1, N18.2, N18.3, N18.30, N18.31, N18.32, N18.4, N18.5, N18.6,
N18.9, N19, N25.0, N25.1, N25.81, N25.89, N25.9, N26.1, N26.9, Q61.02,
Q61.11, Q61.19, Q61.2, Q61.3, Q61.4, Q61.5, Q61.8, Q62.0, Q62.10, Q62.11,
Q62.12, Q62.2, Q62.31, Q62.32, Q62.39, R94.4
Major Depression:
ICD10CM: F32.0, F32.1, F32.2, F32.3, F32.4, F32.9, F33.0, F33.1, F33.2, F33.3,
F33.41, F33.9
Dysthymic Disorder:
ICD10CM: F34.1
Chronic Heart Failure:
ICD10CM: 142.0, 142.1, 142.2, 142.3, 142.4, 142.5, 142.6, 142.7, 142.8, 142.9, 143,
150.1, 150.20, 150.21, 150.22, 150.23, 150.30, 150.31, 150.32, 150.33, 150.40, 150.41,
150.42, 150.43, 150.810, 150.811, 150.812, 150.813, 150.814, 150.82, 150.83, 150.84,
150.89, 150.9
150.05, 150.5
Heart Failure Diagnosis:
ICD10CM: 109.81, 111.0, 113.0, 113.2, 150.1, 150.20, 150.21, 150.22, 150.23, 150.30,
150.31, 150.32, 150.33, 150.40, 150.41, 150.42, 150.43, 150.810, 150.811, 150.812,
150.813, 150.814, 150.82, 150.83, 150.84, 150.89, 150.9
MI:
ICD10CM: I21.01, I21.02, I21.09, I21.11, I21.19, I21.21, I21.29, I21.3, I21.4,
121.9, 121.A1, 121.A9, 122.0, 122.1, 122.2, 122.8, 122.9, 123.0, 123.1, 123.2, 123.3,
123.4, 123.5, 123.6, 123.7, 123.8
Atrial Fibrillation:
ICD10CM: 148.0, 148.21, 148.91
Stroke:
ICD10CM: G45.0, G45.1, G45.2, G45.8, G45.9, G46.0, G46.1, G46.2, G97.31,
G97.32, I60.00, I60.01, I60.02, I61.0, I61.1, I61.2, I61.3, I61.4, I61.5, I61.6, I61.8,
161.9, 163.00, 163.011, 163.012, 163.019, 163.02, 163.031, 163.032, 163.039, 163.09,
163.10, 163.111, 163.112, 163.113, 163.119, 163.12, 163.131, 163.132, 163.133,

163.139, 163.19, 163.20, 163.211, 163.212, 163.213, 163.219, 163.22, 163.231,
163.232, 163.233, 163.239, 163.29, 163.30, 163.311, 163.312, 163.313, 163.319,
163.321, 163.322, 163.323, 163.329, 163.331, 163.332, 163.333, 163.339, 163.341,
163.342, 163.343, 163.349, 163.39, 163.40, 163.411, 163.412, 163.413, 163.419,
163.421, 163.422, 163.423, 163.429, 163.431, 163.432, 163.433, 163.439, 163.441,
163.442, 163.443, 163.449, 163.49, 163.50, 163.511, 163.512, 163.513, 163.519,
163.521, 163.522, 163.523, 163.529, 163.531, 163.532, 163.533, 163.539, 163.541,
163.542, 163.543, 163.549, 163.59, 163.6, 163.81, 163.89, 163.9, 166.01, 166.02,
166.03, 166.09, 166.11, 166.12, 166.13, 166.19, 166.21, 166.22, 166.23, 166.29, 166.3,
166.8, 166.9, 167.841, 167.848, 167.89, 197.810, 197.811, 197.820, 197.821
100.0, 100.0, 107.041, 107.040, 107.00, 107.010, 107.011, 107.020, 107.021
Follow-Up Service:
Outpatient Visit:
CPT: 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215,
99241, 99242, 99243, 99244, 99245, 99341, 99342, 99343, 99344, 99345,
99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386,
99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402,
99403, 99404, 99483
HCPCS: G0402, G0463, T1015
UBREV: 0510, 0511, 0512, 0513, 0514, 0515, 0516, 0517, 0519
CDREV : 0510, 0511, 0512, 0513, 0514, 0515, 0510, 0517, 0515
Outpatient and Telehealth:
CPT: 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99241,
99242, 99243, 99244, 99245, 99341, 99342, 99343, 99344, 99345, 99347,
99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387,
99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403,
99404, 99421, 99422, 99423, 99441, 99442, 99443, 99457, 99458, 99483
HCPCS: G0402, G0463, G2010, G2012, G2251, T1015
Case Management Encounter:
CPT: 99366
Complex Care Management Services:
CPT : 99487, 99489, 99490
Visit Catting Unspecified:
Visit Setting Unspecified:

(with Outpatient POS, Partial Hospitalization POS, Community Mental Health Center POS, or Telehealth POS): CPT: 90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221, 99222, 99223, 99231, 99232, 99233, 99238, 99239, 99252, 99253, 99254, 99255
BH Outpatient: CPT: 98960, 98961, 98962, 99078, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99243, 99244, 99245, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99483 HCPCS: G0463, H0036, H2000, H2017, H2019, H2020, T1015 UBREV: 0510, 0513, 0515, 0516, 0517, 0519, 0520, 0521, 0522, 0523, 0526, 0527, 0528, 0529, 0900, 0902, 0903, 0904, 0911, 0914, 0915, 0916, 0917, 0919, 0982, 0983
Partial Hospitalization or Intensive Outpatient: HCPCS: H2001, H2012, S0201, S9480, S9484, S9485
Electroconvulsive Therapy: (with Ambulatory Surgical Center POS, Community Mental Health Center POS, Outpatient POS, or Partial Hospitalization POS): CPT: 90870 ICD10PCS: GZB0ZZZ, GZB1ZZZ, GZB3ZZZ, GZB4ZZZ
Substance Abuse Counseling and Surveillance: ICD10CM: Z71.41, Z71.51
Substance Use Disorder Services: HCPCS: : G0396, G0397, H0001, H0005, H0015, H2036, UBREV: 0906, 0944, 0945
Online Assessments: CPT: 99421, 99422, 99423, 99457, 99458 HCPCS: G2010, G2012, G2251

Outpatient POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72 Telehealth POS: 02

Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

EFFECTIVENESS OF CARE: C	VERUSE/APPROPRIATENES	S	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Avoidance of Antibiotic	The percentage of	The Intake Period (IP) is the 12-month	Acute Bronchitis Diagnosis:
Treatment for Acute	episodes for members	window that begins 7/1 of the year	ICD10CM: J20.3, J20.4, J20.5, J20.6, J20.7, J20.8, J20.9, J21.0, J21.1, J21.8, J21.9
Bronchitis (AAB)	ages 3 months and older	prior to the MY and ends 6/30 of the	
	with a diagnosis of acute	MY.	AAB Antibiotic Medications:
This is also a measure	bronchitis/bronchiolitis		Aminoglycosides: Amikacin, Gentamicin, Streptomycin, Tobramycin
(AAB-E) collected through	that did not result in an	The Episode Date (EP) is the date of	Aminopenicillins: Amoxicillin, Ampicillin
claims and Electronic	antibiotic dispensing	service for any outpatient, telephone,	Beta-lactamase inhibitors: Amoxicillin-clavulanate, Ampicillin-sulbactam,
Clinical Data Systems.	event.	observation, or ED visit, e-visit or	Piperacillin-tazobactam
Please discuss options for		virtual check-in during the IP, with a	First-generation cephalosporins: Cefadroxil, Cefazolin, Cephalexin
a direct data feed with	Higher rate indicates	diagnosis of acute	Fourth-generation cephalosporins: Cefepime
your Account Executive.	appropriate treatment	bronchitis/bronchiolitis.	Lincomycin derivatives: Clindamycin, Lincomycin
Direct data feeds can	of adults with Acute		Macrolides: Azithromycin, Clarithromycin, Erythromycin
improve provider quality	Bronchitis (i.e., the	Dispensed prescription for an	Miscellaneous antibiotics: Aztreonam, Chloramphenicol, Dalfopristin-
performance and reduce	proportion for whom	antibiotic medication (AAB Antibiotic	quinupristin, Daptomycin, Linezolid, Metronidazole, Vancomycin
the burden of medical	antibiotics were not	Medications List) on or three days	Natural penicillins: Penicillin G benzathine-procaine, Penicillin G potassium,
record requests.	prescribed).	after the EP.	Penicillin G procaine, Penicillin G sodium, Penicillin V potassium, Penicillin G benzathine
		Required Exclusions:	Penicillinase-resistant penicillins: Dicloxacillin, Nafcillin, Oxacillin
		Members who meet any of the	Quinolones: Ciprofloxacin, Gemifloxacin, Levofloxacin, Moxifloxacin, Ofloxacin
		following criteria are excluded from	Rifamycin derivatives: Rifampin
		the measure:	Second-generation cephalosporin: Cefaclor, Cefotetan, Cefoxitin, Cefprozil,
		 In hospice or using hospice 	Cefuroxime
		services any time in the MY.	Sulfonamides: Sulfadiazine, Sulfamethoxazole-trimethoprim
		• Deceased at any time in the MY.	Tetracyclines: Doxycycline, Minocycline, Tetracycline
			Third-generation cephalosporins: Cefdinir, Cefixime, Cefotaxime, Cefpodoxime,
		Common Chart Deficiencies:	Ceftazidime, Ceftriaxone

		 Additional/competing diagnosis requiring antibiotics not documented in visit or coded on claim. 	Urinary anti-infectives: Fosfomycin, Nitrofurantoin, Nitrofurantoin macrocrystals-monohydrate, Trimethoprim Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
Measure	Measure Description	Measure Information/Documentation Required	Coding
Non-Recommended PSA- Based Screening in Older Men (PSA) This is also a measure (PSA-E) collected through claims and Electronic Clinical Data Systems. Please discuss options for a direct data feed with your Account Executive. Direct data feeds can improve provider quality performance and reduce the burden of medical record requests.	The percentage of male members 70 years and older who were screened unnecessarily for prostate cancer using prostate-specific antigen (PSA)–based screening. <i>A lower rate indicates</i> <i>better performance</i> .	 Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. Prostate cancer diagnosis any time during the member's history through December 31 of the MY. Dysplasia of the prostate during the MY or the year prior. A PSA test during the year prior to the MY where lab data indicate an elevated result (>4.0 nanograms/ milliliter) or an abnormal result. Dispensed prescription for a 5- alpha reductase inhibitor during the MY. 	PSA Lab Test: CPT: 84152, 84153, 84154 Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
Measure	Measure Description	Measure Information/Documentation Required	Coding
Appropriate Treatment for Upper Respiratory Infection (URI)	The percentage of episodes for members 3 months of age and older with a diagnosis of upper respiratory	The Intake Period (IP) is the 12-month window that begins July 1 of the year prior to the MY and ends on June 30 of the MY.	URI Diagnosis: ICD10CM: J00, J06.0, J06.9 Antibiotic Medications: Aminoglycosides: Amikacin, Gentamicin, Streptomycin, Tobramycin

This is also a measure (URI-E) collected through claims and Electronic Clinical Data Systems. Please discuss options for a direct data feed with your Account Executive. Direct data feeds can improve provider quality performance and reduce the burden of medical record requests.	infection (URI) that did not result in an antibiotic dispensing event. This is an episode-based event so a member may be included multiple times. Higher rate indicates appropriate treatment (i.e., the proportion for whom antibiotics were NOT prescribed).	The Episode Date (EP) is the Date of Service (DOS) for any outpatient, telephone, observation or ED visit, e- visit, or virtual check-in during the IP with a diagnosis or URI. If a member has more than one EP in a 31-day period, only the first EP will be used. Members with a comorbid condition during the 12 months prior to the EP will be excluded. These include: HIV, HIV Type 2. Malignant neoplasm. Emphysema. COPD. Disorders of the immune system. Other comorbid conditions. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. Additional/Competing diagnosis requiring antibiotics not	 Aminopenicillins: Amoxicillin, Ampicillin Beta-lactamase inhibitors: Amoxicillin-clavulanate, Ampicillin-sulbactam, Piperacillin-tazobactam First-generation cephalosporins: Cefadroxil, Cefazolin, Cephalexin Fourth-generation cephalosporins: Cefepime Lincomycin derivatives: Clindamycin, Lincomycin Macrolides: Azithromycin, Clarithromycin, Erythromycin Miscellaneous antibiotics: Aztreonam, Chloramphenicol, Dalfopristin- quinupristin, Daptomycin, Linezolid, Metronidazole, Vancomycin Natural penicillins: Penicillin G benzathine, Penicillin G benzathine-procaine, Penicillin G potassium, Penicillin G procaine, Penicillin G sodium, Penicillin V potassium Penicillinase-resistant penicillins: Dicloxacillin, Nafcillin, Oxacillin Quinolones: Ciprofloxacin, Gemifloxacin, Levofloxacin, Moxifloxacin, Ofloxacin, Rifamycin derivatives: Rifampin Second-generation cephalosporins: Cefaclor, Cefotetan, Cefoxitin, Cefprozil, Cefuroxime Sulfonamides: Sulfadiazine, Sulfamethoxazole-trimethoprim Tetracyclines: Doxycycline, Minocycline, Tetracycline Third-generation cephalosporins: Cefdinir, Cefixime, Cefotaxime, Cefpodoxime Ceftazidime, Ceftriaxone Urinary anti-infectives: Fosfomycin, Nitrofurantoin, Nitrofurantoin macrocrystals-monohydrate, Trimethoprim Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

Measure	Measure Description	Measure Information/Documentation	Coding
		Required	
Potentially Harmful Drug-	The percentage of	Required Exclusions:	HEDIS rates are based on Diagnosis and Medications/Pharmacy Claims.
Disease Interactions in	Medicare members 65	Members who meet any of the	
Older Adults (DDE)	years of age and older	following criteria are excluded from	Potentially Harmful Drugs — History of Falls Medications:
	who have evidence of	the measure:	Antiepileptics: Carbamazepine, Clobazam, Divalproex sodium, Ethosuximide,
	an underlying disease,	 In hospice or using hospice 	Ethotoin, Felbamate, Fosphenytoin, Gabapentin, Lacosamide, Lamotrigine,
	condition, or health	services any time in the MY.	Levetiracetam, Methsuximide, Oxcarbazepine, Phenobarbital, Phenytoin,
	concern and who were	• Deceased at any time in the MY.	Pregabalin, Primidone, Rufinamide, Tiagabine HCL, Topiramate, Valproic acid,
	dispensed an	 Receiving palliative care any time 	Vigabatrin, Zonisamide
	ambulatory prescription	in the MY.	SNRIs: Desvenlafaxine, Duloxetine, Levomilnacipran, Venlafaxine,
	for a potentially harmful medication concurrent	 History of falls and dementia rates only: A diagnosis of 	SSRIs: Citalopram, Escitalopram, Fluoxetine, Fluvoxamine, Paroxetine, Sertraline
	with or after the	psychosis, schizophrenia,	Potentially Harmful Drugs—History of Falls and Dementia Medications:
	diagnosis.	schizoaffective disorder, or	Antipsychotics: Aripiprazole, Asenapine, Brexpiprazole, Cariprazine,
		bipolar disorder on or between	Chlorpromazine, Clozapine, Fluphenazine, Haloperidol, Iloperidone, Loxapine,
	Three rates are	1/1 of the year prior to the MY	Lurasidone, Molindone, Olanzapine, Paliperidone, Perphenazine, Pimozide,
	reported:	and 12/1 of the MY.	Quetiapine, Risperidone, Thioridazine, Thiothixene, Trifluoperazine, Ziprasidone
	1. A history of falls	 History of falls rate only: A 	Benzodiazepines: Alprazolam, Chlordiazepoxide, Clonazepam, Clorazepate,
	and a prescription	diagnosis of major depressive	Diazepam, Estazolam, Flurazepam, Lorazepam, Midazolam, Oxazepam,
	for anticonvulsants,	disorder or seizure disorder on or	Quazepam, Temazepam, Triazolam
	SSRIs,	between 1/1 of the year prior to	Nonbenzodiazepine hypnotics: Eszopiclone, Zaleplon, Zolpidem
	antipsychotics,	the MY and 12/1 of the MY.	Tricyclic antidepressants: Amitriptyline, Amoxapine, Clomipramine,
	benzodiazepines,		Desipramine, Doxepin (>6 mg), Imipramine, Nortriptyline, Protriptyline,
	non-benzodiazepine		Trimipramine
	hypnotics, or		
	tricyclic		Dementia Medications:
	antidepressants.		Cholinesterase inhibitors: Donepezil, Galantamine, Rivastigmine
	2. Dementia and		Miscellaneous central nervous system agents: Memantine
	prescription for		Dementia combinations: Donepezil-Memantine
	antipsychotics,		
	benzodiazepines,		Potentially Harmful Drugs—Dementia Medications:
	non-benzodiazepine		Anticholinergic agents, antiemetics: Prochlorperazine, Promethazine
	hypnotics, tricyclic		Anticholinergic agents, antihistamines: Brompheniramine, Carbinoxamine,
	antidepressants, H2		Chlorpheniramine, Clemastine, Cyproheptadine, Dexbrompheniramine,
	receptor		Dexchlorpheniramine, Dimenhydrinate, Diphenhydramine, Doxylamine,
	antagonists, or		Pyrilamine, Triprolidine, Hydroxyzine, Meclizine

anticholine	ergic	Anticholinergic agents, antispasmodics: Atropine, Belladonna alkaloids,
agents.		Clidinium-chlordiazepoxide, Dicyclomine, Homatropine, Hyoscyamine,
3. Chronic kid	Iney	Methscopolamine, Propantheline, Scopolamine
disease and	d	Anticholinergic agents, antimuscarinics (oral): Darifenacin, Fesoterodine,
prescriptio	n for	Flavoxate, Oxybutynin, Solifenacin, Tolterodine, Trospium
Cox-2 selec	tive	Anticholinergic agents, anti-Parkinson agents: Benztropine, Trihexyphenidyl
NSAIDs or r	non-	Anticholinergic agents, skeletal muscle relaxants: Cyclobenzaprine,
aspirin NSA	AIDs.	Orphenadrine
		Anticholinergic agents, SSRIs: Paroxetine
Members with	more	Anticholinergic agents, antiarrhythmic: Disopyramide
than one diseas	se or	
condition may a	appear in	Cox-2 Selective NSAIDs and Nonaspirin NSAIDs:
the measure m		Cox-2 Selective NSAIDS: Celecoxib
times.		Nonaspirin NSAIDs: Diclofenac, Etodolac, Fenoprofen, Flurbiprofen, Ibuprofen,
		Indomethacin, Ketoprofen, Ketorolac, Meclofenamate, Mefenamic acid,
		Meloxicam, Nabumetone, Naproxen, Naproxen sodium, Oxaprozin, Piroxicam,
A lower rate in	dicates	Sulindac, Tolmetin
better perform	ance.	
		Dementia:
		ICD10CM: F01.50, F01.51, F02.80, F02.81, F03.90, F03.91, F04, F10.27, F10.97,
		F13.27, F13.97, F18.17, F18.27, F18.97, F19.17, F19.27, F19.97, G30.0, G30.1,
		G30.8, G30.9, G31.83
		Diagnosis of ESRD:
		ICD10CM: N18.5, N18.6, Z99.2
		Dialysis Procedure:
		CPT: 90935, 90937, 90945, 90947, 90997, 90999, 99512
		HCPCS: G0257
		ICD10PCS: 3E1M39Z, 5A1D00Z, 5A1D60Z, 5A1D70Z, 5A1D80Z, 5A1D90Z
		CKD Stage 4 Diagnosis:
		ICD10CM: N18.4
		Total Nephrectomy:
		CPT: 50220, 50225, 50230, 50234, 50236, 50240, 50340, 50370, 50543, 50545,
		50546, 50548
		50546, 50548

			ICD10: OTB00ZZ, OTB03ZZ, OTB04ZZ, OTB07ZZ, OTB08ZZ, OTB10ZZ, OTB13ZZ, OTB14ZZ, OTB17ZZ, OTB18ZZ, OTT00ZZ, OTT04ZZ, OTT10ZZ, OTT14ZZ, OTT20ZZ, OTT24ZZ Kidney Transplant: CPT: 50360, 50365, 50380 ICD10PCS: OTY00Z0, OTY00Z1, OTY00Z2, OTY10Z0, OTY10Z1, OTY10Z2 Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
Measure	Measure Description	Measure Information/Documentation Required	Coding
Risk of Continued Opioid Use (COU)	 The percentage of members 18 years of age and older who have a new episode of opioid use that puts them at risk for continued opioid use. Two rates are reported: 1. The percentage of members whose new episode of opioid use lasts at least 15 days in a 30-day period. 2. The percentage of members whose new episode of opioid use lasts at least 31 days in a 62-day period. 	 The MY is 1/1/-12/31. The Index Prescription Start Date (ISPD) is the earliest prescription dispensing date during the IP. 15-day: Prescriptions covering more than 15 calendar days during the 30-day period beginning on the ISPD through 29 days after the ISPD. 62-day: Prescriptions covering more than 31 calendar days during the 62-day period beginning on the ISPD through 61 days after the ISPD. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. 	 Opioid Medications: Benzhydrocodone, Buprenorphine (transdermal patch and buccal film), Butorphanol, Codeine, Dihydrocodeine, Fentanyl, Hydrocodone, Hydromorphone, Levorphanol, Meperidine, Methadone, Morphine, Opium, Oxycodone, Oxymorphone, Pentazocine, Tapentadol, Tramadol The Opioid Medications List excludes: Injectables. Opioid-containing cough and cold products. Single-agent and combination buprenorphine products used to treat opioid use disorder for medication-assisted treatment (buprenorphine sublingual tablets, buprenorphine subcutaneous implant, and all buprenorphine/naloxone combination products). Ionsys® (fentanyl transdermal patch). This is for inpatient use only and is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS). Methadone when prescribed for the treatment of opioid use disorder. Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

	A lower rate indicates better performance.	 Receiving palliative care during 12 months prior to the IPSD through 61 days after the IPSD. Cancer (Malignant Neoplasm) during 12 months prior to the IPSD through 61 days after the IPSD. Sickle Cell Anemia or HB S Disease during 12 months prior to the IPSD through 61 days after the IPSD. 	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Use of High-Risk Medication in Older Adults (DAE)	 The percentage of Medicare members 67 years of age and older who had at least two dispensing events for high-risk medications. Two rates are reported: At least 2 dispensing events for high-risk medications to avoid from the same drug class. At least 2 dispensing events for high-risk medications to avoid from the same drug class, except for appropriate diagnoses. 	 Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. Receiving palliative care in the MY. Common Chart Deficiencies: No documentation of review of medications at every visit. 	HEDIS rates are based on Diagnosis + Medications/Pharmacy Claims High-Risk Medications: Anticholinergics, first-generation antihistamines: Brompheniramine, Carbinoxamine, Chlorpheniramine, Clemastine. Cyproheptadine, Dexbrompheniramine, Dexchlorpheniramine, Diphenhydramine (oral), Dimenhydrinate, Doxylamine, Hydroxyzine, Meclizine, Promethazine, Pyrilamine, Triprolidine Anticholinergics, anti-Parkinson agents: Benztropine (oral), Trihexyphenidyl Antispasmodics: Atropine (exclude ophthalmic), Belladonna alkaloids, Chlordiazepoxide-clidinium, Dicyclomine, Hyoscyamine, Methscopolamine, Propantheline, Scopolamine Antithrombotic: Dipyridamole (oral excluding extended release) Cardiovascular, alpha agonists, central: Guanfacine, Methyldopa Cardiovascular, other: Disopyramide, Nifedipine, excluding extended release Central nervous system, antidepressants: Amitriptyline, Amoxapine, Clomipramine, Desipramine, Imipramine, Nortriptyline, Paroxetine, Protriptyline, Trimipramine Central nervous system, barbiturates: Amobarbital, Butabarbital, Butalbital, Pentobarbital, Phenobarbital, Secobarbital Central nervous system, vasodilators: Ergoloid mesylates, Isoxsuprine Central nervous system, vasodilators: Ergoloid mesylates, Isoxsuprine Central nervous system, other: Meprobamate Endocrine system, estrogens with or without progestins; include only oral and topical patch products: Conjugated estrogen, Esterified estrogen, Estradiol, Estropipate

3. Total rate (the sum	Endocrine system, sulfonylureas, long-duration: Chlorpropamide, Glimepiride,
of the two	Glyburide
numerators divided	Endocrine system, other: Desiccated thyroid, Megestrol
by the	Nonbenzodiazepine hypnotics: Eszopiclone, Zaleplon, Zolpidem
denominator,	Pain medications, skeletal muscle relaxants: Carisoprodol, Chlorzoxazone,
deduplicating for	Cyclobenzaprine, Metaxalone, Methocarbamol, Orphenadrine
members in both	Pain medications, other: Indomethacin, Ketorolac, includes parenteral,
numerators).	Meperidine
	High-Risk Medications with Days-Supply Criteria (<90 days):
A lower rate indicates	Anti-Infectives, other: Nitrofurantoin, Nitrofurantoin macrocrystals
better performance.	monohydrate
	Link Diel Madientiene with Average Deily Dees Criteries
	High-Risk Medications with Average Daily Dose Criteria:
	Alpha agonists, central: Reserpine >0.1 mg/day
	Cardiovascular, other: Digoxin >0.125 mg/day
	Tertiary TCAs (as single agent or as part of combination products): Doxepin >6
	mg/day
	High-Risk Medications Based on Prescription and Diagnosis Data:
	Antipsychotics, first (conventional) and second (atypical) generation:
	Aripiprazole, Aripiprazole lauroxil, Asenapine, Brexpiprazole, Cariprazine,
	Chlorpromazine, Clozapine, Fluphenazine, Haloperidol, Iloperidone, Loxapine,
	Lurasidone, Molindone, Olanzapine, Paliperidone, Perphenazine, Pimavanserin,
	Pimozide, Quetiapine, Risperidone, Thioridazine, Thiothixene, Trifluoperazine,
	Ziprasidone
	Benzodiazepines, long, short, and intermediate acting: Alprazolam,
	Chlordiazepoxide, Clonazepam, Clorazepate, Diazepam, Estazolam, Flurazepam,
	Lorazepam, Midazolam, Oxazepam, Quazepam, Temazepam, Triazolam
	Note: LOINC and SNOMED codes can be captured through electronic data
	submissions. Please contact your Account Executive for more information.

Measure	Measure Description	Measure Information/Documentation Required	Coding
Deprescribing of Benzodiazepines in Older Adults (DBO) 1 st Year Measure (MY2023)	The percentage of members 67 years of age and older who were dispensed benzodiazepines and achieved a 20% decrease or greater in dose (diazepam milligram equivalent [DME] dose) during the MY.	 The Measurement Year (MY) is 1/1 – 12/31. The Index Treatment Episode (ITE) is the first 30 days of a benzodiazepine prescription occurring during January 1 and September 1 of the MY. The ITE start date is the date of the earliest benzodiazepine prescription dispense date between January 1 and September 1 of the MY that is followed by at least 29 consecutive days with no gaps. The Treatment Period begins the day after the ITE and ends on the last covered day in the MY. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. A diagnosis between January 1 of the MY and the ITE of: Seizure disorder. Rapid eye movement. Benzodiazepine 	HEDIS rates are based on medications/pharmacy claims. Oral Benzodiazepine Medications: Alprazolam: 0.25 MG, 0.5 mg, 1 mg, 1 MGPML, 2 mg, 3 mg Chlordiazepoxide: 5 mg, 10 mg, 25 mg Clonazepam: 0.125 mg, 0.25 mg, 0.5 mg, 1 mg, 2 mg Clorazepate: 3.75 mg, 7.5 mg, 15 mg Diazepam: 1 MGPML, 2 mg, 5 MGPML, 10 mg Estazolam: 1 mg, 2 mg Flurazepam: 15 mg, 30 mg Lorazepam: 1 mg, 2 MGPML Midazolam: 2 MGPML Oxazepam: 10 mg, 30 mg Quazepam: 15 mg Temazepam: 7.5 mg, 15 mg, 22.5 mg, 30 mg Triazolam: 0.125 mg, 0.25 mg
		withdrawal. o Ethanol withdrawal.	

Measure	Measure Description	Measure Information/Documentation Required	Coding
Use of Imaging Studies for Low Back Pain (LBP)	The percentage of members 18 – 75 years of age with a primary diagnosis of low back pain who did not have an imaging study (plain X-ray, MRI, CT scan) within 28 days of the diagnosis.	 An imaging study with a diagnosis of uncomplicated low back pain on the IESD or in the 28 days following the IESD. Do not include outpatient, ED, or observation visits that result in an inpatient stay. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. G6 years of age and older with frailty and advanced illness during the MY. Any of the following anytime in the member's history through 28 days after the IESD: Cancer. HIV. Major organ transplant. Osteoporosis therapy. Lumbar surgery. Spondylopathy. Any of the following during 12 months (1 year) prior to the IESD through 28 days after the IESD: IV drug abuse. Neurologic impairment. Spinal infection. 	Imaging Study: CPT: 72020, 72052, 72100, 72110, 72114, 72120, 72131, 72132, 72133, 72141, 72142, 72146, 72147, 72148, 72149, 72156, 72158, 72200, 72202, 72220 Uncomplicated Low Back Pain: ICD10CM: M47.26, M47.27, M47.28, M47.816, M47.817, M47.818, M47.896, M47.897, M47.898, M48.061, M48.07, M48.08, M51.16, M51.17, M51.26, M51.27, M51.36, M51.37, M51.86, M51.87, M53.2X6, M53.2X7, M53.2X8, M53.3, M53.86, M53.87, M53.88, M54.16, M54.17, M54.18, M54.30, M54.31, M54.32, M54.40, M54.41, M54.42, M54.5, M54.50, M54.51, M54.59, M54.89, M54.9, M99.03, M99.04, M99.23, M99.33, M99.43, M99.53, M99.63, M99.73, M99.83, M99.84, S33.100A, S33.100D, S33.110A, S33.110D, S33.110S, S33.120A, S33.120D, S33.120S, S33.130A, S33.130D, S33.130S, S33.140A, S33.140D, S33.140S, S33.5XXA, S33.6XXA, S33.8XXA, S33.9XXA, S39.002A, S39.002D, S39.002S, S39.012A, S39.012D, S39.012S, S39.092A, S39.092D, S39.092S, S39.82XA, S39.82XD, S39.82XS, S39.92XA, S39.92XD, S39.92XS Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

		 Any of the following during the 3 months (90 days) prior to the IESD through 28 days after the IESD: Trauma. Fragility fracture. 90 consecutive days of corticosteroid treatment any time during the 366-day period that begins 365 days prior to the IESD and ends on the IESD. 	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Use of Opioids at High Dosage (HDO)	The percentage of members 18 years and older who received prescription opioids at a high dosage (average morphine milligram equivalent dose [MME] ≥90) for ≥15 days during the MY. <i>A lower rate indicates</i> <i>better performance.</i>	 Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. Receiving palliative care any time in the MY. Members with cancer (malignant neoplasm) in the MY. Members with sickle cell anemia, or HB S Disease, in the MY. 	Opioid Medications: Benzhydrocodone: Acetaminophen Benzhydrocodone (4.08 mg, 6.12 mg, 8.16 mg) Butorphanol: Butorphanol (10 MGPML) Codeine: Codeine Sulfate (15 mg, 30 mg, 60 mg), Acetaminophen Codeine (2.4 MGPML, 15 mg, 30 mg, 60 mg), Acetaminophen Butalbital Caffeine Codeine (30 mg), Aspirin Carisoprodol Codeine (16 mg) Dihydrocodeine: Acetaminophen Caffeine Dihydrocodeine (16 mg), Aspirin Caffeine Dihydrocodeine (16 mg) Fentanyl buccal or sublingual tablet, transmucosal lozenge (mcg): Fentanyl (100 mcg, 200 mcg, 300 mcg, 400 mcg, 600 mcg, 800 mcg, 1200 mcg, 1600 mcg) Fentanyl oral spray (mcg): Fentanyl (100 MCGPS, 200 MCGPS, 400 MCGPS, 600 MCGPS, 800 MCGPS) Fentanyl nasal spray (mcg): Fentanyl (100 MCGPS, 300 MCGPS, 400 MCGPS) Fentanyl transdermal film/patch (mcg/hr): Fentanyl (12 MCGPH, 25 MCGPH, 37.5 MCGPH, 50 MCGPH, 62.5 MCGPH, 75 MCGPH, 87.5 MCGPH, 100 MCGPH) Hydrocodone: Hydrocodone (10 mg, 15 mg, 20 mg, 30 mg, 40 mg, 50 mg, 60 mg, 80 mg, 100 mg, 120 mg), Acetaminophen Hydrocodone (.5 MGPML, .67 MGPML, 2.5 mg, 5 mg, 7.5 MGPML, 10 mg), Hydrocodone Ibuprofen (2.5 mg, 5 mg, 7.5 mg, 10 mg)

Required	 This is for inpatient use only and is available on program under a Risk Evaluation and Mitigation Methadone for the treatment of opioid use disorder Note: LOINC and SNOMED codes can be captured throw submissions. Please contact your Account Executive for 	Measure	Measure Description	Information/Documentation	 16 mg, 32 mg) Levorphanol: Levorphanol (2 mg, 3 mg) Meperidine: Meperidine (10 MGPML, 50 mg, 75mg, 100 mg, 150 mg), Methadone: Methadone (1 MGPML, 2 MGPML, 5 mg, 10 mg, 10 MGPML, 40 mg) Morphine: Morphine (2 MGPML, 4 MGPML, 5 mg, 10 mg, 15 mg, 20 MGPML, 20 mg, 30 mg, 40 mg, 45 mg, 50 mg, 60 mg, 75 mg, 80 mg, 90 mg, 100 mg, 120 mg, 200 mg), Morphine Naltrexone (20 mg, 30 mg, 50 mg, 60 mg, 80 mg, 100 mg) Opium: Belladonna Opium (30 mg, 60 mg) Oxycodone: Oxycodone (1 MGPML, 5 mg, 7.5 mg, 9 mg, 10 mg, 13.5 mg, 15 mg, 18 mg, 20 mg, 20 MGPML, 27 mg, 30 mg, 36 mg, 40 mg, 60 mg) Oxycodone: Oxycodone (1 MGPML, 2 MGPML, 2.5 mg, 5 mg, 7.5 mg, 10 mg), Aspirin Oxycodone (5 mg, 7.5 mg, 10 mg, 15 mg, 20 mg, 30 mg, 40 mg) Pentazocine: Naloxone Pentazocine (50 mg) Tapentadol: Tapentadol (50 mg, 75 mg, 100 mg, 150 mg, 200 mg, 250 mg) Tramadol: Tramadol (50 mg, 100 mg, 150 mg, 200 mg, 300 mg), Acetaminophen Tramadol (37.5 mg) The HDO Opioid Medications List excludes: Injectables. Opioid cough and cold products. Ionsys[®] (fentanyl transdermal patch). This is for inpatient use only and is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS). Methadone for the treatment of opioid use disorder. Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
			Weasure Description		
 This is for inpatient use only and is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS). Methadone for the treatment of opioid use disorder. Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information. asure Measure Description Measure Coding 					Injectables.Opioid cough and cold products.
Measure Measure Description Measure Measure Description Measure Description Injectables. Injectables. Opioid cough and cold products. Ionsys® (fentanyl transdermal patch). This is for inpatient use only and is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS). Methadone for the treatment of opioid use disorder. Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information. Measure	Injectables.Opioid cough and cold products.				Acetaminophen Tramadol (37.5 mg)
Acetaminophen Tramadol (37.5 mg) The HDO Opioid Medications List excludes: Injectables. Opioid cough and cold products. Ionsys® (fentanyl transdermal patch). This is for inpatient use only and is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS). Methadone for the treatment of opioid use disorder. Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information. Measure Measure Description Measure Coding Coding Coding	Acetaminophen Tramadol (37.5 mg) The HDO Opioid Medications List excludes: Injectables. Opioid cough and cold products. 				Tapentadol: Tapentadol (50 mg, 75 mg, 100 mg, 150 mg, 200 mg, 250 mg)
Tramadol: Tramadol (50 mg, 100 mg, 150 mg, 200 mg, 300 mg), Acetaminophen Tramadol (37.5 mg) The HDO Opioid Medications List excludes: Injectables. Injectables. Opioid cough and cold products. Ionsys® (fentanyl transdermal patch). This if for inpatient use only and is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS). Methadone for the treatment of opioid use disorder. Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information. Measure Measure Coding Measure Coding Measure Coding	Tramadol: Tramadol (50 mg, 100 mg, 150 mg, 200 mg, Acetaminophen Tramadol (37.5 mg) The HDO Opioid Medications List excludes: Injectables. Opioid cough and cold products.				
Pentazocine: Naloxone Pentazocine (50 mg) Tapentadol: Tapentadol: Tapentadol (50 mg, 75 mg, 100 mg, 150 mg, 200 mg, 250 mg) Tramadol: Tramadol: Tramadol (50 mg, 100 mg, 150 mg, 200 mg, 300 mg), Acetaminophen Tramadol (37.5 mg) The HDO Opioid Medications List excludes: Injectables. Opioid cough and cold products. Ionsys® (fentanyl transdermal patch). This for inpatient use only and is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS). Methadone for the treatment of opioid use disorder. Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information. Measure Measure Description	Pentazocine: Naloxone Pentazocine (50 mg) Tapentadol: Tapentadol (50 mg, 75 mg, 100 mg, 150 mg, 200 mg, Acetaminophen Tramadol (37.5 mg) The HDO Opioid Medications List excludes: • Injectables. • Opioid cough and cold products.				Oxymorphone: Oxymorphone (5 mg, 7.5 mg, 10 mg, 15 mg, 20 mg, 30 mg, 40
Oxymorphone: Oxymorphone (5 mg, 7.5 mg, 10 mg, 15 mg, 20 mg, 30 mg, 40 mg) Pentazocine: Naloxone Pentazocine (50 mg) Tapentadol: Tapentadol (50 mg, 100 mg, 150 mg, 200 mg, 250 mg) Tramadol: Tramadol (50 mg, 100 mg, 150 mg, 200 mg, 250 mg) Tramadol: Tramadol (37.5 mg) The HDO Opioid Medications List excludes: Injectables: Opioid cough and cold products. Ionsys* (fentanyl transdermal patch). This is for inpatient use only and is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS). Methadone for the treatment of opioid use disorder. Measure Measure Description	Oxymorphone: Oxymorphone (5 mg, 7.5 mg, 10 mg, 15 mg) Pentazocine: Naloxone Pentazocine (50 mg) Tapentadol: Tapentadol (50 mg, 75 mg, 100 mg, 150 m Tramadol: Tramadol (50 mg, 100 mg, 150 mg, 200 mg, Acetaminophen Tramadol (37.5 mg) The HDO Opioid Medications List excludes: Injectables. Opioid cough and cold products.				Acetaminophen Oxycodone (1 MGPML, 2 MGPML, 2.5 mg, 5 mg, 7.5 mg, 10
Acetaminophen Oxycodone (1 MGPML, 2 MGPML, 2.5 mg, 5 mg, 7.5 mg, 10 mg), Aspirin Oxycodone (4.84 mg), Ibuprofen Oxycodone (5 mg) Oxymorphone: Oxymorphone (5 mg, 7.5 mg, 10 mg, 15 mg, 20 mg, 30 mg, 40 mg) Pentazocine: Naloxone Pentazocine (50 mg) Tapentadol: Tapentadol (50 mg, 75 mg, 100 mg, 150 mg, 200 mg, 250 mg) Tramadol: Tramadol (50 mg, 100 mg, 150 mg, 200 mg, 250 mg) Tramadol: Tramadol: Tramadol (37.5 mg) The HDO Opioid Medications List excludes: Injectables. Opioid cough and cold products. Ionsys* (fentanyl transdermal patch). This is for inpatient use only and is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS). Methadone for the treatment of opioid use disorder. Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information. Measure Measure Description	Acetaminophen Oxycodone (1 MGPML, 2 MGPML, 2.5 mg), Aspirin Oxycodone (4.84 mg), Ibuprofen Oxycodor Oxymorphone: Oxymorphone (5 mg, 7.5 mg, 10 mg, 15 mg) Pentazocine: Naloxone Pentazocine (50 mg) Tapentadol: Tapentadol (50 mg, 75 mg, 100 mg, 150 m Tramadol: Tramadol (50 mg, 100 mg, 150 mg, 200 mg, Acetaminophen Tramadol (37.5 mg) The HDO Opioid Medications List excludes: Injectables. Opioid cough and cold products.				Oxycodone: Oxycodone (1 MGPML, 5 mg, 7.5 mg, 9 mg, 10 mg, 13.5 mg, 15 mg,
Oxycodone: Oxycodone (1 MGPNL, 5 mg, 7.5 mg, 9 mg, 10 mg, 13.5 mg, 18 mg, 20 mg, 20 MGPNL, 27 mg, 30 mg, 36 mg, 40 mg, 60 mg, 80 mg), Acetaminophen Oxycodone (1 MGPNL, 2.7 mg, 30 mg, 36 mg, 40 mg, 90 mg, 75 mg, 10 mg), Aspirin Oxycodone (4.84 mg), Ibuprofen Oxycodone (5 mg) Oxymorphone: Oxymorphone: Oxymorphone (5 mg, 7.5 mg, 10 mg, 15 mg, 20 mg, 30 mg, 40 mg) Pentazocine: Naloxone Pentazocine (50 mg) Tapentadol: Tapentadol (50 mg, 75 mg, 100 mg, 150 mg, 200 mg, 250 mg) Tramadol: Tramadol (50 mg, 75 mg) The HDO Opioid Medications List excludes: Implement Implement Opioid Cough and cold products. Improvement Improvement Implement	Oxycodone: Oxycodone (1 MGPML, 5 mg, 7.5 mg, 9 mg 18 mg, 20 mg, 20 MGPML, 27 mg, 30 mg, 36 mg, 40 mg Acetaminophen Oxycodone (1 MGPML, 2 MGPML, 2.5 mg), Aspirin Oxycodone (4.84 mg), Ibuprofen Oxycodor Oxymorphone: Oxymorphone (5 mg, 7.5 mg, 10 mg, 15 mg) Pentazocine: Naloxone Pentazocine (50 mg) Tapentadol: Tapentadol (50 mg, 75 mg, 100 mg, 150 mg, Acetaminophen Tramadol (50 mg, 100 mg, 150 mg, Acetaminophen Tramadol (37.5 mg) The HDO Opioid Medications List excludes: Injectables. Opioid cough and cold products.				
Opium: Belladonna Opium (30 mg, 60 mg)Oxycodone: Oxycodone (1 MGPML, 5 mg, 7.5 mg, 9 mg, 10 mg, 13.5 mg, 15 mg, 18 mg, 20 mg, 20 MGPML, 27 mg, 30 mg, 36 mg, 40 mg, 60 mg)Acetaminophen Oxycodone (1 MGPML, 2 MGPML, 2.5 mg, 5 mg, 7.5 mg, 10 mg), Aspirin Oxycodone (4.84 mg), Ibuprofen Oxycodone (5 mg)Oxymorphone: Oxymorphone: Oxymorphone (5 mg, 7.5 mg, 10 mg, 15 mg, 20 mg, 30 mg, 40 mg)Pentazocine: Naloxone Pentazocine: Solomg, 150 mg, 200 mg, 200 mg, 200 mg, 200 mg, 250 mg)Tramadol: Tramadol (50 mg, 100 mg, 150 mg, 200 mg, 250 mg)Tramadol: Tramadol (50 mg, 100 mg, 150 mg, 200 mg, 250 mg)Tramadol (37.5 mg)The HDO Opioid Medications List excludes:Injectables.Opioid cough and cold products.Injectables.Opioid cough and cold products.Injectables.Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.MeasureMeasure DescriptionMeasureCoding	 Opium: Belladonna Opium (30 mg, 60 mg) Oxycodone: Oxycodone (1 MGPML, 5 mg, 7.5 mg, 9 mg 18 mg, 20 mg, 20 MGPML, 27 mg, 30 mg, 36 mg, 40 mg Acetaminophen Oxycodone (1 MGPML, 2 MGPML, 2.5 mg), Aspirin Oxycodone (4.84 mg), Ibuprofen Oxycodor Oxymorphone: Oxymorphone (5 mg, 7.5 mg, 10 mg, 15 mg) Pentazocine: Naloxone Pentazocine (50 mg) Tapentadol: Tapentadol (50 mg, 75 mg, 100 mg, 150 mg Acetaminophen Tramadol (50 mg, 100 mg, 150 mg, 200 mg, Acetaminophen Tramadol (37.5 mg) The HDO Opioid Medications List excludes: Injectables. Opioid cough and cold products. 				
mg, 200 mg), Morphine Naîtrexone (20 mg, 30 mg, 50 mg, 60 mg, 80 mg, 100 mg)Opium: Belladonna Opium (30 ng, 60 mg)Oxycodone: Oxycodone (1 MGPML, 5 mg, 7.5 mg, 9 mg, 10 mg, 13.5 mg, 15 mg, 18 mg, 20 mg, 20 MGPML, 27 mg, 30 mg, 36 mg, 40 mg, 60 mg, 80 mg),Acetaminophen Oxycodone (1 MGPML, 2 MGPML, 2.5 mg, 7.5 mg, 10 mg), Aspirin Oxycodone (4.84 mg), Ibuprofen Oxycodone (5 mg)Oxymorphone: Oxymorphone: Oxymorphone: Oxymorphone: (5 mg), 7.5 mg, 10 mg, 15 mg, 20 mg, 30 mg, 40 mg),Pentazocine: Naloxone Pentazocine (50 mg)Tapentadol: Tapentadol (50 mg, 75 mg, 100 mg, 150 mg, 200 mg, 250 mg)Tramadol (50 mg, 75 mg, 100 mg, 150 mg, 200 mg, 250 mg)Tramadol (50 mg, 100 mg, 150 mg, 200 mg, 250 mg)Tapentadol: Tapentadol (50 mg, 100 mg, 150 mg, 200 mg, 250 mg)Tapentadol: Tapentadol (50 mg, 100 mg, 150 mg, 200 mg, 250 mg)Tapentadol: Tapentadol (50 mg, 100 mg, 150 mg, 200 mg, 250 mg)Tapentadol: Tapentadol (50 mg, 100 mg, 150 mg, 200 mg, 250 mg)Tapentadol: Tapentadol (50 mg, 100 mg, 150 mg, 200 mg, 250 mg)Tapentadol: Tapentadol (50 mg, 100 mg, 150 mg, 200 mg, 250 mg)Tapentadol: Tapentadol (37.5 mg)Opioid cough and cold products.Injectables.Opioid cough and cold products.Insis for inpatient use only and is available only through a restricted program under a Risk Evaluation and Mitigation Strategy (REMS).Metabure bescriptionMeasureMeasure DescriptionMeasureKeasureCoding	mg, 200 mg), Morphine Naltrexone (20 mg, 30 mg, 50 mg) Opium: Belladonna Opium (30 mg, 60 mg) Oxycodone: Oxycodone (1 MGPML, 5 mg, 7.5 mg, 9 mg 18 mg, 20 mg, 20 MGPML, 27 mg, 30 mg, 36 mg, 40 mg Acetaminophen Oxycodone (1 MGPML, 2 MGPML, 2.5 mg), Aspirin Oxycodone (1 MGPML, 2 MGPML, 2.5 mg), Aspirin Oxycodone (4.84 mg), Ibuprofen Oxycodor Oxymorphone: Oxymorphone (5 mg, 7.5 mg, 10 mg, 15 mg) Pentazocine: Naloxone Pentazocine (50 mg) Tapentadol: Tapentadol (50 mg, 75 mg, 100 mg, 150 m Tramadol: Tramadol (50 mg, 75 mg, 100 mg, 150 m Acetaminophen Tramadol (37.5 mg) The HDO Opioid Medications List excludes: Injectables. Opioid cough and cold products.				Morphine: Morphine (2 MGPML, 4 MGPML, 5 mg, 10 mg, 15 mg, 20 MGPML,
Morphine: Morphine (2 MGPML, 5 mg, 10 mg, 15 mg, 20 MGPML, 20 mg, 30 mg, 40 mg, 45 mg, 50 mg, 60 mg, 75 mg, 80 mg, 90 mg, 100 mg, 120 mg, 20 mg, Morphine Naltrexone (20 mg, 30 mg, 60 mg, 50 mg, 60 mg, 80 mg, 100 mg) Opjum: Belladonna Opium (30 mg, 00 mg) Opjum: Belladonna Opium (30 mg, 00 mg) Opium: Belladonna Opium (30 mg, 00 mg, 100 mg, 13.5 mg, 15 mg, 18 mg, 20 mg, 20 MGPML, 27 mg, 30 mg, 40 mg, 40 mg, 40 mg, 60 mg, 80 mg), Acetaminophen Oxycodone (1 MGPML, 2 MGPML, 2.5 mg, 5 mg, 7.5 mg, 10 mg), biprofen Oxycodone (1 MGPML, 2.5 mg, 5 mg, 7.5 mg, 10 mg, 15 mg, 20 mg, 20 mg, 30 mg, 40 mg, 60 mg, 80 mg), Acetaminophen Oxycodone (1 MGPML, 2 MGPML, 2.5 mg, 5 mg, 7.5 mg, 10 mg), biprofen Oxycodone (1 MGPML, 2 MGPML, 2.5 mg, 5 mg, 7.5 mg, 10 mg, 15 mg, 20 mg, 30 mg, 40 mg) Pentazocine: Naloxone Pentazocine (50 mg) Tapentadol: Tapentadol (50 mg, 75 mg, 100 mg, 150 mg, 200 mg, 250 mg) Tremadol: Taramadol (37.5 mg) The HDO Opioid Medications List excludes: Injectables. Opioid cough and cold products. Injectables. Opioid cough and cold products. Methadone for the treatment of opioid use disorder.MeasureMeasure DescriptionMeasureCoding	 Morphine: Morphine (2 MGPML, 4 MGPML, 5 mg, 10 n 20 mg, 30 mg, 40 mg, 45 mg, 50 mg, 60 mg, 75 mg, 80 n mg, 200 mg), Morphine Naltrexone (20 mg, 30 mg, 50 n mg) Opium: Belladonna Opium (30 mg, 60 mg) Oxycodone: Oxycodone (1 MGPML, 5 mg, 7.5 mg, 9 mg 18 mg, 20 mg, 20 MGPML, 27 mg, 30 mg, 36 mg, 40 mg Acetaminophen Oxycodone (1 MGPML, 2 MGPML, 2.5 mg), Aspirin Oxycodone (1 MGPML, 2 MGPML, 2.5 mg), Aspirin Oxycodone (5 mg, 7.5 mg, 10 mg, 15 mg) Pentazocine: Naloxone Pentazocine (50 mg) Tapentadol: Tapentadol (50 mg, 75 mg, 100 mg, 150 m Tramadol: Tapentadol (50 mg, 100 mg, 150 mg, Acetaminophen Tramadol (37.5 mg) The HDO Opioid Medications List excludes: Injectables. Opioid cough and cold products. 				
mg) Morphine: Morphine (2 MGPML, 4 MGPML, 5 mg, 10 mg, 15 mg, 20 MGPML, 20 mg, 30 mg, 40 mg, 40 mg, 40 mg, 45 mg, 50 mg, 60 mg, 75 mg, 80 mg, 90 mg, 100 mg, 120 mg, 200 mg), Morphine Naltrexone (20 mg, 30 mg, 50 mg, 60 mg, 80 mg, 100 mg) gy 0pium: Belladonna Opium (30 mg, 60 mg) Oxycodone: Oxycodone (1 MGPML, 2 mg, 75 mg, 9 mg, 10 mg, 13.5 mg, 15 mg, 20 mg, 20 MGPML, 2 MGPML, 2.5 mg, 5 mg, 0 mg, 10 mg, 13.5 mg, 15 mg, 0 mg, 20 MGPML, 2 MGPML, 2.5 mg, 5 mg, 7.5 mg, 10 mg, 13.5 mg, 15 mg, 0 xymorphone: Oxymorphone (5 mg, 7.5 mg, 10 mg, 13 mg, 20 mg, 30 mg, 36 mg, 40 mg, 60 mg, 80 mg), Acetaminophen Oxycodone (1 MGPML, 2 MGPML, 2.5 mg, 5 mg, 7.5 mg, 10 mg), Aspirin Oxycodone (1 MGPML, 2 MGPML, 2.5 mg, 5 mg, 7.5 mg, 10 mg), Aspirin Oxycodone (5 MGPML, 2 MGPML, 2.5 mg, 5 mg, 7.5 mg, 10 mg), Aspirin Oxycodone (5 Mg, 75 mg, 10 mg, 15 mg, 20 mg, 30 mg, 30 mg, 40 mg) Pentazocine: Naloxone Pentazocine (50 mg) Tapentadol: Tapentadol (50 mg, 75 mg, 10 mg, 150 mg, 200 mg, 300 mg), Acetaminophen Tramadol (50 mg, 7.5 mg, 10 mg, 150 mg, 200 mg, 300 mg), Acetaminophen Tramadol (50 mg, 7.5 mg)The HDO Opioid Medications List excludes: • Injectables. • Methadone for the treatment of opioid use disorder.MeasureMeasure DescriptionMeasureKeasure Information.	mg) Morphine: Morphine (2 MGPML, 4 MGPML, 5 mg, 10 n 20 mg, 30 mg, 40 mg, 45 mg, 50 mg, 60 mg, 75 mg, 80 r mg, 200 mg), Morphine Naltrexone (20 mg, 30 mg, 50 r mg) Opium: Belladonna Opium (30 mg, 60 mg) Oxycodone: Oxycodone (1 MGPML, 5 mg, 7.5 mg, 9 mg 18 mg, 20 mg, 20 MGPML, 27 mg, 30 mg, 36 mg, 40 mg Acetaminophen Oxycodone (1 MGPML, 2 MGPML, 2.5 mg), Aspirin Oxycodone (4.84 mg), Ibuprofen Oxycodor Oxymorphone: Oxymorphone (5 mg, 7.5 mg, 10 mg, 15 mg) Pentazocine: Naloxone Pentazocine (50 mg) Tapentadol: Tapentadol (50 mg, 75 mg, 100 mg, 150 m Tramadol: Tramadol (50 mg, 100 mg, 150 mg, 200 mg, Acetaminophen Tramadol (37.5 mg) The HDO Opioid Medications List excludes: • Injectables. • Opioid cough and cold products.				
Meperidine: Meperidine: Meperidine (10 MGPML, 50 mg, 75mg, 100 mg, 150 mg), Methadone: Methadone (1 MGPML, 2 MGPML, 5 mg, 10 mg, 10 MGPML, 40 mg)Morphine: Morphine (2 MGPML, 4 MGPML, 5 mg, 10 mg, 10 MGPML, 40 mg)Morphine: Morphine (2 MGPML, 4 MGPML, 5 mg, 10 mg, 20 mg, 20 mg, Morphine Maltexone (20 mg, 30 mg, 60 mg, 75 mg, 9 mg, 10 mg, 135 mg, 13 mg, 20 mg, 20 mg/ML, 27 mg, 30 mg, 40 mg, 45 mg, 50 mg, 50 mg, 60 mg, 80 mg).Acetaminophen Oxycodone (1 MGPML, 2 MGPML, 25 mg, 5 mg, 10 mg, 135 mg, 13 mg, 20 mg, 20 mg/ML, 27 mg, 30 mg, 40 mg)Oyrcodone: Oxycodone (1 MGPML, 2 MGPML, 25 mg, 5 mg, 10 mg, 135 mg, 10 mg, 13 mg, 20 mg, 20 mg/ML, 27 mg, 30 mg, 40 mg)Pentarocine: Naloxone Pentazocine (50 mg)Tapentadol: Tapentadol (50 mg, 150 mg, 100 mg, 150 mg, 200 mg, 300 mg), Acetaminophen Tramadol (50 mg, 100 mg, 150 mg, 200 mg, 300 mg), Acetaminophen Tramadol (37.5 mg)The HDO Opioid Medications List excludes: Injectables.Opioid cough and cold products.Injectables.Injectables.Opioid cough and cold products.Iomys¹⁰ (Fentanyl transdermal patch).The HDO Opioid Medications List excludes:Injectables.Methadone for the traatment of opioid use disorder. MeasureMeasure DescriptionMeasureCoding	 Meperidine: Meperidine (10 MGPML, 50 mg, 75mg, 10 Methadone: Methadone (1 MGPML, 2 MGPML, 5 mg, 1 mg) Morphine: Morphine (2 MGPML, 4 MGPML, 5 mg, 10 n 20 mg, 30 mg, 40 mg, 45 mg, 50 mg, 60 mg, 75 mg, 80 r mg, 200 mg), Morphine Naltrexone (20 mg, 30 mg, 50 r mg) Opium: Belladonna Opium (30 mg, 60 mg) Oxycodone: Oxycodone (1 MGPML, 5 mg, 7.5 mg, 9 mg 18 mg, 20 mg, 20 MGPML, 27 mg, 30 mg, 36 mg, 40 mg Acetaminophen Oxycodone (1 MGPML, 2.5 mg), Aspirin Oxycodone (4.84 mg), Ibuprofen Oxycodor Oxymorphone: Oxymorphone (5 mg, 7.5 mg, 10 mg, 15 mg) Pentazocine: Naloxone Pentazocine (50 mg) Tapentadol: Tapentadol (50 mg, 75 mg, 100 mg, 150 m Tramadol: Tramadol (50 mg, 150 mg, 200 mg, Acetaminophen Tramadol (37.5 mg) The HDO Opioid Medications List excludes: Injectables. Opioid cough and cold products. 				
Image: constraint of the second sec	Levorphanol: Levorphanol (2 mg, 3 mg) Meperidine: Meperidine (10 MGPML, 50 mg, 75mg, 10 Methadone: Methadone (1 MGPML, 2 MGPML, 5 mg, 10 m g) Morphine: Morphine (2 MGPML, 4 MGPML, 5 mg, 10 n 20 mg, 30 mg, 40 mg, 45 mg, 50 mg, 60 mg, 75 mg, 80 r mg, 200 mg), Morphine Naltrexone (20 mg, 30 mg, 50 r mg) Opium: Belladonna Opium (30 mg, 60 mg) Oxycodone: Oxycodone (1 MGPML, 2 mg, 30 mg, 36 mg, 40 mg Acetaminophen Oxycodone (1 MGPML, 2 MGPML, 2.5 mg), Aspirin Oxycodone (1 MGPML, 2 MGPML, 2.5 mg), Aspirin Oxycodone (1 MGPML, 2 MGPML, 2.5 mg), Aspirin Oxycodone (5 mg, 7.5 mg, 10 mg, 15 mg) Pentazocine: Naloxone Pentazocine (50 mg) Tapentadol: Tamadol (50 mg, 100 mg, 150 m Tramadol: Tramadol (50 mg, 100 mg, 150 mg, 150 m Tramadol (37.5 mg) The HDO Opioid Medications List excludes: Injectables. Opioid cough and cold products.				Hydromorphone: Hydromorphone (1 MGPML, 2 mg, 3 mg, 4 mg, 8 mg, 12 mg,

Use of Opioids From	The percentage of	Required Exclusions:	Opioid Medications:
Multiple Providers (UOP)	members 18 years and	Members who meet any of the	Benzhydrocodone, Buprenorphine (transdermal patch and buccal film),
	older receiving	following criteria are excluded from	Butorphanol, Codeine, Dihydrocodeine, Fentanyl, Hydrocodone,
	prescription opioids for	the measure:	Hydromorphone, Levorphanol, Meperidine, Methadone, Morphine, Opium,
	≥15 days during the MY	In hospice or using hospice	Oxycodone, Oxymorphone, Pentazocine, Tapentadol, Tramadol
	who received opioids	services any time in the MY.	
	from multiple providers.	• Deceased at any time in the MY.	The UOP Opioid Medications List excludes: Injectables.
	Three rates are		Opioid cough and cold products.
	reported:		• Single-agent and combination buprenorphine products used as part of
	1. Multiple		medication-assisted treatment of opioid use (buprenorphine sublingual
	Prescribers: The		tablets, buprenorphine subcutaneous implant, and all
	proportion of		buprenorphine/naloxone combination products).
	members receiving		 Ionsys[®] (fentanyl transdermal patch), because:
	prescriptions for		 It is only for inpatient use.
	opioids from four or		• It is only available through a restricted program under a Risk Evaluation
	more different		and Mitigation Strategy (REMS).
	prescribers during		Methadone when prescribed for the treatment of opioid use disorder.
	the MY.		
	2. Multiple		
	Pharmacies: The		Note: LOINC and SNOMED codes can be captured through electronic data
	proportion of		submissions. Please contact your Account Executive for more information.
	members receiving		
	prescriptions for		
	opioids from four or		
	more different		
	pharmacies during		
	the MY.		
	3. Multiple		
	Prescribers and		
	Multiple		
	Pharmacies: The		
	proportion of		
	members receiving		
	prescriptions for		
	opioids from four or		
	more different		
	prescribers and four		
	or more different		

	pharmacies during the MY (i.e., the proportion of members who are numerator compliant for both the Multiple Prescribers and Multiple Pharmacies rates). A lower rate indicates better performance for all three rates.		
UTILIZATION			
Measure	Measure Description	Measure Information/Documentation Required	Coding
Well-Child Visits in the First 30 Months of Life (W30)	The percentage of members 15 months – 30 months of age who had the recommended well-child visits with a PCP. Two rates are reported: 1. 6 or more visits on or before the 15- month birthday. 2. 2 or more visits between the 15- month birthday plus 1 day and the 30-month birthday.	Documentation from the medical record must include a note indicating a well visit with a PCP and the date the well-child visit occurred. Well-child/EPSDT visit criteria is based on American Academy of Pediatrics Bright Futures: Guidelines for Health Supervision of Infants, Children and Adolescents. <u>https://www.aap.org/en/practice- management/bright-futures/bright- futures-materials-and-tools/</u> Note : Preventive services may be rendered on visits other than well- child visits. Medical records must include documentation of preventive services. Chronic or acute condition assessment and treatment are excluded from this provision.	Use age-appropriate preventive E&M. Encounter for Well Care: ICD10CM: Z00.00, Z00.01, Z00.110, Z00.111, Z00.121, Z00.129, Z00.2, Z00.3, Z01.411, Z01.419, Z02.5, Z76.1, Z76.2 Well Care Visit: CPT: 99381, 99382, 99383, 99384, 99385, 99391, 99392, 99393, 99394, 99395, 99461 HCPCS: Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

		 Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. Common Chart Deficiencies: Children being seen for sick visits only and no documentation/claims/encounter data related to well visit services provided. 	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Child and Adolescent Well-Care Visits (WCV)	The percentage of members 3 – 21 years of age who had at least one comprehensive well-care visit with a PCP or OB/GYN practitioner during the MY.	Documentation from the medical record must include a note indicating a visit with a PCP or OB/GYN, the date when the well-child visit occurred. Well-child/EPSDT visit criteria is based on American Academy of Pediatrics Bright Futures: Guidelines for Health Supervision of Infants, Children and Adolescents. https://www.aap.org/en/practice- management/bright-futures/bright- futures-materials-and-tools/ Note: Preventive services may be rendered on visits other than well- child visits. Medical records must	Use age-appropriate preventive E&M. Encounter for Well Care: ICD10CM: 200.00, 200.01, 200.110, 200.111, 200.121, 200.129, 200.2, 200.3, 201.411, 201.419, 202.5, 276.1, 276.2 Well Care Visit: CPT: 99381, 99382, 99383, 99384, 99385, 99391, 99392, 99393, 99394, 99395, 99461 Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
		child visits. Medical records must include documentation of preventive services. Chronic or acute condition	

		assessment and treatment are	
		excluded from this provision.	
		 Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased at any time in the MY. 	
		 Common Chart Deficiencies: Children or adolescents being seen for sick visits only and no documentation/claims/encounter data related to well-visit services provided. 	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Annual Dental Visit (ADV)			Requires state-specific measure codes.
Retired by NCQA in MY23 but may still apply in state quality reporting. Consult with your Account Executive.			
MEASURES COLLECTED USI	NG ELECTRONIC CLINICAL D	ATA SYSTEMS	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Follow-Up Care for Children Prescribed ADHD Medication (ADD- E)	The percentage of children 6 – 12 years of age who had a newly prescribed ADHD medication and who had	The Intake Period (IP) is the 12-month window starting 3/1 of the year prior to the MY and ending the last calendar day of 2/MY.	Members are identified through administrative and pharmacy claims. ADHD Medications: CNS Stimulants: Dexmethylphenidate, Dextroamphetamine, Lisdexamfetamine, Methylphenidate, Methamphetamine.

This is a measure at le	least three follow-up		
and Electronic Clinical mon Data Systems. Please whi discuss options for a day direct data feed with your Account Executive. Direct data feeds can improve provider quality Two performance and reduce the burden of medical record requests. follo pravider quality follo 2. Met on t leas visit Pha two with	re visits within a 10- onth period, one of hich was within 30 ys of when the first OHD medication was spensed. vo rates are reported: Initiation Phase: embers who had one low-up visit with actitioner with escribing authority ring the 30 days lowing the IPSD. Continuation Phase: embers who remained the medication for at ast 210 days, had a sit in the Initiation ase, and had at least to follow-up visits thin 270 days after e Initiation Phase ded.	 The Index Prescription Start Date (ISPD) is the earliest prescription dispensing date for an ADHD medication in the IP and where there is a negative medication history. Telephone, telehealth visits are acceptable in both the Initiation and Continuation Phases. Only one of the 2 Continuation Phase visits can be e-visit or virtual check-in. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MY. Deceased in the MY. Acute inpatient encounter or discharge with principal diagnosis of mental, behavioral, or neurodevelopmental disorder. Diagnosis of narcolepsy. Common Chart Deficiencies: Follow-up visit more than 30 days after initial medication dispensed date. 2 additional visits within 9 	Alpha-2 receptor agonists: Clonidine, Guanfacine Miscellaneous ADHD Medications: Atomoxetine Visit Setting Unspecified (with Outpatient POS, Partial Hospitalization POS, Community Mental Health Center POS, or Telehealth POS): CPT: 90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90840, 90845, 90847, 90849, 90853, 90875, 90876, 99221, 99222, 99223, 99231, 99232, 99233, 99238, 99239, 99252, 99253, 99254, 99255 Outpatient POS: 03, 05, 07, 09, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 33, 49, 50, 71, 72 Telehealth POS: 02 BH Outpatient: CPT: 98960, 98961, 98962, 99078, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99243, 99244, 99245, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99411, 99412, 99483 HCPCS: G0463, H0036, H2000, H2017, H2019, H2020, T1015 UBREV: 0510, 0513, 0516, 0517, 0519, 0900, 0901, 0911, 0914, 0915, 0916, 0919 Health and Behavior Assessment or Intervention: CPT: 96156, 96158, 96159, 96164, 96165, 96167, 96168 Partial Hospitalization or Intensive Outpatient: HCPCS: H2001, H2012, S0201, S9480, S9484, S9485
2.			
Tone	iowing the irod.	-	
2.	Continuation	In hospice or using hospice	
Me		-	
-			
		•	
			UBREV 0510 0513 0516 0517 0519 0900 0901 0911 0914 0915 0916
	-		Health and Behavior Assessment or Intervention
		Common Chart Deficiencies	
			CK1: 20120, 20123, 20104, 20105, 20107, 20108
end	ueu.		Partial Harritalization or Intensive Outpatients
			HCPCS: H2001, H2012, S0201, S9480, S9484, S9485
		months of starting medication	Telephone Visit:
		are not documented.	CPT: 99441, 99442, 99443
			Online Assessments: (Continuation Phase One of Two Visits):
			CPT: 99421, 99422, 99423, 99457, 99458
			HCPCS: G2010, G2012, G2251

			Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
Adult Immunization Status (AIS-E)	The percentage of members 19 years of	The Measurement Period (MP) is 1/1 through 12/31.	Immunization Administered:
	age and older who are		Adult Influenza Immunization:
This is a measure collected through claims	up to date on recommended routine	Influenza: Members who received an influenza	CVX: 88, 135, 140, 141, 144, 150, 153, 155, 158, 166, 168, 171, 185, 186, 197, 205
and Electronic Clinical	vaccines for influenza;	vaccine on or between 7/1 of the year	
Data Systems. Please	tetanus and diphtheria	prior to the MP and 6/30 of the MP,	Influenza Virus LAIV Immunization:
discuss options for a	(Td) or tetanus,	or with prior influenza virus vaccine-	CVX : 111, 149
direct data feed with your	diphtheria, and acellular	adverse reaction any time during or	
Account Executive. Direct	pertussis (Tdap); zoster;	before the MP.	Adult Pneumococcal Immunization:
data feeds can improve	and pneumococcal.		Herpes Zoster Recombinant Immunization:
provider quality		Td/Tdap:	CVX: 187
performance and reduce		Members who received at least one	
the burden of medical		Td vaccine or one Tdap vaccine	Td Immunization:
record requests.		between nine years prior to the start	CVX: 09, 113, 115, 138, 139
		of the MP and the end of the MP, or	Tdap Immunization:
		with history of at least one of the	CVX: 115
		following contraindications any time	
		during or before the MP:	Vaccine Procedure:
		Anaphylaxis due to Tdap	Adult Influenza Vaccine Procedure:
		vaccine, anaphylaxis due to	CPT: 90630, 90653, 90654, 90656, 90658, 90661, 90662, 90673, 90674, 90682,
		Td vaccine or its	90686, 90688, 90694, 90756
		components.	
		Encephalopathy due to Tdap	Herpes Zoster Live Vaccine Procedure:
		or Td vaccination (post	CPT: 90736
		tetanus vaccination	Herpes Zoster Recombinant Vaccine Procedure:
		encephalitis, post diphtheria	CPT : 90750
		vaccination encephalitis, or	
		post pertussis vaccination	Influenza Virus LAIV Vaccine Procedure:
		encephalitis).	CPT : 90660, 90672
		Zoster:	Adult Pneumococcal Vaccine Procedure:
		Members who received at least one	CPT: 90670, 90671, 90677, 90732
		dose of the herpes zoster live vaccine	
		or two doses of the herpes zoster	Td Vaccine Procedure:
		recombinant vaccine (at least 28 days	CPT : 90714

		apart) anytime on or after the	Tdap Vaccine Procedure:
		member's 50th birthday, or with prior	CPT : 90715
		adverse reaction caused by zoster	
		vaccine or its components any time	
		during or before the MP.	Note: LOINC and SNOMED codes can be captured through electronic data
			submissions. Please contact your Account Executive for more information.
		Pneumococcal:	
		Members who were administered the	
		23-valent pneumococcal	
		polysaccharide vaccine on or after the	
		member's 60 th birthday before or	
		during the MP, or prior pneumococcal	
		vaccine-adverse reaction any time	
		during or before the MP.	
		Required Exclusions:	
		Members who meet any of the	
		following criteria are excluded from	
		the measure:	
		In hospice or using hospice	
		services any time in the MP.	
		• Deceased at any time in the MP.	
Metabolic Monitoring for	1. The percentage of	Both of the following during the MY.	Members are identified through administrative and pharmacy claims.
Children and Adolescents	children and	At least one test for blood	
on Antipsychotics (APM-	adolescents on	glucose or HbA1c, and	Glucose Lab Test:
E)	antipsychotics who	 At least one test for LDL-C or 	CPT: 80047, 80048, 80053, 80069, 82947, 82950, 82951
	received blood glucose	cholesterol	
This is a measure collected	testing.		HbA1C Lab Test:
through claims and		Required Exclusions:	CPT: 83036, 83037
Electronic Clinical Data	2. The percentage of	Members who meet any of the	
Systems. Please discuss	children and	following criteria are excluded from	HbA1C Test Result or Finding:
options for a direct data	adolescents on	the measure:	CPT-CAT-II: 3044F, 3046F, 3051F, 3052F
feed with your Account	antipsychotics who	In hospice or using hospice	
Executive. Direct data	received cholesterol	services any time in the MY.	Cholesterol Lab Test:
feeds can improve	testing.	Deceased in the MY.	CPT: 82465, 83718, 83722, 84478
provider quality			
performance and reduce	3. The percentage of	Common Chart Deficiencies:	LDL-C Lab Test:
	children and		CPT: 80061, 83700, 83701, 83704, 83721

the burden of medical record requests.	adolescents on antipsychotics who received blood glucose and cholesterol testing	• A1C, LDL-C ordered but not completed.	LDL-C Test Result or Finding: CPT-CAT-II: 3048F, 3049F, 3050F Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
Measure	Measure Description	Measure Information/Documentation Required	Coding
Breast Cancer Screening (BCS-E) This is a measure collected through claims and Electronic Clinical Data Systems. Please discuss options for a direct data feed with your Account Executive. Direct data feeds can improve provider quality performance and reduce the burden of medical record requests	The percentage of members 50 – 74 years of age who were recommended for a routine breast cancer screening and had a mammogram to screen for breast cancer.	 All types and methods of mammograms (screening, diagnostic, film, digital, or digital breast tomosynthesis) qualify for numerator compliance. Note: Biopsies, breast ultrasounds, and MRIs do not count toward this measure. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services in the MP. Deceased at any time in the MP. Receiving palliative care any time in the MP. 66 years of age and older with frailty and advanced illness during the MY. Had gender-affirming chest surgery (CPT code 19318) with a diagnosis of gender dysphoria (Gender Dysphoria Value Set) any time during the member's history through the end of the MP. 	Mammography: CPT: 77061, 77062, 77063, 77065, 77066, 77067 Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

		 Bilateral mastectomy or both right and left unilateral mastectomy with bilateral modifier from same procedure any time during the member's history through the end of the MY. 	
Measure	Measure Description	Measure Information/Documentation	Coding
Colorectal Cancer Screening (COL-E) This is a measure collected through claims and Electronic Clinical Data Systems. Please discuss options for a direct data feed with your Account Executive. Direct data feeds can improve provider quality performance and reduce the burden of medical record requests	The percentage of members 45 – 75 years of age who had appropriate screening for colorectal cancer.	 Required The MY is 1/1 – 12/31. Documentation in the medical record must include a note indicating the date when the colorectal cancer screening was performed. A result is not required if the documentation is clearly part of the "medical history" section of the record; if this is not clear, the result or finding must also be present. (This ensures that the screening was performed and not merely ordered.) Colonoscopy in past 10 years (the MY and 9 years prior). Flexible Sigmoidoscopy in past 5 years (the MY and 4 years prior). CT Colonography in past 5 years (the MY and 4 years prior). Stool DNA (sDNA) with FIT test in past 3 years (the MY and 2 years prior). Fecal Occult Blood Test (FOBT) in the MY. 	Colonoscopy: CPT: 44388, 44389, 44390, 44391, 44392, 44394, 44401, 44402, 44403, 44404, 44405, 44406, 44407, 44408, 45378, 45379, 45380, 45381, 45382, 45384, 45385, 45386, 45388, 45389, 45390, 45391, 45392, 45393, 45398 HCPCS: G0105, G0121 Flexible Sigmoidoscopy: CPT: 45330, 45331, 45332, 45333, 45334, 45335, 45337, 45338, 45340, 45341, 45342, 45346, 45347, 45349, 45350 HCPCS: G0104 CT Colonography: CPT: 74261, 74262, 74263 Stool DNA (sDNA) with Fit Lab Test: CPT: 81528 FOBT Lab test: CPT: 82270, 82274 Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
		Required Exclusions:	

Members who meet any of the	
-	
following criteria are excluded from	
the measure:	
In hospice or using hospice	
services any time in the MY.	
Deceased in the MY.	
 Receiving palliative care any time 	
in the MY.	
 66 years of age and older with 	
frailty and advanced illness	
during the MY.	
Colorectal cancer any time in	
member history through 12/31 of	
the MY.	
 Total colectomy any time in 	
member history through 12/31 of	
the MY.	
Common Chart Deficiencies:	
Member-reported data not	
documented with sufficient	
information to show the	
screening was completed in the	
screening was completed in the measure time frame.	
screening was completed in the measure time frame.Documentation not clear on type	
 screening was completed in the measure time frame. Documentation not clear on type of screening (e.g., only "Col" or 	
 screening was completed in the measure time frame. Documentation not clear on type of screening (e.g., only "Col" or "Colon"). 	
 screening was completed in the measure time frame. Documentation not clear on type of screening (e.g., only "Col" or "Colon"). Documentation not clear on 	
 screening was completed in the measure time frame. Documentation not clear on type of screening (e.g., only "Col" or "Colon"). Documentation not clear on location to which scope advanced 	
 screening was completed in the measure time frame. Documentation not clear on type of screening (e.g., only "Col" or "Colon"). Documentation not clear on location to which scope advanced in situations of incomplete 	
 screening was completed in the measure time frame. Documentation not clear on type of screening (e.g., only "Col" or "Colon"). Documentation not clear on location to which scope advanced in situations of incomplete colonoscopy (must advance to 	
 screening was completed in the measure time frame. Documentation not clear on type of screening (e.g., only "Col" or "Colon"). Documentation not clear on location to which scope advanced in situations of incomplete colonoscopy (must advance to the cecum) or flexible 	
 screening was completed in the measure time frame. Documentation not clear on type of screening (e.g., only "Col" or "Colon"). Documentation not clear on location to which scope advanced in situations of incomplete colonoscopy (must advance to the cecum) or flexible sigmoidoscopy (must advance 	
 screening was completed in the measure time frame. Documentation not clear on type of screening (e.g., only "Col" or "Colon"). Documentation not clear on location to which scope advanced in situations of incomplete colonoscopy (must advance to the cecum) or flexible sigmoidoscopy (must advance into the sigmoid colon). 	
 screening was completed in the measure time frame. Documentation not clear on type of screening (e.g., only "Col" or "Colon"). Documentation not clear on location to which scope advanced in situations of incomplete colonoscopy (must advance to the cecum) or flexible sigmoidoscopy (must advance into the sigmoid colon). Most recent screening dates not 	
 screening was completed in the measure time frame. Documentation not clear on type of screening (e.g., only "Col" or "Colon"). Documentation not clear on location to which scope advanced in situations of incomplete colonoscopy (must advance to the cecum) or flexible sigmoidoscopy (must advance into the sigmoid colon). Most recent screening dates not documented in the 	
 screening was completed in the measure time frame. Documentation not clear on type of screening (e.g., only "Col" or "Colon"). Documentation not clear on location to which scope advanced in situations of incomplete colonoscopy (must advance to the cecum) or flexible sigmoidoscopy (must advance into the sigmoid colon). Most recent screening dates not 	

		 Documentation of only "up to date." Documentation of only "next due" dates. FOBTs performed in an office setting. FOBTs performed on a sample collected via Digital Rectal Exam (DRE). Fewer than 3 samples documented for gFOBT. Documentation not clear if Stool-DNA with FIT or FIT FOBT. 	
Depression Screening and Follow-Up for	The percentage of members 12 years of	The MP is 1/1 through 12/31.	Encounter Performed: Behavioral Health Encounter:
Adolescents and Adults	age and older who were	This measure requires the use of an	CPT: 90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90845,
(DSF-E)	screened for clinical	age-appropriate screening	90846, 90847, 90849, 90853, 90865, 90867, 90868, 90869, 90870, 90875,
	depression using a	instrument. The member's age is used	90876, 90880, 90887, 99484, 99492, 99493
This is a measure	standardized instrument	to select the appropriate depression	HCPCS: G0511, H0004, H0036, H0040, H2000, H2001, H2012, H2015, H2017,
collected through claims	and, if screened	screening instrument.	H2019, H2020, S9484, S9485
and Electronic Clinical	positive, received	Acceptable tools for the	UBREV: 0900, 0901, 0911, 0914, 0915, 0916, 0919,
Data Systems. Please	follow-up care.	Adolescent 12-17 population	
discuss options for a		include PHQ-9; PHQ-9M;	Depression Case Management Encounter:
direct data feed with your	Two rates are reported:	PHQ-2; BDI-FS; CESD-R;	СРТ: 99366
Account Executive. Direct data feeds can improve	1. Depression Screening:	EPDS; PROMIS Depression.Acceptable tools for the	The Hannes Line Articles
provider quality	The percentage of	Adult 18+ population include	Follow-Up Visit:
performance and reduce	members who were	PHQ-9; PHQ-2; BDI-FS; BDI-II;	CPT: 98960, 98961, 98962, 98969, 98970, 98971, 98972, 98980, 98981, 99078, 00202, 00202, 00204, 00205, 00211, 00212, 00212, 00214, 00215, 00242
the burden of medical	screened for clinical	CESD-R; DADS; GDS; EPDS;	99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242,
record requests.	depression using a	M-3; PROMIS Depression,	99243, 99244, 99245, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 00350, 00381, 00382, 00382, 00384, 00385, 00385, 00387, 00301, 00303
	standardized	CUDOS.	99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392,
	instrument.		99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99421,
	2. Follow Up on	Follow up which meets criteria:	99422, 99423, 99441, 99442. 99457, 99458, 99483
	Positive Screen:	Outpatient, telephone, or	HCPCS: G0463, G2010, G2012, T1015
	The percentage of	virtual check-in visit.	UBREV: 0510, 0513, 0516, 0517, 0519
	members who received	Depression case	
	follow-up care on or up	management encounter.	
	to 30 days after the date		

	of the first positive screen.	 A behavioral health encounter. Dispensed antidepressant medication. Additional depression screening on a full-length instrument indicating no depression or no symptoms that require follow up on the same day as a positive screen on a brief screening instrument. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP. Bipolar disorder in the year prior to the MP. Depression that starts during the year prior to the MP. 	Dispensed Antidepressant Medication:Miscellaneous antidepressants: Bupropion, Vilazodone, VortioxetineMonoamine oxidase inhibitors: Isocarboxazid, Phenelzine, Selegiline,TranylcyprominePhenylpiperazine antidepressants: Nefazodone, TrazodonePsychotherapeutic combinations: Amitriptyline-chlordiazepoxide,Amitriptyline-perphenazine, Fluoxetine-olanzapineSNRI antidepressants: Desvenlafaxine, Duloxetine, Levomilnacipran,VenlafaxineSSRI antidepressants: Citalopram, Escitalopram, Fluoxetine, Fluvoxamine,Paroxetine, SertralineTetracyclic antidepressants: Maprotiline, MirtazapineTricyclic antidepressants: Amitriptyline, Amoxapine, Clomipramine,Desipramine, Doxepin (>6mg), Imipramine, Nortriptyline, Protriptyline,TrimipramineNote: LOINC and SNOMED codes can be captured through electronic datasubmissions. Please contact your Account Executive for more information.
Measure	Measure Description	Measure Information/Documentation Required	Coding
Utilization of the PHQ-9 to Monitor Depression Symptoms for Adolescents and Adults (DMS-E) This is a measure collected through claims and Electronic Clinical Data Systems. Please	The percentage of members 12 years of age and older with a diagnosis of major depression or dysthymia who had an outpatient encounter with a PHQ-9 score present in their record in the same	 The Measurement Periods (MP) are: January 1 through April 30. May 1 through August 31. September 1 through December 31. The PHQ-9 assessment does not need to occur during a face-to-face encounter; phone-based, e-visit,	Diagnosis: Major Depression or Dysthymia: ICD10CM F32.0, F32.1, F32.2, F32.3, F32.4, F32.5, F32.9, F33.0, F33.1, F33.2, F33.3, F33.40, F33.41, F33.42, F33.9, F34.1 Encounter Performed: Interactive Outpatient Encounter: CPT: 90791, 90792, 90832, 90834, 90837, 98960, 98961, 98962, 98980, 98981, 99078, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99243, 99244, 99245, 99341, 99342, 99344, 99345, 99347, 99348,

discuss options for a direct data feed with your Account Executive. Direct data feeds can improve provider quality performance and reduce the burden of medical record requests. assessment period as the encounter. virtual check-in, or electronic secure messaging is acceptable. 99329, 99330, 99381, 99382, 99383, 99382, 99383, 99340, 99402, 99403, 99402, 99403, 99402, 99403, 99402, 99403, 99402, 99403, 99402, 99403, 99404, 99402, 99403, 99404, 99402, 99403, 99404, 99457, 99457, 99458, 99465, 19004, H003E, H0004, H003E, H0004, H003E, H0004, H003E, H0004, H002B, H0004, H001E, Galda Gillar, Galda			
 For adolescents, guidelines recommend systematic and regular tracking of treatment goals and outcomes, including assessing depressive symptoms. The PHQ-9 tool assesses the nine DSM, Fourth Edition, Text Revision (DSM-IV-TR) criterion symptoms and effects on functioning and has been shown to be highly accurate in diagnosing patients with persistent major depression, partial remission, and full remission. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP. 	direct data feed with your Account Executive. Direct data feeds can improve provider quality performance and reduce the burden of medical	messaging is acceptable. Note: Standardized instruments are useful in identifying meaningful change in clinical outcomes over time. Guidelines for adults recommend that providers establish and maintain regular follow-up with patients diagnosed with depression and use a	99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99421, 99422, 99423, 99441, 99442, 99443, 99457,99458, 99483 HCPCS: G0463, G2010, G2012, G2251, H0004, H0036, H0040, H2000, H2010, H2012, H2017, H2019, H2020, S9484, S9485, T1015 UBREV: 0510, 0513, 0516, 0517, 0519, 0900, 0901, 0911, 0914, 0915, 0916, 0919 Note: LOINC and SNOMED codes can be captured through electronic data
regular tracking of treatment goals and outcomes, including assessing depressive symptoms. The PHQ-9 tool assesses the nine DSM, Fourth Edition, Text Revision (DSM-IV-TR) criterion symptoms and effects on functioning and has been show to be highly accurate in diagnosing patients with persistent major depression, partial remission, and full remission. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP.			
goals and outcomes, including assessing depressive symptoms. The PHQ-9 tool assesses the nine DSM, Fourth Edition, Text Revision (DSM-IV-TR) criterion symptoms and effects on functioning and has been shown to be highly accurate in diagnosing patients with persistent major depression, partial remission, and full remission. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP. 			
depressive symptoms. • The PHQ-9 tool assesses the nine DSM, Fourth Edition, Text Revision (DSM-IV-TR) criterion symptoms and effects on functioning and has been shown to be highly accurate in diagnosing patients with persistent major depression, partial remission, and full remission. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: • In hospice or using hospice services any time in the MP. • Decreased at any time in the MP.		goals and outcomes,	
 The PHQ-9 tool assesses the nine DSM, Fourth Edition, Text Revision (DSM-IV-TR) criterion symptoms and effects on functioning and has been shown to be highly accurate in diagnosing patients with persistent major depression, partial remission, and full remission. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP. 			
nine DSM, Fourth Edition, Text Revision (DSM-IV-TR) criterion symptoms and effects on functioning and has been shown to be highly accurate in diagnosing patients with persistent major depression, partial remission, and full remission. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP.			
Text Revision (DSM-IV-TR) criterion symptoms and effects on functioning and has been shown to be highly accurate in diagnosing patients with persistent major depression, partial remission, and full remission. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP.			
effects on functioning and has been shown to be highly accurate in diagnosing patients with persistent major depression, partial remission, and full remission. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP.			
has been shown to be highly accurate in diagnosing patients with persistent major depression, partial remission, and full remission. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP.			
accurate in diagnosing patients with persistent major depression, partial remission, and full remission. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP.		•	
patients with persistent major depression, partial remission, and full remission. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP.			
major depression, partial remission, and full remission. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP. 			
remission, and full remission. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP.			
Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP.			
Members who meet any of the following criteria are excluded from the measure: • In hospice or using hospice services any time in the MP. • Deceased at any time in the MP.			
following criteria are excluded from the measure: • In hospice or using hospice services any time in the MP. • Deceased at any time in the MP.		Required Exclusions:	
 the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP. 		-	
 In hospice or using hospice services any time in the MP. Deceased at any time in the MP. 		-	
 services any time in the MP. Deceased at any time in the MP. MP. 			
Deceased at any time in the MP.			
MP.		-	
		-	
		Bipolar disorder in the MP.	

		 Personality disorder in the MP. Psychotic disorder in the MP. Pervasive development disorder in the MP. 	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Depression Remission or Response for Adolescents and Adults (DRR-E)	The percentage of members 12 years of age and older with a diagnosis of depression	The Measurement Period (MP) is 1/1 through 12/31. The Intake Period (IP) is 5/1 of the	Diagnosis: Major Depression or Dysthymia: ICD10CM: F32.0, F32.1, F32.2, F32.3, F32.4, F32.5, F32.9, F33.0, F33.1, F33.2, F33.3, F33.40, F33.41, F33.42, F33.9, F34.1
This is a measure collected through claims and Electronic Clinical	and an elevated PHQ-9 score who had evidence of response or remission	year prior to the MP through 4/30 of the MP.	Encounter Performed: Interactive Outpatient Encounter:
Data Systems. Please discuss options for a direct data feed with your Account Executive. Direct	within 120-240 days (4 – 8 months) of the elevated score.	The Episode Intake Start Date (EISD) is the earliest date in the IP where a member has a diagnosis of major depression or dysthymia and a PHQ-9	CPT: 90791, 90792, 90832, 90834, 90837, 98960, 98961, 98962, 98980, 98981, 99078, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99243, 99244, 99245, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391,
data feeds can improve provider quality performance and reduce	Three rates are reported: 1. Follow-Up PHQ-9:	total score >9 documented. Required Exclusions:	99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99421, 99422, 99423, 99441, 99442, 99443, 99457,99458, 99483
the burden of medical record requests.	The percentage of members who have a follow-up PHQ-9 score	Members who meet any of the following criteria during the IP or during the MP are excluded from the	HCPCS: G0463, G2010, G2012, G2251, H0004, H0036, H0040, H2000, H2010, H2012, H2017, H2019, H2020, S9484, S9485, T1015
	documented within 120- 240 days (4 – 8 months) after the initial elevated PHQ-9 score.	 measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP. 	UBREV: 0510, 0513, 0516, 0517, 0519, 0900, 0901, 0911, 0914, 0915, 0916, 0919
	2. Depression Remission: The percentage of members who achieved	 Bipolar disorder. Personality disorder. Psychotic disorder. Pervasive development disorder. 	Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
	remission within 120- 240 days (4 – 8 months) after the initial elevated PHQ-9 score.		

Measure	 3. Depression Response: The percentage of members who showed response within 120-240 days (4 – 8 months) after the initial elevated PHQ-9 score. Measure Description 	Measure	Coding
incusure		Information/Documentation Required	
Unhealthy Alcohol Use Screening and Follow-Up (ASF-E) This is a measure collected through claims and Electronic Clinical Data Systems. Please discuss options for a direct data feed with your Account Executive. Direct data feeds can improve provider quality performance and reduce the burden of medical record requests.	The percentage of members 18 years of age and older who were screened for unhealthy alcohol use using a standardized instrument and, if screened positive, received appropriate follow-up care. Two rates are reported: 1. Unhealthy Alcohol Use Screening: The percentage of members who had a systematic screening for unhealthy alcohol use. 2. Alcohol Counseling or Other Follow-up Care: The percentage of members receiving brief counseling or other follow-up care within 60 days (2 months) of	 The Measurement Period (MP) is 1/1 through 12/31. Follow-up is an encounter on, or up to 60 days after, the date of the first positive screening that includes at least one of the following: Feedback on alcohol use and harms. Identification of high-risk situations for drinking and coping strategies. Increase the motivation to reduce drinking. Development of a personal plan to reduce drinking. Documentation of receiving alcohol misuse treatment. Required Exclusions: Members who meet any of the following criteria during the MP are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP. 	Diagnosis Alcohol Use Disorder: ICD10CM: F10.10, F10.120, F10.121, F10.129, F10.130, F10.131, F10.132, F10.139, F10.14, F10.150, F10.151, F10.159, F10.180, F10.181, F10.182, F10.188, F10.19, F10.20, F10.220, F10.221, F10.229, F10.230, F10.231, F10.232, F10.239, F10.24, F10.250, F10.251, F10.259, F10.26, F10.27, F10.280, F10.281, F10.982, F10.939, F10.90, F10.900, F10.921, F10.929, F10.930, F10.931, F10.932, F10.939, F10.94, F10.950, F10.951, F10.959, F10.96, F10.97, F10.980, F10.981, F10.982, F10.988, F10.99, K29.20, K29.21, K70.10, K70.11 Intervention Performed: Alcohol Counseling or Other Follow-Up Care: CPT: HCPCS: G0396, G0397, G2011, H0005, H0015, H2036 Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

	screening positive for unhealthy alcohol use.	 Alcohol use disorder that starts during the year prior to the MP. History of dementia any time during the member's history through the end of the MP. 	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Prenatal Immunization Status (PRS-E) This is a measure collected through claims and Electronic Clinical Data Systems. Please discuss options for a direct data feed with your Account Executive. Direct data feeds can improve provider quality performance and reduce the burden of medical record requests.	The percentage of deliveries in which the member received influenza and tetanus, diphtheria toxoids, and acellular pertussis (Tdap) vaccinations.	 The Measurement Period (MP) is 1/1 through 12/31. Influenza: Deliveries where members received an adult influenza vaccine on or between July 1 of the year prior to the MP and the delivery date; or Deliveries where members had an influenza virus vaccine- adverse reaction any time during or before the MP. Tdap: Deliveries where the members had any of the following: At least one Tdap vaccine during the pregnancy (including the delivery date). Anaphylactic reaction to Tdap or Td vaccine or its components any 	Immunization Administered: Adult Influenza Immunization: CVX: 88, 135, 140, 141, 144, 150, 153, 155, 158, 166, 168, 171, 185, 186, 197, 205 Tdap Immunization: CVX: 115 Vaccine Procedure: Adult Influenza Vaccine Procedure: CPT: 90630, 90653, 90654, 90656, 90658, 90661, 90662, 90673, 90674, 90682, 90686, 90688, 90694, 90756 Tdap Vaccine Procedure: CPT: 90715 Deliveries: CPT: 59400, 59409, 59410, 59514, 59515, 59612, 59614, 59620, 59622 HCPCS: 10D0020, 10D0021, 10D0022, 10D0723, 10D0724, 10D0725, 10D0726, 10D0727, 10D0728, 10E0XZZ
		 time during or before the MP. Encephalopathy due to Td or Tdap vaccination any time during or before the MP. A note indicating the specific antigen name and the immunization date, or an immunization certificate prepared 	Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.

		 by a healthcare provider that has the dates of administration. Documented history of specific disease, anaphylactic reactions, or contraindications for a specific vaccine. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP. Delivered at less than 37 weeks gestation. 	
Measure	Measure Description	Measure Information/Documentation Required	Coding
Prenatal Depression	The percentage of	The Measurement Period (MP) is 1/1	Encounter Performed:
Screening and Follow-Up	deliveries in which	- 12/31.	Behavioral Health Encounter:
(PND-E)	members were screened		CPT: 90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90845,
	for clinical depression	This measure requires the use of an	90846, 90847, 90849, 90853, 90865, 90867, 90868, 90869, 90870, 90875,
This is a measure	while pregnant and, if	age-appropriate screening	90876, 90880, 90887, 99484, 99492, 99493
collected through claims	screened positive,	instrument. The member's age is used	
and Electronic Clinical	received follow-up care.	to select the appropriate depression	HCPCS: G0511, H0004, H0036, H0040, H2000, H2001, H2012, H2015, H2017,
Data Systems. Please		screening instrument.	H2019, H2020, S9484, S9485
discuss options for a	Two rates are reported:	Acceptable tools for the	
direct data feed with your	1. Depression	Adolescent 12-17 population	UBREV: 0900, 0901, 0911, 0914, 0915, 0916 , 0919
Account Executive. Direct	Screening:	include PHQ-9; PHQ-9M; PHQ-2;	
aata tooda can improvo	The percentage of	BDI-FS; CESD-R; EPDS; PROMIS	Depression Case Management Encounter:
data feeds can improve			ADT 000.00
provider quality	deliveries in which	Depression.	СРТ: 99366
provider quality performance and reduce	deliveries in which members were screened	Depression.Acceptable tools for the Adult	
provider quality performance and reduce the burden of medical	deliveries in which members were screened for clinical depression	 Depression. Acceptable tools for the Adult 18+ population include PHQ-9; 	Follow-Up Visit:
provider quality performance and reduce	deliveries in which members were screened for clinical depression using a standardized	 Depression. Acceptable tools for the Adult 18+ population include PHQ-9; PHQ-2; BDI-FS; BDI-II; CESD-R; 	Follow-Up Visit: CPT: 98960, 98961, 98962, 98970, 98971, 98972, 98980, 98981, 99078, 99202,
provider quality performance and reduce the burden of medical	deliveries in which members were screened for clinical depression	 Depression. Acceptable tools for the Adult 18+ population include PHQ-9; 	Follow-Up Visit:

2. Follow up on Positive Screen: The percentage of deliveries in which members received follow-up care within 30 days of screening positive for depression.	 Follow up which meets criteria: Outpatient, telephone, or virtual check-in visit. Depression case management encounter. A behavioral health encounter. Dispensed antidepressant medication. Additional depression screening on a full-length instrument indicating no depression or no symptoms that require follow up on the same day as a positive screen on a brief screening instrument. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP. Delivered at less than 37 weeks gestation. 	 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99421, 99422, 99423, 99441, 99442, 99457, 99458, 99483 HCPCS: G0463, G2010, G2012, T1015 UBREV: 0510, 0513, 0516, 0517, 0519 Dispensed Antidepressant Medication: Miscellaneous antidepressants: Bupropion, Vilazodone, Vortioxetine Monoamine oxidase inhibitors: Isocarboxazid, Phenelzine, Selegiline, Tranylcypromine Phenylpiperazine antidepressants: Nefazodone, Trazodone Psychotherapeutic combinations: Amitriptyline-chlordiazepoxide, Amitriptyline-perphenazine, Fluoxetine-olanzapine SNRI antidepressants: Desvenlafaxine, Duloxetine, Levomilnacipran, Venlafaxine SSRI antidepressants: Maprotiline, Mirtazapine Tricyclic antidepressants: Amitriptyline, Amoxapine, Clomipramine, Desipramine, Doxepin (>6mg), Imipramine, Nortriptyline, Protriptyline, Trimipramine Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
Measure Description	Measure Information/Documentation Required	Coding
The percentage of deliveries in which members were screened for clinical depression during the postpartum period, and if screened positive, received follow-up care.	The Measurement Period (MP) is 1/1 – 12/31. This measure requires the use of an age-appropriate screening instrument. The member's age is used to select the appropriate depression screening instrument.	Encounter Performed: Behavioral Health Encounter: CPT: 90791, 90792, 90832, 90833, 90834, 90836, 90837, 90838, 90839, 90845, 90846, 90847, 90849, 90853, 90865, 90867, 90868, 90869, 90870, 90875, 90876, 90880, 90887, 99484, 99492, 99493 HCPCS: G0511, H0004, H0036, H0040, H2000, H2001, H2012, H2015, H2017, H2019, H2020, S9484, S9485
	Positive Screen: The percentage of deliveries in which members received follow-up care within 30 days of screening positive for depression. Measure Description The percentage of deliveries in which members were screened for clinical depression during the postpartum period, and if screened positive, received	Positive Screen: The percentage of deliveries in which members received follow-up care within 30 days of screening positive for depression.• Outpatient, telephone, or virtual check-in visit.• Depression case management encounter.• Depression case management encounter.• A behavioral health encounter.• Dispensed antidepressant medication.• Additional depression screening on a full-length instrument indicating no depression or no symptoms that require follow up on the same day as a positive screen on a brief screening instrument.Required Exclusions: Members who meet any of the following criteria are excluded from the measure:• In hospice or using hospice services any time in the MP.• Delivered at less than 37 weeks gestation.Measure DescriptionMeasure Information/Documentation RequiredThe percentage of deliveries in which members were screened for clinical depression during the postpartum period, and if screened positive, receivedMeasure requires the use of an age-appropriate screening instrument. The member's age is used to select the appropriate depression

discuss options for a direct data feed with your Account Executive. Direct data feeds can improve provider quality performance and reduce the burden of medical record requests.	Two rates are reported. 1. Depression Screening: The percentage of deliveries in which members were screened for clinical depression using a standardized instrument during the postpartum period. 2. Follow up on Positive Screen: The percentage of deliveries in which members received follow-up care within 30 days of screening positive for depression.	 Acceptable tools for the Adolescent 12-17 population include PHQ-9; PHQ-9M; PHQ-2; BDI-FS; CESD-R; EPDS; PROMIS Depression. Acceptable tools for the Adult 18+ population include PHQ-9; PHQ-2; BDI-FS; BDI-II; CESD-R; DADS; EPDS; M-3; PROMIS Depression, CUDOS. Follow up which meets criteria: Outpatient, telephone, or virtual check-in visit. Depression case management encounter. A behavioral health encounter. Dispensed antidepressant medication. Additional depression screening on a full-length instrument indicating no depression or no symptoms that require follow up on the same day as a positive screen on a brief screening instrument. Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP. 	UBREV: 0900, 0901, 0911, 0914, 0915, 0916, 0919 Depression Case Management Encounter: CPT: 99366 Follow-Up Visit: CPT: : 98960, 98961, 98962, 98970, 98971, 98972, 98980, 98981, 99078, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99242, 99243, 99244, 99245, 99341, 99342, 99344, 99345, 99347, 99348, 99394, 99350, 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397, 99401, 99402, 99403, 99404, 99421, 99422, 99423, 99441, 99442, 99457, 99458, 99483 HCPCS: G0463, G2010, G2012, T1015 UBREV: 0510, 0513, 0516, 0517, 0519 Dispensed Antidepressant Medication: Miscellaneous antidepressants: Bupropion, Vilazodone, Vortioxetine Monoamine oxidase inhibitors: Isocarboxazid, Phenelzine, Selegiline, Tranylcypromine Phenylpiperazine antidepressants: Nefazodone, Trazodone Psychotherapeutic combinations: Amitriptyline-chlordiazepoxide, Amitriptyline-perphenazine, Fluoxetine-olanzapine SNRI antidepressants: Desvenlafaxine, Duloxetine, Levomilnacipran, Venlafaxine SSRI antidepressants: Citalopram, Escitalopram, Fluoxetine, Fluvoxamine, Paroxetine, Sertraline Tetracyclic antidepressants: Maprotiline, Mirtazapine Tricyclic antidepressants: Maprotiline, Mirtazapine Tricyclic antidepressants: Maprotiline, Mirtazapine Tricyclic antidepressants: Maprotiline, Mirtazapine Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
---	---	---	---

Measure	Measure Description	Measure Information/Documentation Required	Coding
Social Need Screening and Intervention (SNS-E) This is a measure	The percentage of members who were screened, using pre- specified instruments, at least once in the	Accountable Health	Food Intervention: CPT: 96156, 96160, 96161, 97802, 97803, 97804 HCPCS: S9470 Homelessness/Housing Intervention:
collected through claims and Electronic Clinical Data Systems. Please discuss options for a direct data feed with your Account Executive. Direct	measurement period (MP) for unmet food, housing, and transportation needs and received a corresponding	Communities (AHC) Health- Related Social Needs (HRSN) Screening Tool. • American Academy of Family Physicians (AAFP) Social Needs Screening Tool.	CPT: 96156, 96160, 96161 Transportation Interventions: CPT: 96156, 96160, 96161
data feeds can improve provider quality performance and reduce the burden of medical record requests.	 intervention within 30 days of screening positive. Food Screening: The percentage of members who were screened for food insecurity. Food Intervention: The percentage of members who received a corresponding intervention within 30 days (1 month) of screening positive for food insecurity. Housing Screening: The 	 Health Leads Screening Panel. Hunger Vital Sign (HVS). Protocol for Responding to and Assessing Patients' Assets, Risks & Experiences (PRAPARE). Safe Environment for Every Kid (SEEK). U.S. Household Food Security Survey. U.S. Adult Food Security Survey. U.S. Child Food Security Survey. U.S. Household Food Security Survey – Six-Item Short Form. 	Note: LOINC and SNOMED codes can be captured through electronic data submissions. Please contact your Account Executive for more information.
	percentage of members who were screened for housing instability, homelessness or housing inadequacy.	 We Care Survey. WellRx Questionnaire. Housing Instability, homelessness, and housing inadequacy: Accountable Health Communities (AHC) Health- 	

The measureme	· · · · · · · · · · · · · · · · · · ·	
period (MP) is 1,	/1 – Screening Tool.	
12/31.	American Academy of Family	
	Physicians (AAFP) Social	
	Needs Screening Tool.	
	Children's HealthWatch	
	Housing Stability Vital Signs.	
	Health Leads Screening Panel	
	 Protocol for Responding to 	
	and Assessing Patients'	
	Assets, Risks & Experiences	
	(PRAPARE).	
	We Care Survey.	
	WellRx Questionnaire.	
	Transportation insecurity:	
	Accountable Health	
	Communities (AHC) Health-	
	Related Social Needs (HRSN)	
	Screening Tool.	
	American Academy of Family	
	Physicians (AAFP) Social	
	Needs Screening Tool.	
	Comprehensive Universal	
	Behavior Screen (CUBS).	
	Health Leads Screening	
	Panel.	
	 Protocol for Responding to 	
	and Assessing Patients'	
	Assets, Risks & Experiences	
	(PRAPARE).	
	PROMIS	
	WellRx Questionnaire	
	Interventions are required for any	
	element (food, housing, and	
	transportation) found positive upon	
	screening. Interventions must	
	correspond to the positive screening	
		L/

and must be within 30 days of positive screen (day of screen and 30
days following for a total of 31 days. Interventions include:
Assistance.
• Assessment.
Counseling.
Coordination.
Education.
Evaluation of Eligibility.
Provision.
Referral.
Required Exclusions: Members who meet any of the following criteria are excluded from the measure: In hospice or using hospice services any time in the MP. Deceased at any time in the MP.
Deceased at any time in the MP.



All images are used under license for illustrative purposes only. Any individual depicted is a model.